

Review

VAR Models with an Index Structure: A Survey with New Results

Gianluca Cubadda 

Dipartimento di Economia e Finanza, Università di Roma “Tor Vergata”, Via Columbia 2, 00133 Roma, Italy;
gianluca.cubadda@uniroma2.it

Abstract

The main aim of this paper is to review recent advances in the multivariate autoregressive index model [MAI] and their applications to economic and financial time series. MAI has recently gained momentum because it can be seen as a link between two popular but distinct multivariate time series approaches: vector autoregressive modeling [VAR] and the dynamic factor model [DFM]. Indeed, on the one hand, MAI is a VAR model with a peculiar reduced-rank structure that can lead to a significant dimension reduction; on the other hand, it allows for the identification of common components and common shocks in a similar way as the DFM. Our focus is on recent developments of the MAI, which include extending the original model with individual autoregressive structures, stochastic volatility, time-varying parameters, high-dimensionality, and co-integration. In addition, some gaps in the literature are filled by providing new results on the representation theory underlying previous contributions, and a novel model is provided.

Keywords: multivariate autoregressive index models; vector autoregressive models; dynamic factor models; reduced-rank regression

JEL Classification: C32



Academic Editor: Julien Chevallier

Received: 31 August 2025

Revised: 29 September 2025

Accepted: 16 October 2025

Published: 22 October 2025

Citation: Cubadda, G. (2025). VAR Models with an Index Structure: A Survey with New Results. *Econometrics*, 13(4), 40. <https://doi.org/10.3390/econometrics13040040>

Copyright: © 2025 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The vector autoregressive model [VAR] and the dynamic factor model [DFM] are arguably among the most popular tools for analyzing economic and financial variables over time. Since the seminal contribution of Sims (1980), VARs have been theoretically extended and practically implemented to forecast, structurally analyze, and detect comovements in multivariate time series. DFMs were introduced more recently (Forni et al., 2000; Forni & Lippi, 2001; Stock & Watson, 2002a, 2002b; Bai & Ng, 2002; Bai, 2003), but they rapidly contested the role of the workhorse in empirical macroeconomics.

The main reason for the success of the DFM is two-fold. First, it allows handling a much larger number of variables than those that are generally employed in traditional small-scale VARs, thus potentially boosting forecasting accuracy and solving the informational deficiency problems that arise in structural analyses when the agent's information set is richer than the econometrician's information set (see, e.g., Forni & Gambetti, 2014). Second, DFM allows for disentangling the shocks that drive the common component of high-dimensional time series and recovering the structural shocks from these common shocks only. Hence, in structural DFMs, the number of shocks is smaller than the number of variables (Forni et al., 2009), which is in line with dynamic, stochastic general equilibrium models (see Fernández-Villaverde et al., 2016 and the references therein) and, more

generally, with the standard macroeconomic view that a low number of shocks drives aggregate fluctuations.

Efforts have recently been made to endow the VAR with the above-mentioned features of the DFM. On the one hand, shrinkage estimators have been proposed for medium–large VARs, both from a Bayesian perspective (e.g., [Bańbura et al., 2010](#); [Koop, 2013](#); [Carriero et al., 2015](#)) and from a classical standpoint (e.g., [Hsu et al., 2008](#); [Kock & Callot, 2015](#); [Hecq et al., 2023](#)). On the other hand, the multivariate autoregressive index (MAI) model—originally proposed by [Reinsel \(1983\)](#) as a convenient approach to dimension reduction in stationary VARs—has recently gained renewed attention.¹

Late advances have shown that MAI and its variants allow for both forecasting variables and identifying shocks analogously to the DFM but without encountering some issues in model identification and statistical inference that characterize the latter, such as the requirement that the number of variables diverges at a given rate and the need for specific assumptions on both the correlation structure of the idiosyncratic components and the factor loadings (see, i.e., [Bai, 2003](#); [Bai & Ng, 2006](#)). Moreover, VARs with index structures have been shown to be able to accommodate features such as stochastic volatility ([Carriero et al., 2022](#)) and time-varying parameters ([Cubadda et al., 2025](#)), which are not easy to handle within the DFM framework. The MAI falls within reduced-rank VARs, a general class of models that include, as special cases, both the cointegrated VAR (see [Johansen, 1995](#) and the references therein) and the common serial correlation [CSC] models (see [Cubadda & Hecq, 2022b](#) and the references therein). Although CSC models and MAI have similar mathematical formulations, their respective goals and properties are rather different; whereas the former are based on the existence of (possibly dynamic) linear combinations of autocorrelated time series that are white noise, VARs with an index structure assume that there is a limited number of channels through which information from the past is transmitted to the variables of interest. The main aim of this paper is twofold. First, recent developments in MAI are reviewed, such as the structural MAI ([Carriero et al., 2016](#)), the vector heterogeneous index model for realized volatilities ([Cubadda et al., 2017](#)), and augmentations of the original model with individual autoregressive structures ([Cubadda & Guardabascio, 2019](#)), stochastic volatility ([Carriero et al., 2022](#)), time-varying parameters ([Cubadda et al., 2025](#)), high-dimensionality ([Cubadda & Hecq, 2022a](#)), and cointegration ([Cubadda & Mazzali, 2024](#)). Second, new results are provided in terms of representation theory for various models, and a novel modeling is proposed, namely the cointegrated index-augmented autoregressive model, which combines and extends the results in [Cubadda and Guardabascio \(2019\)](#) and [Cubadda and Mazzali \(2024\)](#).

This paper is organized as follows. Focusing on representation theory, Section 2 reviews previous contributions and provides new insights into some of them. Section 3 presents the new model and deals with its estimation, whereas some details of the estimation procedure are relegated to Appendix A. Finally, Section 4 provides some conclusions.

2. VAR Models with an Index Structure

In this section, we review models that are rooted in the original MAI formulation and provide new results with respect to representation theory for some of them. Analogies and differences with the DFM are discussed in detail. Estimation and identification issues are also covered.

2.1. The Structural Multivariate Autoregressive Index Model

Let us assume that the n -vector time series $Y_t = (y_{1t}, \dots, y_{nt})'$ is generated by the following stationary VAR(p) model:

$$\Phi(L)Y_t = \varepsilon_t, \quad t = 1 \dots T, \quad (1)$$

where L is the lag operator, $\Phi(L) = I_n - \sum_{j=1}^p \Phi_j L^j$; ε_t is a vector of n errors with $E(\varepsilon_t \varepsilon_t') = \Sigma$ (positive definite) and finite fourth moments, $E(\varepsilon_t | F_{t-1}) = 0$; and F_t is the natural filtration of the process Y_t . For simplicity, deterministic elements are ignored.

The key assumption of MAI (Reinsel, 1983) is the following:

Assumption 1. *The following holds:*

$$[\Phi_1', \dots, \Phi_p']' = [\alpha_1', \dots, \alpha_p']' \omega',$$

where ω is a full-rank $n \times q$ -matrix with $q < n$, and α_j is a $n \times q$ -matrix for $j = 1, \dots, p$.

Under Assumption 1, Model (1) can be rewritten as

$$Y_t = \sum_{j=1}^p \alpha_j \underbrace{\omega' Y_{t-j}}_{f_{t-j}} + \varepsilon_t \quad (2)$$

where linear combinations of $f_t = \omega' Y_t$ are called indexes. The MAI has at most $nq(p+1) - q^2$ mean parameters, which implies a significant dimension reduction when p is small with respect to n .²

By premultiplying both sides of Equation (2) with ω' , we get

$$f_t = \sum_{j=1}^p \omega' \alpha_j f_{t-j} + \omega' \varepsilon_t, \quad (3)$$

which shows that the indexes follow a VAR(p) process and not a VARMA process, as is generally the case for linear combinations of elements of a VAR (see Cubadda et al., 2009 and the references therein).

Remark 1. *In view of Equations (2) and (3), the MAI resembles the exact DFM [EDFM] (see Lippi, 2019 and the references therein), but there are also some relevant differences. First, in the EDFM series, Y_t loads the factors contemporaneously and not only with lags. Second, the factors and the idiosyncratic terms in the EDFM are uncorrelated at any lag lead, whereas in the MAI, we have $E(f_t \varepsilon_{t+j}) = 0$ only for $j > 0$. Third, the contemporaneous variance matrix of the idiosyncratic terms in the EDFM is diagonal, whereas Σ is generally not diagonal.*

Placing emphasis on the analogies between MAI and EDFM, Carriero et al. (2016) propose identifying structural shocks as linear transformations of the index shocks only. Starting from the Wold representation of series Y_t

$$Y_t = \Psi(L)\varepsilon_t,$$

and inserting the decomposition of the identity matrix between $\Psi(L)$ and ε_t , as in Centoni and Cubadda (2003),

$$I_n = \Sigma \omega (\omega' \Sigma \omega)^{-1} \omega' + \omega_{\perp} (\omega_{\perp}' \Sigma^{-1} \omega_{\perp})^{-1} \omega_{\perp}' \Sigma^{-1}, \quad (4)$$

one obtains the following decomposition of series Y_t :

$$Y_t = \chi_t + \iota_t \quad (5)$$

where

$$\chi_t = \Psi(L)\Sigma\omega\Sigma^{-1}\varepsilon_t^\chi, \quad (6)$$

$$\iota_t = \Psi(L)\omega_\perp(\omega'_\perp\Sigma^{-1}\omega_\perp)^{-1}\varepsilon_t^\iota, \quad (7)$$

$\Sigma = \omega'\Sigma\omega$, $\varepsilon_t^\chi = \omega'\varepsilon_t$, $\varepsilon_t^\iota = \omega'_\perp\Sigma^{-1}\varepsilon_t$, $E(\varepsilon_t^\chi\varepsilon_t^{\chi'}) = 0$, and $E(\chi_t\iota_{t-j}') = 0$ for $\forall j$.

Since the shock ε_t^χ is one of the indexes, ε_t^χ may be interpreted as the common shock and χ_t as the common components of the series Y_t . Similarly, ε_t^ι and ι_t can be labeled, respectively, as uncommon shocks and an uncommon component.

Interestingly, post-multiplying with $\omega_\perp$ both sides of relation

$$\Psi(L)(I_n - \sum_{j=1}^{p-1} \alpha_j \omega' L^j) = I_n$$

we obtain $\Psi(L)\omega_\perp = \omega_\perp$, which in turn implies that the Wold polynomial matrix of the MAI has the form

$$\Psi(L) = I_n + \sum_{j=1}^{\infty} \theta_j \omega' L^j \quad (8)$$

where θ_j is an $n \times q$ -matrix for $j > 0$.

Having substituted $\Psi(L)$ in Equations (6) and (7) with the RHS of Equation (8), we can finally prove the following proposition.

Proposition 1. *In the MAI, the components of Y_t in (5) read as follows:*

$$\begin{aligned} \chi_t &= (\Sigma\omega\Sigma^{-1} + \sum_{j=1}^{\infty} \theta_j L^j) \varepsilon_t^\chi, \\ \iota_t &= \omega_\perp(\omega'_\perp\Sigma^{-1}\omega_\perp)^{-1}\varepsilon_t^\iota, \end{aligned}$$

where the uncommon component ι_t denotes n -dimensional white noise such that $\text{Rank}(E(\iota_t\iota_t')) = n - q$.

Corollary 1. *The indexes and the common component are linked through the relation $f_t = \omega'\chi_t$, which trivially follows from Proposition 1.*

Remark 2. *In view of Proposition 1, the decomposition (5) has clear analogies with the analogous decomposition in the EDFM. However, differently from the idiosyncratic terms in the EDFM, the uncommon component ι_t is obviously cross-sectionally dependent.³*

Carriero et al. (2016) suggest recovering the structural shocks as linear transformations of the common shock ε_t^χ only. Hence, for most $q < n$, structural shocks can be recovered, as observed in DFMs and in dynamic stochastic general equilibrium models. In principle, all identification strategies that are available for structural VARs or structural DFMs (see Stock & Watson, 2016 and the references therein) can be adopted.

On the estimation side, Carriero et al. (2016) prove that the iterative maximum likelihood procedure proposed by Reinsel (1983) is consistent when $n = o(\sqrt{T})$. Moreover, they provide an MCMC algorithm for Bayesian estimation and show, by simulations, that the Bayesian approach outperforms the classical one when $n = 15, 20$. Finally, they document the practical value of the structural MAI by two empirical applications: the transmission mechanism of monetary policies and the propagation of demand and supply shocks.

2.2. The Vector Heterogeneous Autoregressive Index Model

The univariate heterogeneous AR model [HAR], originally proposed by [Corsi \(2009\)](#) is a popular tool for analyzing and forecasting daily realized volatility [RV] measures without resorting to more involved long-memory models. Technically speaking, the HAR is a constrained AR(22) model where the predictors are the first lags of the following: (i) the daily RV; (ii) the weekly (5 days) average of the daily RV; (iii) the monthly (22 days) average of the daily RV.

[Cubadda et al. \(2017\)](#) propose a multivariate HAR for a set of n daily RV measures, $Y_t^{(d)} \equiv (Y_{1,t}^{(d)}, \dots, Y_{n,t}^{(d)})'$, that is endowed with an index structure. In particular, the vector heterogeneous autoregressive index model [VHARI] reads as follows:

$$Y_t^{(d)} = \alpha^{(d)} \omega' Y_{t-1d}^{(d)} + \alpha^{(w)} \omega' Y_{t-1d}^{(w)} + \alpha^{(m)} \omega' Y_{t-1d}^{(m)} + \varepsilon_t,$$

where (d) , (w) , and (m) denote, respectively, the time horizons of one day, one week, and one month such that

$$Y_t^{(w)} = \frac{1}{5} \sum_{j=0}^4 Y_{t-jd}^{(d)}, \quad Y_t^{(m)} = \frac{1}{22} \sum_{j=0}^{21} Y_{t-jd}^{(d)}$$

The VHARI enjoys two important properties that are not shared by alternative approaches, and they induce dimensional reductions in the vector HAR⁴: First, the index $f_t^{(d)} = \omega' Y_{t-1d}^{(d)}$ preserves the temporal cascade structure of the HAR model since

$$f_t^{(w)} = \frac{1}{5} \sum_{j=0}^4 f_{t-jd}^{(d)}, \quad f_t^{(m)} = \frac{1}{22} \sum_{j=0}^{21} f_{t-jd}^{(d)}.$$

Second, pre-multiplying both sides of the VHARI by ω' yields the following:

$$f_t^{(d)} = \omega' \alpha + \omega' \alpha^{(d)} f_{t-1d}^{(d)} + \omega' \alpha^{(w)} f_{t-1d}^{(w)} + \omega' \alpha^{(m)} f_{t-1d}^{(m)} + \omega' \varepsilon_t,$$

which shows that the indexes follow a multivariate HAR model. In particular, when $q = 1$, a univariate HAR model generated all the dynamics of the n RVs.

On the estimation side, [Cubadda et al. \(2017\)](#) suggest using a switching algorithm [SA], an iterative method for the numerical maximization of the log-likelihood of complex models that has a long tradition in time series analysis (see [Boswijk & Doornik, 2004](#) and the references therein). In particular, the proposed SA requires the following steps:

1. Given an (initial) estimate of ω , maximize the conditional Gaussian likelihood $\ell(A, \Sigma | \omega)$, where $A = [\alpha^{(d)'}, \alpha^{(w)'}, \alpha^{(m)'}]'$.
2. Given the previously obtained estimates of A and Σ , maximize the conditional likelihood $\ell(\omega | A, \Sigma)$.
3. Repeat steps 1 and 2 until numerical convergence occurs.⁵

A key point of the above SA is that both steps 1 and 2 require running OLS regressions only. This feature provides the SA with several advantages over Newton-type optimization methods: computational simplicity, with no need for normalization conditions in ω ; explicit optimization at each step; and the ease of application of regularization schemes or linear restrictions on parameters (see [Cubadda and Guardabascio \(2019\)](#) for additional discussions).

Furthermore, when the SA is initialized with consistent estimates and is iterated sufficiently, the resulting estimator is asymptotically equivalent to the ML one ([Hautsch et al., 2023](#)). [Cubadda et al. \(2017\)](#) show by simulation that the suggested SA

performs well even when elements of ε_t have a log-normal error distribution with GARCH variances.

Following [Patton and Sheppard \(2009\)](#), [Cubadda et al. \(2017\)](#) use a VHARI to build the optimal linear combination of ten different estimators of the volatility of the same market to evaluate its merits through an out-of-sample forecasting exercise. The VHARI model performs well, often outperforming previously existing methods.

2.3. The Index-Augmented Autoregressive Model

A possible limitation of MAI as a forecasting tool is that the only predictors of the series $y_{i,t}$ for $i = 1, \dots, n$ are the lagged indexes, whereas the forecasts obtained through the DFM exploit information coming from the past of both factors and the series $y_{i,t}$ itself (see the seminal contributions by [Stock and Watson \(2002a, 2002b\)](#)). Although the indexes may be interpreted as ‘supervised’ factors that are constructed for emphasizing the comovements between the present and the past of the system, some variables are better predicted by their own lags rather than by any linear combination of all variables only.

In order to overcome such limitations, [Cubadda and Guardabascio \(2019\)](#) extended the basic MAI model by allowing individual AR structures for each element of Y_t . Their key assumption is the following.

Assumption 2. *It holds*

$$\phi_{ik}^{(j)} = \sum_{m=1}^q \alpha_{im}^{(j)} \omega_{km},$$

where $\phi_{ik}^{(j)}$ is the generic element of the polynomial matrix Φ_j , ω_{km} is the generic element of ω , and $\alpha_{im}^{(j)}$ is the generic element of α_j for $j = 1, \dots, p$, $i = 1, \dots, n$, $k = 1, \dots, i-1, i+1, \dots, n$.

In other words, Assumption 2 states that there is a reduced number of channels p through which each variable is influenced by the past of other variables in the system, which is consistent with the common view that few shocks are responsible for most macroeconomic fluctuations.

Under Assumption 2 and using the reparametrization $\delta_{ii}^{(j)} = \phi_{ii}^{(j)} - \sum_{m=1}^q \alpha_{im}^{(j)} \omega_{im}$, Model (1) can be rewritten into the following index-augmented autoregressive model [IAAR]:

$$Y_t = \sum_{j=1}^p D_j Y_{t-j} + \sum_{j=1}^s \alpha_j f_{t-j} + \varepsilon_t, \quad (9)$$

where D_j is a $n \times n$ diagonal matrix with $\delta_{ii}^{(j)}$ as a generic diagonal element, and for greater generality, $s \leq p$.

Remark 3. *Since the number of parameters of Model (9) is equal to $n(qs + q + p) - q^2$, it is necessary to impose proper upper bounds to either q or s to ensure that the MIAAR is more parsimonious than the VAR. To this end, it is easy to see that sufficient conditions are $q < n - 1$ for $s = p \geq 2$ or $s < p - 1$ for any p and $q < n$. However, in empirical applications, the estimated values of q are typically much smaller than n (see [Cubadda & Guardabascio, 2019](#); [Carriero et al., 2022](#)).*

Remark 4. *The individual forecasting equation of the IAAR reads*

$$y_{it+1} = \sum_{j=0}^{p-1} \delta_{ii}^{(j)} y_{it-j} + \sum_{j=0}^{s-1} \alpha_i^{(j)} f_{t-j} + \varepsilon_{it+1}, \quad (10)$$

where $\alpha_{i\cdot}^{(j)'} is the i -th row of matrix α_j . Equation (10) is entirely analogous to the individual forecasting equation of the DFM, with one important difference. Whereas factors are typically estimated using principal component methods, which aim to maximize the contemporaneous variability of series Y_t , the indexes in (10) are constructed explicitly by taking into account the covariability between each series y_{it} and the lags of other elements of Y_t conditionally on the lags of the series y_{it} .$

Remark 5. Interestingly, by the same argument underlying Proposition 1, we see that, differently from the MAI, $\Psi(L)\omega_{\perp} \neq \omega_{\perp}$, which implies, in view of Equation (7), that the uncommon component ι_t is generally autocorrelated in the case of the IAAR. Hence, the decomposition (5) for the IAAR closely resembles the analogous decomposition in the approximate DFM (see Lippi (2019) and the references therein). However, the estimation of the index f_t does not require that $n \rightarrow \infty$, nor does it impose conditions on the autocorrelations and cross-correlations of the elements of ι_t or on the loading α_j as in the approximate DFM.

Cubadda and Guardabascio (2019) proposed a two-step SA for the estimation of the IAAR, along with a variant where a ℓ_2 regularization scheme is applied in both steps. They show, by simulations, that the regularized version of the SA outperforms the standard one with $n = 20$. Regarding model specification, they opt for the use of information criteria [IC], in line with previous contributions showing that IC outperforms likelihood ratio tests in the selection of reduced-rank VAR models (see, e.g., Gonzalo & Pitarakis, 1999; Cavaliere et al., 2015; Cavaliere et al., 2018). Finally, the IAAR proves to outperform well-known macroeconomic forecasting methods when applied to systems with n ranging from 4 to 40.

Carriero et al. (2022) endowed the IAAR with Stochastic Volatility [IAAR-SV] in the error ε_t and offered Bayesian estimations using Markov Chain Monte Carlo [MCMC] techniques. Furthermore, they used (4) to decompose the time-varying volatility $E(\varepsilon_t \varepsilon_t') = \Sigma_t$ as follows:

$$\Sigma_t = \underbrace{\Sigma_t \omega (\omega' \Sigma_t \omega)^{-1} \omega' \Sigma_t}_{\text{common}} + \underbrace{\omega_{\perp} (\omega_{\perp}' \Sigma_t^{-1} \omega_{\perp})^{-1} \omega_{\perp}'}_{\text{uncommon}}$$

Carriero et al. (2022) applied the IAAR-SV to analyze the commonality in both levels and volatilities of inflation rates in several countries, and their main finding is that a substantial fraction of inflation volatility can be attributed to a global factor that also drives inflation levels and their persistence.

2.4. The Time-Varying Multivariate Autoregressive Index Model

A further step towards taking parameter instabilities over time into account was made by Cubadda et al. (2025), who proposed the following MAI with time-varying parameters and time-varying volatility [MAI-TVP-TVV]:

$$Y_t = \sum_{j=1}^p \alpha_{j,t} \omega' Y_{t-j} + \varepsilon_t,$$

$$\alpha_t = \alpha_t + \kappa_t,$$

where $\alpha_t = \text{Vec}(\alpha'_{1,t}, \dots, \alpha'_{n,t})'$, $\varepsilon_t \sim N(0, \Sigma_t)$, $\kappa_t \sim N(0, Q_t)$; ε_t and κ_t are independent at any lag and lead. Notice that it is assumed that the index loadings evolve over time as random walks, while the index weight ω remain stable.

In order to overcome the computational limitation related to MCMC procedures, Cubadda et al. (2025) offer a hybrid estimation method that combines the SA, Kalman filter with forgetting factors (Koop & Korobilis, 2014), and exponentially weighted moving average techniques (Johansson et al., 2023) for the time-varying volatility.

An empirical application, where 25 US quarterly time series are used to forecast three key macroeconomic variables, shows that the MAI-TVP-TVV is one of the best models in a large set of competitors for all targets, improving upon its counterparts, especially for short horizons. Other interesting findings are that once the MAI is endowed with time-varying volatility [MAI-TVV], there are no clear improvements in adding time-varying parameters for point forecasting, but the MAI-TVP-TVV always outperforms the MAI-TVV in density forecasting.

2.5. The Dimension-Reducible VAR

Cubadda and Hecq (2022a) studied the conditions under which the dynamics in a large-dimension VAR are entirely generated by a small-scale VAR. They show that such conditions are met when the coefficient matrices of the large VAR have the same common right space and a common left null space. This entails combining Assumption 1 with the following.

Assumption 3. *The following holds:*

$$\omega'_{\perp} [\Phi_1, \dots, \Phi_p] = 0$$

Assumption 3 is popularly known in time series econometrics as the CSC (see Cubadda & Hecq, 2022b and the reference therein) given that

$$\omega'_{\perp} Y_t = \omega'_{\perp} \varepsilon_t,$$

That is, there exist $(n - q)$ linear combinations of variables Y_t that are white noise, and as such, cyclical behavior cannot be exhibited.

Taking Assumptions 1 and 3 together leads to the dimension-reducible VAR model [DRVAR]:

$$Y_t = \sum_{j=1}^p \omega \phi_j f_{t-j} + \varepsilon_t, \quad (11)$$

where ϕ_j is a $q \times q$ matrix for $j = 1, \dots, p$.

Assuming, without loss of generality, that $\omega' \omega = I_q$ and $\omega'_{\perp} \omega_{\perp} = I_{n-q}$, we can decompose series Y_t as follows:

$$Y_t = \omega f_t + \omega_{\perp} \eta_t, \quad (12)$$

where f_t is the dynamic component, and $\eta_t = \omega'_{\perp} \varepsilon_t$ is the static one. Premultiplying both sides of DRVAR by ω' one obtains

$$f_t = \sum_{j=1}^p \phi_j f_{t-j} + \varepsilon_t^{\chi},$$

where $\varepsilon_t^{\chi} = \omega' \varepsilon_t$, which shows that f_t is generated by a q -dimensional VAR (p) process.

By inserting the Wold representation of the dynamic component f_t in Equation (12), it follows that

$$Y_t = \omega \gamma(L) \varepsilon_t^{\chi} + \omega_{\perp} \eta_t, \quad (13)$$

where $\gamma(L)^{-1} = I_n - \sum_{j=1}^p \phi_j L^j$. Finally, by linearly projecting $\omega_{\perp} \eta_t$ on ε_t^{χ} , we obtain $\omega_{\perp} \eta_t = \rho \varepsilon_t^{\chi} + v_t$ with $E(\varepsilon_t^{\chi} v_t') = 0$, which can be inserted into Equation (13) to obtain

$$Y_t = C(L) \varepsilon_t^{\chi} + v_t, \quad (14)$$

where $C_0 = \omega + \rho$ and $C_j = \omega \gamma_j$ for $j > 0$.

Representation (14) highlights that system dynamics are completely generated by common reduced form errors ε_t^χ . Consequently, Cubadda and Hecq (2022a) label v_t as the ignorable errors, as they are noise without structural interpretation. Since errors ε_t^χ and v_t are uncorrelated at any lead and lag, it is then possible to recover the structural shocks solely from the reduced form errors ε_t^χ of the common component χ_t using any of the procedures that are commonly employed in structural VARs or structural DFMs (see Stock and Watson (2016) and the references therein).

In order to estimate the matrix ω , one may rely on a nonparametric estimator proposed by Lam et al. (2011). The underlying intuition is that the matrix ω lies in the space generated by the eigenvectors associated with the q nonzero eigenvalues of the symmetric and semipositive definite matrix:

$$M = \sum_{j=1}^{p_0} \Sigma_y(j) \Sigma_y(j)',$$

where $p_0 \geq p$, and $\Sigma_y(j)$ is the autocovariance matrix of series Y_t in lag j . Under some regularity conditions, the matrix formed by the eigenvectors associated with the q largest eigenvalues of the sample estimate of M is a $\sqrt{T}$ -consistent estimator of ω (up to an orthonormal transformation) when q is fixed: $n, T \rightarrow \infty$, and $\omega_i' \omega_i = O(n)$ for $i = 1, \dots, q$, where $\omega = [\omega_1, \dots, \omega_q]$. Remarkably, the speed of convergence of the estimator, namely $\sqrt{T}$, is the same as when the dimension n is finite.

Moreover, Cubadda and Hecq (2022a) provide both the OLS and GLS estimators of the coefficient ϕ in Equation (11) and consistent information criteria for the selection of q , and they show by simulations that the proposed methodology works well with the temporal and cross-sectional sizes that are typical in macroeconomics. Finally, the approach is applied to analyze a large set of US economic time series and to identify the shock that is responsible for most of the common volatility in the business cycle frequency band.

2.6. The Vector Error-Correction Index Model

The models considered so far do not explicitly deal with the possible presence of unit roots. Given that most macroeconomic and financial time series are characterized by stochastic trends, it is important to understand how a cointegrated VAR model can be augmented with an index structure.

Let us assume that series Y_t follows the vector error-correction model [VECM]

$$\Delta Y_t = \alpha_0 \beta' Y_{t-1} + \sum_{j=1}^{p-1} \Pi_j \Delta Y_{t-j} + \varepsilon_t, \quad (15)$$

where α_0 and β are full-rank $n \times r$ ($r < n$) matrices such that $\alpha_0 \beta' = -\Phi(1)$, $\Pi_j = -\sum_{i>j} \Phi_i$ for $j = 1, \dots, p-1$, $\alpha_0' \bar{\Pi} \beta_\perp$ is non-singular, and $\bar{\Pi} = I_n - \sum_{j=1}^{p-1} \Pi_j$. Under such assumptions, it is well known that the elements of Y_t are individually at most $I(1)$ and that they are jointly cointegrated with respect to an order 1 in the sense that $\beta' Y_{t-1}$ is $I(0)$ (see Johansen (1995) and the references therein).

To possibly reduce the number of parameters in the VECM, Cubadda and Mazzali (2024) made the following assumptions:

Assumption 4. For $\Pi = [\Pi_1', \dots, \Pi_{p-1}']'$, the following holds:

$$\Pi = A\omega',$$

where ω is a full-rank $n \times q$ matrix with $q < n$, and A is a full-rank $n(p-1) \times q$ matrix.

Assumption 5. The following holds:

$$\beta = \omega\gamma,$$

where γ is a full-rank $q \times r$ matrix with $q \geq r$.

Under Assumptions 4 and 5, Model (15) can be rewritten in the following vector error-correction index model [VECIM]:

$$\Delta Y_t = \alpha_0 \gamma' f_{t-1} + \sum_{j=1}^{p-1} \alpha_j \Delta f_{t-j} + \varepsilon_t,$$

where γ is a full-rank $q \times r$ matrix ($q \geq r$), and α_j is an $n \times q$ matrix for $j = 1, \dots, p-1$ such that $\text{rank}([\alpha_1', \dots, \alpha_{p-1}']') = q$. Notice that the cointegration matrix is given by $\beta = \omega\gamma$.

Interestingly, the indexes f_t themselves are generated by a q -dimensional VECM:

$$\Delta f_t = \alpha_0 \gamma' f_{t-1} + \sum_{j=1}^{p-1} \alpha_j \Delta f_{t-j} + \varepsilon_t^\chi,$$

where $\alpha_j = \omega' \alpha_j$, for $j = 0, 1, \dots, p-1$.

By first inserting the decomposition (5) between $\Psi(L)$ and ε_t into the Wold representation of the first differences ΔY_t :

$$\Delta Y_t = \Psi(L) \varepsilon_t,$$

and then further decomposing the common component χ_t into permanent and transitory subcomponents as in Centoni and Cubadda (2003), we obtain the following:

$$Y_t = \chi_t + \iota_t = \pi_t + \tau_t + \iota_t, \quad (16)$$

where

$$\Delta \pi_t = \Psi(L) \Sigma \omega \Sigma^{-1} \Sigma \alpha_{0\perp} (\alpha_{0\perp}' \Sigma \alpha_{0\perp})^{-1} \underbrace{\alpha_{0\perp}' \varepsilon_t^\chi}_{\varepsilon_t^\pi}, \quad (17)$$

$$\Delta \tau_t = \Psi(L) \Sigma \omega \Sigma^{-1} \alpha_0 (\alpha_0' \Sigma^{-1} \alpha_0)^{-1} \underbrace{\alpha_0' \Sigma^{-1} \varepsilon_t^\chi}_{\varepsilon_t^\tau}, \quad (18)$$

$$\Delta \iota_t = \Psi(L) \omega_\perp (\omega_\perp' \Sigma^{-1} \omega_\perp)^{-1} \underbrace{\omega_\perp' \Sigma^{-1} \varepsilon_t}_{\varepsilon_t^\iota} \quad (19)$$

Since errors ε_t^π are the innovations of the common trends in the indexes f_t (see, e.g., Johansen, 1995) and errors ε_t^τ are such that $E(\varepsilon_t^\pi \varepsilon_t^{\tau'}) = 0$, Cubadda and Mazzali (2024) labeled π_t as the common permanent component and τ_t as the common transitory component, whereas ι_t is the uncommon component given that $E(\varepsilon_t^\iota \varepsilon_t^{\pi'}) = 0$ and $E(\varepsilon_t^\iota \varepsilon_t^{\tau'}) = 0$.

Following a similar reasoning as the one leading to Proposition 1, post-multiplying, with $\omega_\perp$ both sides of the relation

$$\Psi(L) (\Delta I_n - \sum_{j=1}^{p-1} \alpha_j \omega' \Delta L^j - \alpha_0 \gamma' \omega' L) = \Delta I_n$$

we again obtain $\Psi(L) \omega_\perp = \omega_\perp$, which in turn implies that the Wold polynomial matrix of the VECIM has the same form as (8). Finally, inserting (8) in Equations (17)–(19), we can prove the following proposition:

Proposition 2. In the VECIM, the first differences of the components of Y_t in (16) read

$$\begin{aligned}\Delta\pi_t &= (\Sigma\omega\Sigma^{-1} + \sum_{j=1}^{\infty} \theta_j L^j) \Sigma \alpha_{0\perp} (\alpha'_{0\perp} \Sigma \alpha_{0\perp})^{-1} \varepsilon_t^\pi \equiv P(L) \varepsilon_t^\pi, \\ \Delta\tau_t &= (\Sigma\omega\Sigma^{-1} + \sum_{j=1}^{\infty} \theta_j L^j) \alpha_0 (\alpha'_0 \Sigma^{-1} \alpha_0)^{-1} \varepsilon_t^\tau \equiv T(L) \varepsilon_t^\tau \\ \Delta\iota_t &= \omega_\perp (\omega'_\perp \Sigma^{-1} \omega_\perp)^{-1} \varepsilon_t^\iota,\end{aligned}$$

where the uncommon component ι_t is an n -dimensional random walk such that $\text{Rank}(E(\Delta\iota_t \Delta\iota'_t)) = n - q$.

Notice that Proposition 2 implies that Corollary 1 applies to the VECIM as well.

Remark 6. Given that the components in (16) are not correlated with each other at any lag and lead, the VECIM allows one to perform a structural analysis, taking advantage of the features of both the DFM—namely isolating shocks that are common among variables—and the VECM—namely disentangling shocks having transitory or permanent effects. For instance, one may identify the structural transitory shocks as $u_t = C^{-1} D \varepsilon_t^\tau$ and the impulse response functions as $\Theta(L) = T(L) D^{-1} C$, where D is the matrix formed by the first r rows of $T(0)$, and C is a lower triangular matrix such that

$$CC' = D \alpha'_0 \Sigma^{-1} \omega' \Sigma \omega \Sigma^{-1} \alpha_0 D'$$

Since the first r rows of $\Theta(0)$, being equal to C , form a lower triangular matrix, the usual interpretation of structural shocks obtained through a Cholesky factorization applies to u_t .

Cubadda and Mazzali (2024) offered a three-step SA for the estimation of the VECIM and proposed selecting the triple (p, q, r) in a unique search by IC. An extensive Monte Carlo study shows that the proposed methodology works reasonably well for n , ranging from 6 to 18 when the model is identified by the Hannan–Quinn IC. Moreover, in an empirical application, they identified a shock that maximizes the variability of the common transitory component of unemployment at business cycle frequencies and another one that does the same, but for the common permanent component of unemployment. These two shocks are endowed with a neater economic interpretation than compared to a unique main business cycle shock identified according to Angeletos et al. (2020).

3. A New Proposal: The Cointegrated Index-Augmented Autoregressive Model

A possible limitation of the VECIM is that the uncommon component ι_t is necessarily a random walk, which may be considered restrictive for some applications. For example, Barigozzi et al. (2021) proposed a DFM where the idiosyncratic components may be $I(0)$ or $I(1)$.

In order to overcome this issue, one can combine the VECIM with the IAAR. Formally, this involves using Assumption 5 along with the following one:

Assumption 6. For the VECM (15), the following holds:

$$\pi_{ik}^{(j)} = \sum_{m=1}^q \alpha_{im}^{(j)} \omega_{km},$$

where $\pi_{ik}^{(j)}$ is the generic element of the polynomial matrix Π_j for $j = 1, \dots, p-1$, $i = 1, \dots, n$, $k = 1, \dots, i-1, i+1, \dots, n$.

Taking Assumptions 5 and 6, the model (15) can be rewritten into the following cointegrated index-augmented auto-regressive model [CIAAR]:

$$\Delta Y_t = \sum_{j=1}^{p-1} D_j \Delta Y_{t-j} + \alpha_0 \gamma' \omega' Y_{t-1} + \sum_{j=1}^{s-1} \alpha_j \omega' \Delta Y_{t-j} + \varepsilon_t, \quad (20)$$

where D_j is an $n \times n$ diagonal matrix with $\delta_{ii}^{(j)} = \pi_{ii}^{(j)} - \sum_{m=1}^q \alpha_{im}^{(j)} \omega_{im}$ as a generic diagonal element.

When the elements of series Y_t are $I(1)$, Model (20) includes several earlier models for series ΔY_t as special cases, as summarized in Table 1.

Table 1. Previous models as special cases of the CIAAR.

Model	Restrictions on Model (20)	N. of Restrictions
MAI	$\alpha_0 = 0_{n \times r}, \gamma = 0_{q \times r}, D_j = 0_{n \times n}$ for $j = 1, \dots, p-1$	$n(p-1) + r(n+q-r)$
IAAR	$\alpha_0 = 0_{n \times r}, \gamma = 0_{q \times r}$	$r(n+q-r)$
VECIM	$D_j = 0_{n \times n}$ for $j = 1, \dots, p-1$	$n(p-1)$

Remark 7. Interestingly, by the same argument underlying Proposition 2, we see that, differently from the VECIM, $\Psi(L)\omega_{\perp} \neq \omega_{\perp}$, which implies, in view of Equation (19), that the first differences of the uncommon component $\Delta \iota_t$ are generally autocorrelated in the case of the CIAAR. The uncommon component ι_t is still stochastically singular with rank $n - q$. Since system (20) has overall $n - r$ unit roots, while the common component χ_t has $q - r$ unit roots, uncommon component ι_t has $n - q$ unit roots (see [Deistler & Wagner, 2017](#); [Barigozzi et al., 2020](#) on the properties of singular $I(1)$ stochastic processes).

Following [Cubadda and Guardabascio \(2019\)](#) and [Cubadda and Mazzali \(2024\)](#), the estimation procedure is based on an SA where each step is designed to increase the Gaussian likelihood of Model (20). In detail, when $0 < r < q$, the procedure goes as follows:

1. Given (initial) estimates of γ , ω , and $D = [D_1, \dots, D_{p-1}]'$, maximize the conditional Gaussian likelihood $\mathcal{L}(A^\dagger, \Sigma | \gamma, \omega, D)$ by estimating $A^\dagger = [\alpha'_0, A']'$, where $A = [\alpha'_1, \dots, \alpha'_{s-1}]'$, and Σ is applied on OLS with respect to the following equation:

$$\Delta Y_t - \sum_{j=1}^{p-1} D_j \Delta Y_{t-j} = \alpha_0 \gamma' \omega' Y_{t-1} + \sum_{j=1}^{s-1} \alpha_j \omega' \Delta Y_{t-j} + \varepsilon_t$$

2. Premultiply with $\Sigma^{-1/2}$ and apply the Vec operator to both sides of Equation (20); then, use the property $\text{Vec}(ABC) = (C' \otimes A)\text{Vec}(B)$ to obtain

$$\begin{aligned} \Sigma^{-1/2} \Delta Y_t &= \sum_{j=1}^{p-1} (Y'_{t-j} \otimes \Sigma^{-1/2}) \text{Vec}(D_j) + \\ &\quad (Y'_{t-1} \otimes \Sigma^{-1/2} \alpha_0 \gamma' + \sum_{j=1}^{s-1} \Delta Y'_{t-j} \otimes \Sigma^{-1/2} \alpha_j) \text{Vec}(\omega') + \Sigma^{-1/2} \varepsilon_t, \end{aligned}$$

and reparametrize the above model as

$$\begin{aligned} \Sigma^{-1/2} \Delta Y_t &= \sum_{h=1}^{p-1} [(Y'_{t-h} \otimes \Sigma^{-1/2}) M] \delta_h + \\ &\quad (Y'_{t-1} \otimes \Sigma^{-1/2} \alpha_0 \gamma' + \sum_{j=1}^{s-1} \Delta Y'_{t-j} \otimes \Sigma^{-1/2} \alpha_j) \text{Vec}(\omega') + \Sigma^{-1/2} \varepsilon_t, \end{aligned} \quad (21)$$

where δ_j is a n -vector such that $D_j = \text{diag}(\delta_j)$, and M is a binary $n^2 \times n$ -matrix for which its generic element m_{ik} is such that

$$m_{ik} = \begin{cases} 1 & \text{if } i = 1 + (k-1)(n+1), \quad k = 1, \dots, N \\ 0 & \text{otherwise} \end{cases}$$

Given the previously obtained estimates of $A^\dagger$, γ , and Σ , maximize $\mathcal{L}(\omega, D|A^\dagger, \gamma, \Sigma)$ by estimating $\text{Vec}(\omega')$ and $\delta = [\delta'_1, \dots, \delta'_{p-1}]'$ with OLS in Equation (21).

3. Given the previously obtained estimates of ω and D , maximize $\mathcal{L}(\gamma|\omega, D)$ by estimating γ as the eigenvectors that correspond to the r largest eigenvalues of the matrix

$$S_{11}^{-1}S_{10}S_{00}^{-1}S_{01}$$

where $S_{ij} = \sum_{t=p+1}^T R_{i,t}R'_{j,t}$ for $i, j = 0, 1$; $R_{0,t}$ and $R_{1,t}$ are, respectively, the residuals of an OLS regression of $\Delta Y_t - \sum_{j=1}^{p-1} D_j \Delta Y_{t-j}$ and $\omega' Y_{t-1}$ on $[\Delta Y'_{t-1}\omega, \dots, \Delta Y'_{t-s+1}\omega]'$.

4. Repeat steps 1 to 3 until numerical convergence occurs.

When $r = 0$, step 3 is clearly not needed, and steps 1 and 2 must be modified as follows:

- 1.1 Given (initial) estimates of ω and D , maximize $\mathcal{L}(A, \Sigma|\omega, D)$ by estimating A and Σ with OLS on the following model:

$$\Delta Y_t - \sum_{j=1}^{p-1} D_j \Delta Y_{t-j} = \sum_{j=1}^{s-1} \alpha_j \omega' \Delta Y_{t-j} + \varepsilon_t$$

- 2.1 Given the previously obtained estimates of A and Σ , maximize $\mathcal{L}(\omega|A, \Sigma)$ by estimating $\text{Vec}(\omega')$ and δ with OLS on the following model:

$$\Sigma^{-1/2} \Delta Y_t = \sum_{h=1}^{p-1} [(Y'_{t-j} \otimes \Sigma^{-1/2})M] \delta_j + \left(\sum_{j=1}^{s-1} \Delta Y'_{t-j} \otimes \Sigma^{-1/2} \alpha_j \right) \text{Vec}(\omega') + \Sigma^{-1/2} \varepsilon_t,$$

Finally, when $r = q$, we can assume, without loss of generality, that $\gamma = I_q$. Then, step 3 is, again, not needed, whereas steps 1 and 2 must be modified as follows:

- 1.3 Given (initial) estimates of ω and D , maximize $\mathcal{L}(A^\dagger, \Sigma|\omega, D)$ by estimating $A^\dagger$ and Σ with OLS in the following model:

$$\Delta Y_t - \sum_{j=1}^{p-1} D_j \Delta Y_{t-j} = \alpha_0 \omega' Y_{t-1} + \sum_{j=1}^{s-1} \alpha_j \omega' \Delta Y_{t-j} + \varepsilon_t$$

- 2.3 Given the previously obtained estimates of $A^\dagger$ and Σ , maximize $\mathcal{L}(\omega, D|A^\dagger, \Sigma)$ by estimating $\text{Vec}(\omega')$ and δ with OLS in the following model:

$$\begin{aligned} \Sigma^{-1/2} \Delta Y_t = & \sum_{h=1}^{p-1} [(Y'_{t-j} \otimes \Sigma^{-1/2})M] \delta_j + \\ & (Y'_{t-1} \otimes \Sigma^{-1/2} \alpha_0 + \sum_{j=1}^{s-1} \Delta Y'_{t-j} \otimes \Sigma^{-1/2} \alpha_j) \text{Vec}(\omega') + \Sigma^{-1/2} \varepsilon_t \end{aligned}$$

The choice of initial values for the above procedures is discussed in Appendix A, whereas the selection of the quadruple (p, s, q, r) can be carried out by IC sequentially or in a unique search, as suggested by Cubadda and Mazzali (2024).

4. Conclusions and Future Research Directions

The DFM and VAR are, arguably, among the most popular tools in macroeconometrics and financial econometrics. The two approaches should be considered complementary rather than substitutive, since each has its own merits. The MAI represents a link between these two methodologies: On the one hand, it is a VAR with a specific reduced-rank structure that alleviates the dimensionality problem; on the other hand, the MAI and its variants have several analogies with the DFM; in particular, they allow for identifying a small number of common reduced-form errors and for recovering structural shocks from those errors only.

However, the MAI is not affected by some theoretical limitations of the DFM, such as the requirement that the cross-sectional dimension diverges to infinity and the need for specific assumptions on the dynamic correlation structure of the idiosyncratic component and on the factor loadings. In a more practical perspective, VARs with an index structure can also handle features such as stochastic volatility (Carriero et al., 2022) and time-varying parameters (Cubadda et al., 2025), which are not easily accommodated in DFMs.

Recent developments in VAR models with index structures have considerably extended the original MAI formulation, endowing the model with individual autoregressive structures, stochastic volatility, time-varying parameters, high dimensionality, and cointegration. These extensions have proven to be useful tools for detecting common components, obtaining efficiency gains through the imposition of parameter restrictions, performing structural analysis, and boosting forecast accuracy.

Having reviewed most of the recent advances on the MAI and provided new insights on the representation theory underlying the IAAR and the VECIM, a new model, namely the CIAAR, was proposed along with an estimation procedure. The CIAAR extends previous contributions by allowing the VECIM for individual AR structures and the IAAR for cointegration.

There is plenty of room for future research that could be developed in at least three directions. First, the practical relevance of the CIAAR must be investigated both empirically and by simulations. Second, sparsity could be introduced in the MAI and its variants, employing regularized ℓ_1 regressions in the SA in place of OLS. This would open up the possibility of tackling both dimension reduction, through the index structure, and sparsity in the model coefficients, through Lasso and its variants. Third, the approaches considered in this survey could be applied to data with more elaborate dependence structures than vector time series, such as spatiotemporal processes or matrix-tensor time series. First contributions along these lines were provided by Pu et al. (2025), Wang et al. (2022), and Hecq et al. (2024).

Funding: The financial support of MUR under the 20223725WE (PRIN 2022) grant is gratefully acknowledged.

Data Availability Statement: No new data were created or analyzed in this study.

Acknowledgments: Previous versions of this paper were presented at an intermediate workshop on methodological and computational issues in large-scale time series models for economics and finance in Messina, the Villa Mondragone time series symposium in honour of Marco Lippi in Monte Porzio Catone (Rome); the 11th ICEEE in Palermo; and the final workshop on methodological and computational issues in large-scale time series models for economics and finance in Monte Porzio Catone (Rome). The author thanks the participants, as well as three anonymous referees, for their helpful comments and suggestions. The usual disclaimers apply.

Conflicts of Interest: The author declares no conflicts of interest.

Appendix A

The choice of the initial values for an SA is important. Not only is an accurate initialization necessary to boost numerical convergence but the SA is also asymptotically equivalent to the ML one when the parameters to be initialized are consistently estimated (Hautsch et al., 2023).

With reference to the SA in Section 3, the initial values for γ , ω , and D can be obtained as follows:

1. Use the usual Johansen procedure on the model (15) and obtain estimates $\hat{\alpha}_0$, $\hat{\beta}$, and $\hat{\Pi}_j$ for $j = 1, \dots, m$, where $m = \max\{p, s\} - 1$.
2. Construct matrices $\tilde{\Pi}_j = \hat{\Pi}_j - \text{diag}[\hat{\pi}_{11}^{(j)}, \dots, \hat{\pi}_{nn}^{(j)}]'$ for $j = 1, \dots, m$.
3. Construct the matrix $\tilde{\Phi} = [\tilde{\Pi}'_1, \dots, \tilde{\Pi}'_m, \hat{\beta}\hat{\alpha}'_0]'$.
4. Compute the singular-value decomposition $\tilde{\Phi} = U\Lambda V'$, where the singular values are not increasingly ordered, and obtain $\hat{\omega}$ as the matrix formed by the first q columns of V .
5. Compute the q -rank approximation of $\tilde{\Phi}$ as $\tilde{\Phi} = U\bar{\Lambda}V'$, where $\bar{\Lambda}$ is obtained from Λ by setting the smallest $n - q$ singular values to 0.
6. Construct $\bar{\Pi} = [\bar{\Pi}'_1, \dots, \bar{\Pi}'_{p-1}]'$ as the matrix formed by the first $n(p - 1)$ rows of $\tilde{\Phi}$.
7. Construct $\hat{D}_j$ as a diagonal matrix with the diagonal equal to $\text{diag}[\hat{\pi}_{11}^{(j)} - \bar{\pi}_{11}^{(j)}, \dots, \hat{\pi}_{nn}^{(j)} - \bar{\pi}_{nn}^{(j)}]'$ for $j = 1, \dots, s - 1$.

The motivation for the above choices is twofold. First, the asymptotic distribution of the Johansen estimator of β is not affected by restrictions on the short-run parameters (Johansen, 1995), which implies that $\hat{\alpha}_0$, $\hat{\Pi}_j$, and $\tilde{\Pi}_j$ are consistent, although inefficient, estimators of the associated parameters. Second, the right-singular vectors that correspond to the q largest singular values of the matrix $\tilde{\Phi}$ consistently estimate ω (see, e.g., Reinsel et al., 2022). By the same argument, $\bar{\Pi}$ provides a consistent estimator of Π . Finally, the consistency of $\hat{D} = [\hat{D}_1, \dots, \hat{D}_{p-1}]'$ trivially follows from the ones of $\hat{\Pi}$ and $\bar{\Pi}$.

Notes

- ¹ At the end of 2024, the annual citation rate of Reinsel (1983) in Scopus has increased by about 54% in the last 9 years, with the majority of recent citations coming from econometric journals.
- ² Indeed, the matrix ω , once identified through normalizing restrictions, has $q(n - q)$ free parameters.
- ³ Remarkably, when the factors in the EDFM are estimated by some principal components of series Y_t , the sample variance matrix of the estimated idiosyncratic component has a reduced rank as well.
- ⁴ The most obvious alternatives to the VHARI are likely multivariate principal component regression and reduced-rank regression.
- ⁵ A general proof of the convergence of this family of iterative procedures is given by Oberhofer and Kmenta (1974).

References

- Angeletos, G. M., Collard, F., & Dellas, H. (2020). Business-cycle anatomy. *American Economic Review*, 110(10), 3030–3070. [CrossRef]
- Bai, J. (2003). Inferential theory for factor models of large dimensions. *Econometrica*, 71(1), 135–171. [CrossRef]
- Bai, J., & Ng, S. (2002). Determining the number of factors in approximate factor models. *Econometrica*, 70(1), 191–221. [CrossRef]
- Bai, J., & Ng, S. (2006). Confidence intervals for diffusion index forecasts and inference for factor-augmented regressions. *Econometrica*, 74(4), 1133–1150. [CrossRef]
- Bañbura, M., Giannone, D., & Reichlin, L. (2010). Large Bayesian vector auto regressions. *Journal of Applied Econometrics*, 25(1), 71–92. [CrossRef]
- Barigozzi, M., Lippi, M., & Luciani, M. (2020). Cointegration and error correction mechanisms for singular stochastic vectors. *Econometrics*, 8(1), 3. [CrossRef]
- Barigozzi, M., Lippi, M., & Luciani, M. (2021). Large-dimensional dynamic factor models: Estimation of impulse–response functions with I(1) cointegrated factors. *Journal of Econometrics*, 221(2), 455–482. [CrossRef]
- Boswijk, H., & Doornik, J. (2004). Identifying, estimating and testing restricted cointegrated systems: An overview. *Statistica Neerlandica*, 58(4), 440–465. [CrossRef]

- Carriero, A., Clark, T. E., & Marcellino, M. (2015). Bayesian VARs: Specification choices and forecast accuracy. *Journal of Applied Econometrics*, 30(1), 46–73. [CrossRef]
- Carriero, A., Corsello, F., & Marcellino, M. (2022). The global component of inflation volatility. *Journal of Applied Econometrics*, 37(4), 700–721. [CrossRef]
- Carriero, A., Kapetanios, G., & Marcellino, M. (2016). Structural analysis with multivariate autoregressive index models. *Journal of Econometrics*, 192(2), 332–348. [CrossRef]
- Cavaliere, G., De Angelis, L., Rahbek, A., & Taylor, A. (2015). A comparison of sequential and information-based methods for determining the co-integration rank in heteroskedastic VAR models. *Oxford Bulletin of Economics and Statistics*, 77(1), 106–128. [CrossRef]
- Cavaliere, G., De Angelis, L., Rahbek, A., & Taylor, A. (2018). Determining the cointegration rank in heteroskedastic VAR models of unknown order. *Econometric Theory*, 34(2), 349–382. [CrossRef]
- Centoni, M., & Cubadda, G. (2003). Measuring the business cycle effects of permanent and transitory shocks in cointegrated time series. *Economics Letters*, 80(1), 45–51. [CrossRef]
- Corsi, F. (2009). A simple approximate long-memory model of realized volatility. *Journal of Financial Econometrics*, 7(2), 174–196. [CrossRef]
- Cubadda, G., & Guardabascio, B. (2019). Representation, estimation and forecasting of the multivariate index-augmented autoregressive model. *International Journal of Forecasting*, 35(1), 67–79. [CrossRef]
- Cubadda, G., Guardabascio, B., & Grassi, S. (2025). The time-varying multivariate autoregressive index model. *International Journal of Forecasting*, 41(1), 175–190. [CrossRef]
- Cubadda, G., Guardabascio, B., & Hecq, A. (2017). A vector heterogeneous autoregressive index model for realized volatility measures. *International Journal of Forecasting*, 33(2), 337–344. [CrossRef]
- Cubadda, G., & Hecq, A. (2022a). Dimension reduction for high dimensional vector autoregressive models. *Oxford Bulletin of Economics and Statistics*, 84(5), 1123–1152. [CrossRef]
- Cubadda, G., & Hecq, A. (2022b). Reduced rank regression models in economics and finance. In *Oxford handbook of economic forecasting*. Oxford University Press.
- Cubadda, G., Hecq, A., & Palm, F. (2009). Studying co-movements in large multivariate data prior to multivariate modelling. *Journal of Econometrics*, 148(1), 25–35. [CrossRef]
- Cubadda, G., & Mazzali, M. (2024). The vector error correction index model: Representation, estimation and identification. *Econometrics Journal*, 27, 126–150. [CrossRef]
- Deistler, M., & Wagner, M. (2017). Cointegration in singular ARMA models. *Economics Letters*, 155(4), 39–42. [CrossRef]
- Fernández-Villaverde, J., Rubio-Ramírez, J., & Schorfheide, F. (2016). Solution and estimation methods for DSGE models. In J. Taylor, & H. Uhlig (Eds.), *Handbook of macroeconomics* (Vol. 2, pp. 527–724). North Holland.
- Forni, M., & Gambetti, L. (2014). Sufficient information in structural VARs. *Journal of Monetary Economics*, 66(1), 124–136. [CrossRef]
- Forni, M., Hallin, M., Lippi, M., & Reichlin, L. (2000). The generalized dynamic-factor model: Identification and estimation. *Review of Economics and Statistics*, 82(4), 540–554. [CrossRef]
- Forni, M., Hallin, M., Lippi, M., & Reichlin, L. (2009). Opening the black box: Structural factor models with large cross sections. *Econometric Theory*, 25(5), 1319–1347. [CrossRef]
- Forni, M., & Lippi, M. (2001). The generalized dynamic factor model: Representation theory. *Econometric Theory*, 17(6), 1113–1141. [CrossRef]
- Gonzalo, J., & Pitarakis, J. (1999). Dimensionality effect in cointegration analysis. In *Cointegration, causality, and forecasting. A festschrift in honour of Clive WJ Granger* (pp. 212–229). Oxford University Press. [CrossRef]
- Hautsch, N., Okhrin, O., & Ristig, A. (2023). Maximum-likelihood estimation using the zig-zag algorithm. *Journal of Financial Econometrics*, 21(4), 1346–1375. [CrossRef]
- Hecq, A., Margaritella, L., & Smeekes, S. (2023). Granger causality testing in high-dimensional VARs: A post-double-selection procedure. *Journal of Financial Econometrics*, 21(3), 915–958. [CrossRef]
- Hecq, A., Ricardo, I., & Wilms, I. (2024). *Reduced-rank matrix autoregressive models: A medium n approach*. Available online: <https://arxiv.org/abs/2407.07973> (accessed on 20 January 2025).
- Hsu, N. J., Hung, H. L., & Chang, Y. M. (2008). Subset selection for vector autoregressive processes using Lasso. *Computational Statistics & Data Analysis*, 52(7), 3645–3657. [CrossRef]
- Johansen, S. (1995). *Likelihood-based inference in cointegrated vector autoregressive models*. Oxford University Press.
- Johansson, K., Ogut, M. G., Pelger, M., Schmelzer, T., & Boyd, S. (2023). A simple method for predicting covariance matrices of financial returns. *Foundations and Trends in Econometrics*, 12(4), 324–407. [CrossRef]
- Kock, A. B., & Callot, L. (2015). Oracle inequalities for high dimensional vector autoregressions. *Journal of Econometrics*, 186(2), 325–344. [CrossRef]
- Koop, G. M. (2013). Forecasting with medium and large Bayesian VARs. *Journal of Applied Econometrics*, 28(2), 177–203. [CrossRef]

- Koop, G. M., & Korobilis, D. (2014). A new index of financial conditions. *European Economic Review*, 71, 101–116. [\[CrossRef\]](#)
- Lam, C., Yao, Q., & Bathia, N. (2011). Estimation of latent factors for high-dimensional time series. *Biometrika*, 98(4), 901–918. [\[CrossRef\]](#)
- Lippi, M. (2019). Time-domain approach in high-dimensional dynamic factor models. In *Oxford research encyclopedia of economics and finance*. Oxford University Press.
- Oberhofer, W., & Kmenta, J. (1974). A general procedure for obtaining maximum likelihood estimates in generalized regression models. *Econometrica*, 42(3), 579–590. [\[CrossRef\]](#)
- Patton, A., & Sheppard, K. (2009). Optimal combinations of realised volatility estimators. *International Journal of Forecasting*, 25(2), 218–238. [\[CrossRef\]](#)
- Pu, D., Fang, K., Lan, W., Yu, J., & Zhang, Q. (2025). Reduced rank spatio-temporal models. *Journal of Business & Economic Statistics*, 43(1), 98–109.
- Reinsel, G. (1983). Some results on multivariate autoregressive index models. *Biometrika*, 70(1), 145–156. [\[CrossRef\]](#)
- Reinsel, G., Velu, R., & Chen, K. (2022). *Multivariate reduced-rank regression. Theory, methods and applications*. Springer Nature.
- Sims, C. A. (1980). Macroeconomics and reality. *Econometrica*, 48(1), 1–48. [\[CrossRef\]](#)
- Stock, J. H., & Watson, M. W. (2002a). Forecasting using principal components from a large number of predictors. *Journal of the American Statistical Association*, 97(460), 1167–1179. [\[CrossRef\]](#)
- Stock, J. H., & Watson, M. W. (2002b). Macroeconomic forecasting using diffusion indexes. *Journal of Business & Economic Statistics*, 20(2), 147–162.
- Stock, J. H., & Watson, M. W. (2016). Dynamic factor models, factor-augmented vector autoregressions and structural vector autoregressions in macroeconomics. In J. Taylor, & H. Uhlig (Eds.), *Handbook of macroeconomics* (Vol. 2A, pp. 415–525). North Holland.
- Wang, D., Zheng, Y., Lian, H., & Li, G. (2022). High-dimensional vector autoregressive time series modeling via tensor decomposition. *Journal of the American Statistical Association*, 117(539), 1338–1356. [\[CrossRef\]](#)

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.