

The Impact of Stress on Health in Childhood and Adolescence in the Era of the COVID-19 Pandemic

Stefano Cianfarani^{a, b, c} Valentina Pampanini^{a, c}

^aDipartimento Pediatrico Universitario Ospedaliero, IRCCS “Bambino Gesù” Children’s Hospital, Rome, Italy;

^bDepartment of Systems Medicine, Tor Vergata University, Rome, Italy; ^cDepartment of Women’s and Children’s Health, Karolinska Institutet, Stockholm, Sweden

Keywords

Coronavirus disease 2019 · Severe acute respiratory syndrome coronavirus 2 · Stress · Childhood · Adolescence

Abstract

Background: The ongoing pandemic of coronavirus disease 2019 (COVID-19) caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is posing many challenges to global health. Efforts from the whole scientific community have shed light on the pathogenetic mechanisms and the clinical features of SARS-CoV-2 infection as well as on potential therapeutic strategies. **Summary:** The consequences of stress related to social isolation and anxiety generated by the pandemic on mental and physical health are collateral effects that are yet poorly investigated.

© 2021 S. Karger AG, Basel

Introduction

The coronavirus disease 2019 (COVID-19) pandemic as well as the measures taken to contain the outbreak have deeply affected health, economics, and social life worldwide. In comparison with adults, children and adoles-

cents have lower morbidity and mortality rates when infected with the novel severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) [1, 2]. Nevertheless, children are inevitably affected, both directly and indirectly, by the stress induced by the ongoing COVID-19 pandemic [3].

Survival depends on the ability to maintain homeostasis by coping with changing and, in some cases, stressing circumstances. However, stress may induce long-term neuroendocrine and epigenetic changes that may affect health [4]. Indeed, the neuroendocrine response to stress is an excellent example of a plastic system that responds to threats to homeostasis, altering its function to meet current and expected future demands [5]. The key role played by the endocrine system in this homeostatic regulation has been known since the early studies in the 1930s, when the activation of the sympatho-adrenomedullary and hypothalamic-pituitary-adrenal (HPA) axes was shown in response to physical injury as well as perceived psychological threats [6].

The individual response to stressors is heterogeneous depending on both genetics, as elegantly shown in animal models [7], and early programming of HPA, as shown in the seminal work of Levine in 1967 [8]. The neonatal exposure to stressors (i.e., endotoxin) induces a long-term

effect on the mechanisms generating basal pulsatile HPA activity in adult animals [9]. Herein, we review the available evidence on the potential short- and long-term consequences of being exposed to COVID-19 pandemic in childhood and adolescence.

Immediate Consequences

Probably the main indirect effect of the pandemic affecting children well-being is mediated by the impact of epidemic on parents and caregivers. The stress experienced by caregivers may undermine their ability to provide adequate nurturing care. In the present global health and socioeconomic crisis, which affects disadvantaged people more deeply, primary caregivers and parents find increasing difficulty in providing adequate health, nutrition, safety, and care of children. Therefore, this crisis may have particularly harmful effects on children already living in conditions of adversity and stress or with disabilities [10].

Another immediate effect of the pandemic-related stress is the impact on child and adolescent mental health [11]. Social isolation, combined with fear of contagion and quarantine, as well as with potential (mis)information overload, causes chronic stress and is associated with a higher risk of developing anxiety and depression. Social isolation can be defined as an impairment in the quality and/or quantity of the interactions experienced by an individual with other people. While social isolation before COVID-19 pandemic affected most commonly the older population, measures to minimize the spread of SARS-CoV-2 adopted worldwide have exposed all the age groups transversally to the effects of social isolation [12]. Pandemics and related public measures, such as quarantine, have been associated with several negative psychological outcomes, as comprehensively reviewed by Brooks and colleagues [13]. Studies exploring the effects of COVID-19 pandemic on mental health are flourishing with time, showing that the psychological impact of the pandemic deserves immediate strategies of prevention as well as early and long-term intervention [14].

Social isolation has been identified as a potent stressor in species dependent on social interactions. Increased activation of the HPA axis, enhanced glucocorticoid stress responses, and increased levels of corticosterone have been reported in rats following experimental social isolation [15–18]. Furthermore, dysregulation of brain neurotransmitters has been recognized as the mechanism responsible for the development of neuropsychological dis-

orders associated with social isolation [19]. Recently, the impact of COVID-19 and quarantine on children's and adolescents' mental health has been reviewed by Singh and colleagues [20], reporting a significant increase in behavioral and emotional disorders in children as a consequence of school closures. Post-traumatic stress disorder has been identified in 30% of children in quarantine or isolation [21]. Age, socioeconomic/educational status, and preexisting mental health conditions influence the extent of burden on children and adolescents. Children with preexisting mental health conditions are particularly vulnerable, both for the loss of school routines, which represent fundamental coping mechanisms in these conditions, and for the lack of access to mental health services and support groups during quarantine [22].

Moreover, the restriction of physical activity, especially outdoor and in group, represents a further burden on mental health of children and adolescents [23]. Whether these consequences of pandemic on mental health may have an impact on the behavior and particularly on suicide risk in childhood and adolescence is object of investigation [24–26].

The change of life style induced by the pandemic has inevitably led to an increased risk of overweight and obesity in children and adolescents [27, 28]. There is a close interplay between stress and eating behavior, which often interact each other, leading to a vicious circle worsening both stress and adiposity [29]. Adipose tissue is an active endocrine gland and secretes a wide variety of hormones and regulatory factors. These factors reflect the metabolic status of the adipocytes and exert effects on the brain as well as other organs to control energy intake and expenditure [30]. A known stressing condition such as sleep deprivation is associated with increased appetite and insulin resistance [31]. An impairment in sleep quality has been reported by several authors across the world in connection with stress generated by COVID-19-related containment measures [32–34]. Despite these data refer to adult subjects, it is reasonable to speculate that sleep disturbances could also affect children. Chronic exposure to high cortisol levels in turn stimulates mesolimbic reward pathways within the brain, leading to increased intake of palatable food such as sucrose solutions [30]. Therefore, it is not difficult to foresee a further increase of obesity epidemic in the coming years [35].

Finally, a significant increase of cases of central precocious puberty and a faster pubertal progression in girls has been reported since the beginning of COVID-19 pandemic [36, 37]. This might be related to multiple environmental triggers, including the increase in fat mass, the




Fig. 1. Schematic representation of the consequences on mental and physical health associated with COVID-19 pandemic. COVID-19, coronavirus disease 2019.

prolonged exposure to electronic devices, as well as the psychological stress itself [36], whose role in acceleration of puberty timing proposed in past studies [38] remains to be demonstrated.

Although negative consequences of the pandemic deserve attention in order to tailor early and long-term intervention strategies, it is noteworthy that positive effects of COVID-19 pandemic on mental health in children are also existing. In particular, increased proximity with parents and close relatives in appropriate and healthy familiar environment can be of benefit for children. Furthermore, we recently reported an improved glucometabolic control in very young children with type 1 diabetes as an effect of the increased time spent with parents during quarantine [39].

Long-Term Consequences

The long-term impact of COVID-19 epidemic-related stress on health of children and adolescents is still unknown. According to the concept of the Developmental Origins of Health and Disease (DOHaD), early exposure (during fetal and early postnatal life) to a suboptimal en-

vironment may predispose the individual to develop certain diseases in later life [40, 41]. According to the DOHaD concept, parental environmental factors, including diet, body composition, metabolism, and stress, affect the health and chronic disease risk of the offspring throughout their lives. In this context, the period around conception is considered crucial for the long-term parental influences on the health of the next generation. In particular, the exposure to an adverse environment can influence long-term risks of offspring cardiovascular, metabolic, immune, and neurological morbidities [42, 43].

Psychological stress itself is an environmental stressor that can affect DOHaD with a variety of long-term consequences. Maternal stress and suboptimal intrauterine environment may program fetal development by inducing epigenetic changes, ultimately leading to rearrangement of offspring metabolic status [44, 45]. For instance, epigenetic changes involving methylation of genes playing a key role in adiposity regulation account for the observed relationship between intrafamilial childhood adversity and the risk of obesity in young adulthood [46]. A recent systematic review has shown that maternal stress exposes the offspring to a higher risk of obesity, hyperglycemia, insulin resistance, diabetes mellitus, metabolic

syndrome, cardiovascular disease, hypertension, restricted fetal growth, as well as reduced birth, adrenal, and pancreas weights [47]. Furthermore, maternal stress has been associated with increase in incidence of infections and in noninfectious illnesses as well as the risk of asthma in the offspring [48, 49].

As with other stressors, such as environmental chemicals or nutritional imbalance, the effects depend on the type of stress, as well as the strength, timing, and duration of the exposure. A recent meta-analysis explored anxiety and depression in pregnant women during COVID-19 pandemic, identifying a significant increase in the risk of these disorders during pandemic compared to non-pandemic times [50].

Follow-up studies are needed to verify the long-term effects of fetal exposure to maternal stress related to the COVID-19 pandemic. Interestingly, accelerated shortening of telomeres has been identified as a potential biomarker of stress environments, including social stress, and social disadvantage has been linked to accelerated telomere shortening in children [51].

Conclusions

Living in the time of COVID-19 pandemic exposes children and adolescents to stress in both a direct and an indirect way (Fig. 1). Pediatricians and pediatric endocrinologists should be informed about the potential consequences of the exposure to the stress related to the pandemic in order to provide a more careful monitoring of

children health in the coming years. Media should be involved in an awareness campaign to inform parents and caregivers, in a communication tailored to meet children needs, and in a global educational strategy aimed at mitigating the short- and long-term impact of stress on health of children and adolescents.

Key Messages

In this minireview, we report the available evidence on the potential short- and long-term consequences of being exposed to COVID-19 pandemic in childhood and adolescence.

Conflict of Interest Statement

S.C. is the Editor-in-Chief of *Hormone Research in Paediatrics*. V.P. has no conflicts of interest to declare.

Funding Sources

This work was supported by Ministero della Salute (RRC-2020-23669481).

Author Contributions

Both S.C and V.P. participated in the conception, design, and writing of the manuscript.

References

- 1 Zimmermann P, Curtis N. Why is COVID-19 less severe in children? A review of the proposed mechanisms underlying the age-related difference in severity of SARS-CoV-2 infections. *Arch Dis Child*. 2020.
- 2 Ludvigsson JF. Systematic review of COVID-19 in children shows milder cases and a better prognosis than adults. *Acta Paediatr*. 2020;109(6):1088–95.
- 3 Jiao WY, Wang LN, Liu J, Fang SF, Jiao FY, Pettoello-Mantovani M, et al. Behavioral and emotional disorders in children during the COVID-19 epidemic. *J Pediatr*. 2020;221:264–e1.
- 4 Russell G, Lightman S. The human stress response. *Nat Rev Endocrinol*. 2019;15(9):525–34.
- 5 Lightman SL. The neuroendocrinology of stress: a never ending story. *J Neuroendocrinol*. 2008;20(6):880–4.
- 6 Szabo S, Tache Y, Somogyi A. The legacy of Hans Selye and the origins of stress research: a retrospective 75 years after his landmark brief “letter” to the editor# of nature. *Stress*. 2012;15(5):472–8.
- 7 Sternberg EM, Young WS, Bernardini R, Calogero AE, Chrousos GP, Gold PW, et al. A central nervous system defect in biosynthesis of corticotropin-releasing hormone is associated with susceptibility to streptococcal cell wall-induced arthritis in Lewis rats. *Proc Natl Acad Sci U S A*. 1989;86(12):4771–5.
- 8 Levine S. Maternal and environmental influences on the adrenocortical response to stress in weanling rats. *Science*. 1967;156(3772):258–60.
- 9 Shanks N, Windle RJ, Perks PA, Harbuz MS, Jessop DS, Ingram CD, et al. Early-life exposure to endotoxin alters hypothalamic-pituitary-adrenal function and predisposition to inflammation. *Proc Natl Acad Sci U S A*. 2000;97(10):5645–50.
- 10 Yoshikawa H, Wuermli AJ, Britto PR, Dreyer B, Leckman JF, Lye SJ, et al. Effects of the global coronavirus disease-2019 pandemic on early childhood development: short- and long-term risks and mitigating program and policy actions. *J Pediatr*. 2020;223:188–93.
- 11 Fegert JM, Vitiello B, Plener PL, Clemens V. Challenges and burden of the Coronavirus 2019 (COVID-19) pandemic for child and adolescent mental health: a narrative review to highlight clinical and research needs in the acute phase and the long return to normality. *Child Adolesc Psychiatry Ment Health*. 2020;14:20.
- 12 Clair R, Gordon M, Kroon M, Reilly C. The effects of social isolation on well-being and life satisfaction during pandemic. *Humanit Soc Sci Commun*. 2021;8.

- 13 Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet*. 2020;395(10227):912–20.
- 14 Galea S, Merchant RM, Lurie N. The mental health consequences of COVID-19 and physical distancing: the need for prevention and early intervention. *JAMA Intern Med*. 2020; 180(6):817–8.
- 15 Gamallo A, Villanua A, Trancho G, Fraile A. Stress adaptation and adrenal activity in isolated and crowded rats. *Physiol Behav*. 1986; 36(2):217–21.
- 16 Hawkey LC, Cole SW, Capitanio JP, Norman GJ, Cacioppo JT. Effects of social isolation on glucocorticoid regulation in social mammals. *Horm Behav*. 2012;62(3):314–23.
- 17 Toth M, Mikics E, Tulogdi A, Aliczki M, Haller J. Post-weaning social isolation induces abnormal forms of aggression in conjunction with increased glucocorticoid and autonomic stress responses. *Horm Behav*. 2011; 60(1):28–36.
- 18 Weiss IC, Pryce CR, Jongen-Rêlo AL, Nanz-Bahr NI, Feldon J. Effect of social isolation on stress-related behavioural and neuroendocrine state in the rat. *Behav Brain Res*. 2004; 152(2):279–95.
- 19 Mumtaz F, Khan MI, Zubair M, Dehpour AR. Neurobiology and consequences of social isolation stress in animal model-A comprehensive review. *Biomed Pharmacother*. 2018;105: 1205–22.
- 20 Singh S, Roy D, Sinha K, Parveen S, Sharma G, Joshi G. Impact of COVID-19 and lockdown on mental health of children and adolescents: a narrative review with recommendations. *Psychiatry Res*. 2020;293:113429.
- 21 Demaria F, Vicari S. COVID-19 quarantine: psychological impact and support for children and parents. *Ital J Pediatr*. 2021;47(1):58.
- 22 Lee J. Mental health effects of school closures during COVID-19. *Lancet Child Adolesc Health*. 2020;4(6):421.
- 23 Burtcher J, Burtcher M, Millet GP. (Indoor) isolation, stress, and physical inactivity: vicious circles accelerated by COVID-19? *Scand J Med Sci Sports*. 2020;30(8):1544–5.
- 24 Balachandran AK, Alagarsamy S, Mehroliia S. Suicide among children during Covid-19 pandemic: an alarming social issue. *Asian J Psychiatr*. 2020;54:102420.
- 25 Isumi A, Doi S, Yamaoka Y, Takahashi K, Fujiwara T. Do suicide rates in children and adolescents change during school closure in Japan? The acute effect of the first wave of COVID-19 pandemic on child and adolescent mental health. *Child Abuse Negl*. 2020;110(Pt 2):104680.
- 26 Mourouvaye M, Botteman H, Bonny G, Fourcade L, Angoulvant F, Cohen JF, et al. Association between suicide behaviours in children and adolescents and the COVID-19 lockdown in Paris, France: a retrospective observational study. *Arch Dis Child*. 2020.
- 27 Pietrobelli A, Pecoraro L, Ferruzzi A, Heo M, Faith M, Zoller T, et al. Effects of COVID-19 lockdown on lifestyle behaviors in children with obesity living in Verona, Italy: a Longitudinal Study. *Obesity*. 2020;28(8):1382–5.
- 28 Cuschieri S, Grech S. COVID-19: a one-way ticket to a global childhood obesity crisis? *J Diabetes Metab Disord*. 2020:1–4.
- 29 Doom JR, Lumeng JC, Sturza J, Kaciroti N, Vazquez DM, Miller AL. Longitudinal associations between overweight/obesity and stress biology in low-income children. *Int J Obes*. 2020;44(3):646–55.
- 30 Dallman MF, Pecoraro NC, La Fleur SE, Warne JP, Ginsberg AB, Akana SF, et al. Glucocorticoids, chronic stress, and obesity. *Prog Brain Res*. 2006;153:75–105.
- 31 McEwen BS. Sleep deprivation as a neurobiologic and physiologic stressor: allostasis and allostatic load. *Metabolism*. 2006;55(10):S20–3.
- 32 Batool-Anwar S, Robbins R, Ali SH, Capasso A, Foreman J, Jones AM, et al. Examining changes in sleep duration associated with the onset of the COVID-19 pandemic: who is sleeping and who is not? *medRxiv*. 2021.
- 33 Franceschini C, Musetti A, Zenesini C, Palagini L, Scarpelli S, Quattropiani MC, et al. Poor sleep quality and its consequences on mental health during the COVID-19 lockdown in Italy. *Front Psychol*. 2020;11:574475.
- 34 Lin LY, Wang J, Ou-Yang XY, Miao Q, Chen R, Liang FX, et al. The immediate impact of the 2019 novel coronavirus (COVID-19) outbreak on subjective sleep status. *Sleep Med*. 2021;77:348–54.
- 35 Dietz WH. The obesity epidemic in young children. Reduce television viewing and promote playing. *BMJ*. 2001;322(7282):313–4.
- 36 Stagi S, De Masi S, Bencini E, Losi S, Paci S, Parpagnoli M, et al. Increased incidence of precocious and accelerated puberty in females during and after the Italian lockdown for the coronavirus 2019 (COVID-19) pandemic. *Ital J Pediatr*. 2020;46(1):165.
- 37 Verzani M, Bizzarri C, Chioma L, Bottaro G, Pedicelli S, Cappa M. Impact of COVID-19 pandemic lockdown on early onset of puberty: experience of an Italian tertiary center. *Ital J Pediatr*. 2021;47(1):52.
- 38 Wierson M, Long PJ, Forehand RL. Toward a new understanding of early menarche: the role of environmental stress in pubertal timing. *Adolescence*. 1993;28(112):913–24.
- 39 Schiaffini R, Barbetti F, Rapini N, Inzaghi E, Deodati A, Patera IP, et al. School and pre-school children with type 1 diabetes during Covid-19 quarantine: the synergic effect of parental care and technology. *Diabetes Res Clin Pract*. 2020;166:108302.
- 40 Gluckman PD, Hanson MA, Pinal C. The developmental origins of adult disease. *Matern Child Nutr*. 2005;1(3):130–41.
- 41 Fleming TP, Watkins AJ, Velazquez MA, Mathers JC, Prentice AM, Stephenson J, et al. Origins of lifetime health around the time of conception: causes and consequences. *Lancet*. 2018;391(10132):1842–52.
- 42 Gluckman PD, Hanson MA, Cooper C, Thornburg KL. Effect of in utero and early-life conditions on adult health and disease. *N Engl J Med*. 2008;359(1):61–73.
- 43 Grandjean P, Barouki R, Bellinger DC, Castelleyn L, Chadwick LH, Cordier S, et al. Life-long implications of developmental exposure to environmental stressors: new perspectives. *Endocrinology*. 2015;156(10):3408–15.
- 44 Fowden AL. Stress during pregnancy and its life-long consequences for the infant. *J Physiol*. 2017;595(15):5055–6.
- 45 Deodati A, Inzaghi E, Cianfarani S. Epigenetics and in utero acquired predisposition to metabolic disease. *Front Genet*. 2019;10: 1270.
- 46 Kaufman J, Montalvo-Ortiz JL, Holbrook H, O'Loughlin K, Orr C, Kearney C, et al. Adverse childhood experiences, epigenetic measures, and obesity in youth. *J Pediatr*. 2018; 202:150–e3.
- 47 Eberle C, Fasig T, Brüseke F, Stichling S. Impact of maternal prenatal stress by glucocorticoids on metabolic and cardiovascular outcomes in their offspring: a systematic scoping review. *PLoS One*. 2021;16(1):e0245386.
- 48 Bush NR, Savitz J, Coccia M, Jones-Mason K, Adler N, Boyce WT, et al. Maternal stress during pregnancy predicts infant infectious and noninfectious illness. *J Pediatr*. 2021;228: 117–e2.
- 49 Al-Hussainy A, Mohammed R. Consequences of maternal psychological stress during pregnancy for the risk of asthma in the offspring. *Scand J Immunol*. 2021;93(1):e12919.
- 50 Hessami K, Romanelli C, Chiurazzi M, Cozzolino M. COVID-19 pandemic and maternal mental health: a systematic review and meta-analysis. *J Matern Fetal Neonatal Med*. 2020: 1–8.
- 51 Mitchell C, Hobcraft J, McLanahan SS, Siegel SR, Berg A, Brooks-Gunn J, et al. Social disadvantage, genetic sensitivity, and children's telomere length. *Proc Natl Acad Sci U S A*. 2014;111(16):5944–9.