

ARTICLE OPEN



Pharmacogenetic testing in Italy: results of a nationwide survey by the Joint Working Group for the pharmacogenetics implementation in Italy

Rossana Roncato^{1,2}, Samantha Perfler¹, Martina Gambon¹, Enrico Zoroddu³, Monica Rosa Miozzo⁴, Sabrina Angelini⁵, Massimo Gennarelli^{6,7}, Gabriele Stocco^{8,9}, Valeria Conti¹⁰, Amelia Filippelli¹⁰, Paola Borgiani¹¹, Giuseppe Novelli¹¹, Alessio Squassina¹², Matteo Floris³, Erika Cecchin¹ and The Joint Italian Working Group on Pharmacogenetics Implementation*

© The Author(s) 2026

Pharmacogenetics enables personalization of drug therapy based on an individual's genetic profile. Despite clinical relevance, implementation of pharmacogenetics remains limited. In Italy, integration is fragmented, with heterogeneous practices and a lack of national coordination. A comprehensive assessment of the current landscape is essential. A nationwide survey was conducted between January and October 2025 to map laboratories providing pharmacogenetic testing. A structured questionnaire collected data on institutional characteristics, testing workflows, pharmacogene panels, analytical methodologies, interpretation procedures, and reimbursement. Forty-nine laboratories participated (response rate: 65%). Most were part of public institutions (82%), primarily general or research hospitals. Testing was predominantly performed in medical genetics units (39%) and focused on oncology, specifically *DPYD* (94%) and *UGT1A1* (84%) for fluoropyrimidine and irinotecan therapies. Adherence to national (SIF/AIOM) and international (CPIC/DPWG) guidelines was generally high; compliance with AMP Tier 1 analytical standards varied substantially. Pharmacological counseling was provided by only 29% of laboratories, mainly by clinical pharmacology units. Considerable heterogeneity emerged in testing platforms, bioinformatics tools, and the use of CE-IVD-certified kits. Marked geographical disparities were evident, with pharmacogenetic activity concentrated in Northern Italy. This survey provides the first national overview of pharmacogenetics implementation in Italy, revealing variability in laboratory practices, interpretation standards, and clinical integration. While oncology-related testing is widely adopted and guideline adherence is increasing, the lack of a coordinated national framework restricts consistency and equitable access. Establishing a coordinated network of pharmacogenetic laboratories with harmonized standards for testing, reporting, and education is crucial for evidence-based pharmacogenetic care.

European Journal of Human Genetics; <https://doi.org/10.1038/s41431-026-02063-4>

INTRODUCTION

Pharmacogenetics examines the influence of genetic variability on drug response, providing the opportunity to tailor pharmacological treatments, thereby enhancing efficacy and reducing the risk of adverse drug reactions [1]. In recent years, multiple drivers, including decreasing sequencing costs, the availability of large-scale genomic resources, and substantial evidence from clinical trials, have accelerated the integration of pharmacogenetics into healthcare systems [2–4]. These advances are reflected in the growing number of international guidelines developed by scientific societies and expert consortia, which provide dosing

recommendations for a wide range of clinically relevant gene–drug pairs [5,6].

However, despite this favorable scenario, the implementation of pharmacogenetics remains fragmented in several countries, including Italy. Although pharmacogenetic information is increasingly available on drug labels and national guidelines have been issued for selected gene–drug pairs, its translation into clinical practice remains inconsistent [7]. Furthermore, the language used in regulatory documents is not always sufficiently explicit to guide healthcare professionals on *when* and *how* to apply pharmacogenetic testing [8,9].

¹Experimental and Clinical Pharmacology, Centro di Riferimento Oncologico di Aviano IRCCS, Aviano, Italy. ²Department of Medicine, University of Udine, Udine, Italy. ³Department of Biomedical Sciences, University of Sassari, Sassari, Italy. ⁴Medical Genetics, Department of Health Sciences, Università degli Studi di Milano, Milano, Italy. ⁵Department of Pharmacy and Biotechnology (FABIT), University of Bologna, Bologna, Italy. ⁶Department of Molecular and Translational Medicine, University of Brescia, Brescia, Italy. ⁷Genetics Unit, IRCCS Istituto Centro San Giovanni di Dio Fatebenefratelli, Brescia, Italy. ⁸Department of translational and advanced diagnostics, Institute for maternal and child health IRCCS Burlo Garofalo, Trieste, Italy. ⁹Department of medical, surgical and health sciences, University of Trieste, Trieste, Italy. ¹⁰Department of Medicine, Surgery, and Dentistry “Scuola Medica Salernitana”, University of Salerno, Baronissi (SA), Italy. ¹¹Department of Biomedicine and Prevention, Section of Genetics, University of Rome “Tor Vergata”, Roma, Italy. ¹²Section of Neuroscience and Clinical Pharmacology, Department of Biomedical Sciences, University of Cagliari, Cagliari, Italy. *A list of authors and their affiliations appears at the end of the paper. ✉email: matteo.floris@gmail.com; eccechin@cro.it

Received: 20 December 2025 Revised: 5 February 2026 Accepted: 12 February 2026

Published online: 03 April 2026

A recent nationwide assessment of laboratories offering pharmacogenetic services (from the point of view of an external quality provider), documented a marked increase in the uptake of *DPYD* and *UGT1A1* testing over the past five years in Italy, together with improved adherence to guidelines and greater accuracy in clinical interpretation [10,11]. Nonetheless, considerable heterogeneity remains in terms of the panels used, timing of testing, the format in which results are communicated, and whether or not genetic counseling and pharmacological recommendations are provided alongside the results.

Overall, these findings converge on a critical issue: Italy lacks a standardized national framework for pharmacogenetic testing. Practices differ widely in terms of which tests are performed, when they are requested, who is responsible for them, turnaround times, and whether reports should be accompanied by clinical interpretation or pharmacological consultation [10].

This fragmentation underscores the need for a structured approach to harmonization. As a first and fundamental task, we prioritized conducting a national survey to map laboratories offering pharmacogenetic testing across Italy. This initiative aims to provide a systematic overview of existing services, promote networking among laboratories, and generate the evidence base required to inform national recommendations. Building on these premises, the objectives of the present work are to: (i) conduct a comprehensive nationwide survey of pharmacogenetic services in Italy; (ii) establish a multidisciplinary network of professionals in pharmacogenetics; (iii) promote integration of pharmacogenetics into clinical practice; (iv) harmonize infrastructure development across the country; (v) raise awareness among Italian stakeholders regarding pharmacogenetics implementation and regulation; (vi) increase international visibility for Italian pharmacogenetics initiatives; and (vii) create a collaborative network to foster the exchange of pharmacological, genetic, and clinical expertise.

METHODS

Survey design

A structured questionnaire was developed to systematically collect information on the organization and delivery of pharmacogenetic testing in Italy (Supplementary Material). The survey was designed to capture: (i) the range of pharmacogenetic tests offered; (ii) clinical contexts in which testing is requested; (iii) healthcare professionals responsible for ordering, signing the pharmacogenetic report and interpreting results; (iv) turnaround times; and (v) whether reports were accompanied by a clinical interpretation or pharmacological consultation. The design was based on by previous national experiences of implementing pharmacogenetic testing [12].

Identification of laboratories

A comprehensive list of laboratories potentially offering pharmacogenetic services was established from multiple sources, including institutional and hospital websites, national and regional registries, published reports, original articles and professional networks- complemented by targeted online search aimed at identifying germline pharmacogenetic laboratories operating in Italy. Both public and private laboratories were deemed eligible for inclusion.

Recruitment and data collection

Each identified laboratory was contacted by email and invited to participate in the survey. The invitation included a cover letter outlining the project's objectives and providing a secure link to the online questionnaire. Data collection was conducted between January and October 2025.

To maximize participation, a structured follow-up strategy was applied. Non-responding laboratories received up to two reminder emails at two-week intervals. If no response was obtained, telephone calls were made to ensure that the invitation reached the appropriate contact person within the institution.

Data handling and analysis

Survey responses were collected electronically and stored in a secure database. Data were analyzed descriptively to summarize the distribution

of pharmacogenetic testing practices across institutions. Results are reported in aggregate to ensure the confidentiality of participating laboratories.

RESULTS

Survey respondent demographics

A total of 75 laboratories were contacted, of which 49 participated in the nationwide survey (response rate = 65%) (Fig. 1A). General hospitals were the most represented institutions ($n = 20$; 41%), followed by national research hospitals (IRCCS) ($n = 16$; 33%) and university hospitals ($n = 10$; 20%). Private laboratories ($n = 2$; 4%) and other institutions ($n = 1$; 2%) were only marginally represented (Fig. 1B). The majority of responders were public institutions ($n = 40$; 82%), while a smaller proportion were private facilities ($n = 9$; 18%) (Fig. 1C). Among private facilities, 67% ($n = 6$) were located in the Lombardy region.

Laboratory characteristics

Medical Genetics laboratories accounted for the largest proportion of pharmacogenetic testing ($n = 19$; 39%), followed by Clinical Pathology and Biochemistry units ($n = 9$; 18%) and Clinical Pharmacology departments ($n = 6$; 12%), together representing about two-thirds of all testing activities. Pathology units accounted for only 4% ($n = 2$), while 27% ($n = 13$) of responses were classified as "other" (Supplementary Table 1).

The training background of laboratory directors was diverse but mainly represented by geneticists ($n = 23$; 47%), followed by clinical pathologists/biochemists ($n = 9$; 18%) and pharmacologists ($n = 5$; 10%). Smaller proportions were pathologists ($n = 2$; 4%), hematologists ($n = 2$; 4%), and oncologists ($n = 1$; 2%), while 14% ($n = 7$) reported other specialties (Fig. 2A).

Pharmacogenetic testing was performed across multiple care settings: 90% ($n = 44$) of laboratories tested outpatients, 86% ($n = 42$) inpatients, and 45% ($n = 22$) private-care patients. Most laboratories ($n = 31$; 63%) processed both internal and external to the hospital requests, while 27% ($n = 13$) handled only internal requests and 10% ($n = 5$) only external ones, mainly from private providers (Supplementary Table 1).

Pharmacogenetic testing is primarily provided to support anticancer therapy. As shown in Fig. 2B The vast majority of laboratories tested patients to receive either fluoropyrimidines ($n = 46$; 94%) or irinotecan ($n = 41$; 84%), reflecting the widespread adoption of *DPYD* and *UGT1A1* testing. Thiopurines ($n = 16$; 33%), clopidogrel ($n = 8$; 16%), antidepressants ($n = 4$; 8%), and siponimod ($n = 13$; 27%) followed, while abacavir ($n = 9$; 18%), opioids ($n = 6$; 12%), voriconazole ($n = 3$; 6%) tamoxifen ($n = 2$; 4%), and a small number of other drugs including atazanavir, antiemetics, proton pump inhibitors, isoniazid, and tacrolimus ($n = 1$; 2% each) were less common. As shown in Fig. 2C, the implementation of pharmacogenetic testing among the laboratories participating in the survey was predominantly focused on oncology. Specifically, *DPYD* (fluoropyrimidine-related) and *UGT1A1* (irinotecan-related) genotyping represented 59% of the tests offered as diagnostic service.

Accordingly, requests originated predominantly from oncologists ($n = 46$; 94%), with smaller proportions from infectious diseases specialists and surgeons ($n = 5$; 10% each), neurologists ($n = 4$; 8%), cardiologists and radiotherapists ($n = 3$; 6% each), and a few from rheumatologists, geneticists, psychiatrists, and general practitioners (Supplementary Table 1).

Testing strategies

The selection of variant panels for each tested gene was evaluated in comparison with the recently published AMP (Association for Molecular Pathology) guidelines Tier 1 panels with variable results across pharmacogenes. *TPMT* showed the highest alignment (91%), followed by *CYP2C19* (63%). *CYP2C9* (17%) and *CYP2D6*




Fig. 1 Geographic distribution and institutional characteristics of participating laboratories. **A** Map showing survey response rate by region. **B** Distribution of participating laboratories by institution type. **C** Distribution by institutional setting: public vs. private.

(20%) showed lower concordance. No laboratory adopted the AMP Tier 1 panel for *DPYD*, but instead, the selection of variants was largely consistent with the national SIF-AIOM panel (81%). For *UGT1A1*, alignment with the SIF-AIOM panel was 32%, whereas agreement with DPWG recommended variants reached 59%. Even when alignment with the AMP Tier 1 panel was low for *CYP2C19* and *CYP2C9*, the laboratories showed higher concordance rates to other panels, at 63% and 67%, respectively. Notably, *CYP2D6* exhibited marked variability in the selection of testing panels across laboratories (Table 1).

Considering the current availability of Italian pharmacogenetic guidelines for *DPYD*/ fluoropyrimidines and *UGT1A1*/ irinotecan, we evaluated their consistency with the variant panels recommended by various scientific societies/consortia (i.e., AMP, CPIC, DPWG, and RNPgX) for these two gene/drug interactions (Supplementary Table 2). For *DPYD*, a high degree of concordance was observed between the different panels adopted, with the *DPYD**2 A, *DPYD**13, c.1236 G > A (or c.1129-5923 C > G), and c.2846 A > T included in all except RNPgX, which does not include any HapB3 marker polymorphisms. The SIF-AIOM panel also includes the *DPYD**6 variant (to be tested in a post-toxicity context), while the AMP Tier 1 panel is the most comprehensive of all. For *UGT1A1*, all panels include *UGT1A1**28/*36/*37 and *UGT1A1**6, except for the SIF-AIOM panel, which only includes *UGT1A1**28 and *UGT1A1**6 and RNPgX, which does not include *UGT1A1**6. Currently, there are no CPIC guidelines available for *UGT1A1*/ irinotecan.

Testing was almost universally provided pre-treatment for *DPYD* (100%) and *UGT1A1* (97%), while semi-preemptive strategies were reported for *CYP2C19* (53%) and *HLA-B* (50%). A reactive testing strategy is the least frequent overall, though it appears quite common for *CYP2D6* (39%), which is frequently tested in response to drug-related adverse events or for therapy adjustments.

qPCR was the most commonly used method (38–90% depending on the gene), while targeted next-generation sequencing (NGS) was mainly applied to *HLA-B* (40%). International CPIC and DPWG guideline adherence for result interpretation was highest for *CYP2C19* (88%), *CYP2D6* (80%), and *CYP2C9* (83%), and lower for *DPYD* and *UGT1A1*, reflecting a major reliance on national SIF-AIOM recommendations.


Adoption of CE-IVD-certified kits varied widely across pharmacogenes (Supplementary Table 3). The highest level of implementation was reported for *DPYD* (38 centers, 11 different kits) and *UGT1A1* (31 centers, 7 kits), confirming their established role in oncology. In contrast, CE-IVD use was limited for *CYP2C19* (3 out of 13 centers) and *CYP2C9* (2 out of 13 centers), each employing two distinct kits. *TPMT* and *HLA-B* were tested with certified assays in only two centers, while a single-center use was reported for *ABCB1*, *CYP2D6*, and *HLA*. *F5L* and *NAT2* were analyzed exclusively with non-certified methods. Overall, CE-IVD implementation predominated for *DPYD* and *UGT1A1* but remained fragmented for other pharmacogenes. Only one laboratory reported using whole-exome sequencing (WES) for pharmacogenetic testing.

A comparable heterogeneity emerged in the use of bioinformatics tools (Supplementary Table 4). Overall, 18 different software programs were reported, with the most significant variability observed for *DPYD* and *UGT1A1*, analyzed by 31 and 28 centers, respectively, using approximately ten distinct software solutions each. These findings underscore the lack of standardization in data interpretation across laboratories.

Interpretation and reporting

The personnel responsible for signing pharmacogenetic reports are most frequently geneticists by training, accounting for nearly two-thirds of cases ($n = 32$; 65%). Clinical pathologists and biochemists represented the second most common group ($n = 15$; 31%), followed by pharmacologists ($n = 6$; 12%), while 8% ($n = 4$) of laboratories reported other training backgrounds for the personnel allowed to sign pharmacogenetic reports.

Nearly all laboratories ($n = 44$; 90%) usually provide a written clinical interpretation of the results embedded in the pharmacogenetic report. These comments most frequently include an indication of the presence of a risk phenotype for toxicity or lack of efficacy ($n = 36$; 73%), followed by the interpretation of the predicted phenotype based on the genotype ($n = 23$; 47%). Only about one quarter ($n = 12$; 24%) of laboratories include specific dosing or therapeutic recommendations (pharmacological counseling), and 31% ($n = 15$) reported including a reference to relevant clinical guidelines or redirecting the patient to a specialist (Supplementary Table 1).



Most laboratories ($n = 36$; 73%) reported obtaining informed consent before performing pharmacogenetic testing. Among these, 12 laboratories used a consent form specifically designed for pharmacogenetic testing, while the remaining laboratories reported using a general consent form for germline analysis. The vast majority of laboratories ($n = 44$; 90%) indicated that pharmacogenetic incidental findings (e.g., those obtained through clinical exome sequencing) are not communicated to patients (Supplementary table 1).

Nineteen responders (40%) provided suggestions for improving pharmacogenetics implementation in Italy, selecting from various proposed items. The most frequent item was "Collaboration and

Most laboratories (73%) reported full reimbursement of pharmacogenetic testing by the National Health System, 22% reported partial reimbursement, and 4% no reimbursement. However, reimbursement policies vary across regions, reflecting

Table 1. Compliance of tested variant panels and clinical interpretation with different pharmacogenetic guidelines, and testing approaches for key pharmacogenes.

Gene	Panel AMP Tier 1 compliance	Validated panel compliance*	SIF-AIOM panel compliance [§]	Testing setting			Methodology mainly utilized	Guidelines for interpretation	
				Pre-emptive	Semi-preemptive	Pre-treatment		SIF-AIOM	CPIC/DPWG
DPYD	0%	93%	81%	0%	2%	100%	19%	90%	5%
UGT1A1	-	59%	32%	0%	3%	97%	8%	65%	27%
TPMT	91%	91%	-	9%	0%	91%	0%	-	91%
CYP2C19	63%	63%	-	13%	0%	75%	38%	-	88%
CYP2C9	17%	67%	-	17%	0%	83%	17%	-	83%
CYP2D6	20%	20%	-	20%	0%	20%	60%	-	80%
HLA-B	-	80%	-	20%	20%	60%	0%	-	60%

For each gene, the table summarizes AMP Tier 1 compliance, adherence to panels proposed by different pharmacogenetic guidelines (SIF-AIOM, CPIC, DPWG), predominant testing settings (pre-emptive, semi-preemptive, pre-treatment, reactive), most commonly used methodology, and pharmacogenetic guideline adherence for the results interpretation.

*Validated according to either CPIC, DPWG, or SIF-AIOM guidelines.

[§] Only for DPYD and UGT1A1.

the decentralized structure of the Italian National Health System. Within the Italian reimbursement system, (pharmaco)genetic testing is reimbursed according to procedural codes (e.g., single-gene tests, gene panels, or NGS tests), rather than on an individual gene-by-gene basis. Several laboratories also reported the absence of dedicated reimbursement codes specific to pharmacogenetic testing, which may contribute to heterogeneous administrative and reimbursement practices across institutions.

DISCUSSION

The application of pharmacogenetics to pharmacological treatment management has seen a significant advancement in the last few years; however, its clinical implementation appears to be still challenged by a poor harmonization of clinical and laboratory procedures as well as a still deficient regulatory framework in the field [7,8,10]. This nationwide survey provides the first comprehensive overview of pharmacogenetic testing practices in Italy. The results reveal a substantial heterogeneity in service organization, testing procedures, and clinical integration across different institutions as well as a high plethora of educational and harmonization needs expressed by professionals in the field.

Pharmacogenetic testing emerged as predominantly applied in the oncology therapeutic area, particularly for *DPYD* and *UGT1A1* in support of fluoropyrimidine and irinotecan treatment. Regulatory initiatives have played a central role in promoting clinical implementation in this context. In 2020, the European Medicines Agency (EMA) recommended pre-treatment *DPYD* testing [13], and the Italian Medicines Agency (AIFA) adopted this guidance [14], formally recognizing the clinical utility of both *DPYD* and *UGT1A1* testing. These developments have been facilitated by the availability of national guidelines jointly issued by SIF and AIOM, which promote standardized genotyping and dosing strategies, with a well established clinical utility supported by recent studies [15].

However, national pharmacogenetic guidance remains limited outside the oncological field. Expanding national recommendations to other gene-drug pairs could help define testing priorities and support broader implementation. Moreover, the involvement of medical professional societies, as done for the *DPYD/UGT1A1* national guidelines, developed in a collaboration between pharmacologists and oncologists, can boost the awareness of the value of pharmacogenetics among medical practitioners, thereby facilitating the translation of pharmacogenetic guidelines into improved clinical practices. Continued collaboration among national professional societies, along with the alignment to international frameworks like CPIC and DPWG, will be essential to advance pharmacogenetics integration across therapeutic areas.

Compared with similar surveys conducted in other European countries [4,12,16,17], the Italian pharmacogenetic ecosystem appears less standardized and more fragmented, particularly regarding testing panels, software tools, and result interpretation. The marked heterogeneity in testing platforms and interpretation software further highlights the absence of harmonized technical standards. The use of 18 different bioinformatics tools, along with the coexistence of numerous CE-IVD and non-certified kits, may lead to variability in genotyping accuracy and phenotype prediction. Developing consensus on validated assays, reference variant lists, and interpretation pipelines would therefore represent a critical step toward improving reproducibility and clinical reliability across sites. Moreover, while most laboratories primarily rely on targeted panels, an increasing number of groups are shifting toward broader approaches such as NGS, which enable a more comprehensive assessment of individual pharmacogenetic profiles. This evolution may facilitate the detection of rare or previously untested variants, thereby expanding the clinical scope of pharmacogenetics, an approach that has already shown




Fig. 3 Characteristics of laboratories performing pharmacogenetic testing and overview of survey feedback. **A** Type of unit performing pharmacogenetic testing. Bars indicate the number of laboratories per unit type; the line shows the percentage of laboratories providing pharmacological counseling within each type of unit. **B** Overview of feedback and annotations submitted by laboratories participating in the survey, grouped by thematic area.




Fig. 4 Map of pharmacogenetic laboratories across Italy. Background colors represent geographic macro-areas, and circle size indicates the average number of pharmacogenetic tests performed per laboratory, per year.

potential clinical advantages [18–20]. However, the transition toward such technologies also underscores the need for enhanced bioinformatics infrastructures and more standardized analysis pipelines [21], as also highlighted by feedback collected in the survey. An opportunity in this context is offered in oncology, where repurposing NGS data on tumor tissue generated within Molecular Tumor Boards (MTBs) could represent a cost-effective opportunity. The germline component of tumor–normal sequencing is routinely produced but remains largely underutilized for pharmacogenetic purposes, despite its potential to support personalized treatment decisions [22].

At both national and international levels, several consortia have emerged to standardize and harmonize pharmacogenomics laboratory procedures, in order to ensure analytical consistency, clinical reliability, and cross-laboratory comparability. Among these, the AMP has developed comprehensive recommendations defining the minimum variant sets (Tier 1 panels) required for accurate genotype-to-phenotype translation. These panels were designed according to criteria such as allele frequency across multiple populations, evidence of functional impact, and clinical validity supported by peer-reviewed studies, thereby promoting the inclusion of variants across diverse ethnic groups [23–26].

In parallel, international expert bodies such as the CPIC and the DPWG have focused on the clinical interpretation of pharmacogenetic results, providing evidence-based, gene–drug–specific recommendations for dose adjustment or therapeutic selection [27]. Together, these initiatives constitute a complementary framework: while AMP establishes the analytical foundation to ensure that pharmacogenetic assays capture clinically relevant variants, CPIC and DPWG translate these data into actionable clinical guidance.

A key finding concerns the inconsistency in the genetic content of pharmacogenetic panels used across laboratories. The present survey highlights that AMP compliance is achieved for some pharmacogenes (e.g., *TPMT*), but remains low or variable for others (*CYP2C9*, *CYP2D6*, *DPYD*), underscoring the need for consensus on variant selection and standardized analytical validation criteria, and highlighting once more how the availability of clinical guidelines mediated by the national societies remains the major reference for the laboratory activity.

Testing settings also showed important trends. Oncology-related genes such as *DPYD* and *UGT1A1* are almost exclusively tested pre-treatment, in line with recent studies and publications that support the positive impact of preemptive genotyping on treatment outcomes [2].

Conversely, semi-preemptive strategies are emerging for *CYP2C19* and *HLA-B*, indicating a gradual shift toward broader clinical implementation. This higher prevalence of semi-preemptive testing for *CYP2C19* and *HLA-B* may be due to their frequent inclusion in broader pharmacogenetic panels, which are often performed independently of an immediate therapeutic decision but can still provide clinical benefit later. However, reactive testing—performed only after an adverse event or treatment failure—remains prevalent for genes like *CYP2D6*, suggesting that routine pre-emptive or panel-based approaches are not yet integrated into Italian clinical workflows.

The predominance of genetics and clinical pathology units in performing pharmacogenetic tests likely reflects the historical evolution of molecular diagnostics within the Italian healthcare system. However, the limited involvement of pharmacology units—despite their expertise in dose adjustment and therapy optimization—suggests a need to strengthen interdisciplinary collaboration. The relatively low proportion of laboratories providing explicit pharmacological counseling (24%) underscores this gap. Notably, pharmacological counseling is most often provided by pharmacology units, which, however, are limited in number and unevenly distributed across the country. This highlights the potential value of establishing a national network of specialized laboratories and experts to provide pharmacological

support in the interpretation of pharmacogenetic data. Furthermore, concentrating pharmacogenetic analyses in accredited genetic laboratories could ensure high-quality, standardized testing while enabling hub pharmacological centers to support spoke healthcare facilities with comprehensive pharmacological consultations.

A direct consequence of this fragmented landscape is the considerable variability observed across the different regions of the country. Testing activity was indeed found to be concentrated in the northern regions, with marked geographical disparities in availability and volume, as well as variable access to pharmacological counseling and reimbursement mechanisms. At the system level, the uneven distribution of pharmacogenetic services in the territory, with the majority of high-volume laboratories located in Northern Italy, raises concerns about equitable access to testing nationwide. Similarly, while most laboratories report full or partial reimbursement by the National Health Service, differences in coverage across regions contribute to inconsistent integration of pharmacogenetics into routine care. This clearly reflects the organization of the Italian health care system, which delegates to individual regions the responsibility for defining many aspects of the healthcare delivery, including the allocation of resources, laboratory accreditation, and reimbursement mechanisms. While this regional autonomy can promote flexibility and innovation, it also results in marked heterogeneity in the implementation of pharmacogenetic testing and in patient access to these services. A coordinated national framework is urgently needed to address these disparities, by defining minimum testing requirements, accreditation criteria, and pathways for sustainable reimbursement.

The high rate of responders providing feedback on the needs to advance pharmacogenetics implementation in Italy demonstrates strong interest and motivation among professionals in the field. All the feedback received from the laboratories participating in the survey can be summarized as highlighting the need for structured training and networking opportunities to support the implementation of pharmacogenetics in clinical practice. This clearly underscores the urgent need for dedicated training and education programs in pharmacogenetics. Consistent with other surveys, the lack of specific pharmacogenetic training is recognized as one of the significant barriers to its widespread clinical implementation [4,12,16,17]. To address this gap, ongoing initiatives promoting pharmacogenetics in Italy should be actively supported and coordinated by scientific organizations such as the Italian Society of Pharmacology (SIF) and the Italian Society of Human Genetics (SIGU) to foster education, standardization, and interdisciplinary collaboration across clinical and research settings.

Overall, the insights gained from this survey provide a foundation for the activities of the *Joint Working Group for the Implementation of Pharmacogenetics in Italy*. The next steps should include the establishment of a permanent national network of laboratories, the harmonization of testing and reporting standards, and the promotion of education and awareness initiatives for clinicians and healthcare stakeholders. Enhanced collaboration between geneticists, pharmacologists, and clinicians will be essential to embed pharmacogenetic testing into patient-centered therapeutic strategies, to align Italy with international best practices.

The main strengths of this study lie in its nationwide coverage and high response rate, providing a representative picture of pharmacogenetic laboratory activity across Italy. Limitations include its self-reported design and the possibility of participation bias, as laboratories with more structured pharmacogenetic programs may have been more inclined to respond.

In conclusion, this survey establishes the first national map of pharmacogenetic services in Italy. It highlights key challenges—heterogeneity, fragmentation, and regional disparities—but also substantial potential and motivation for integration, improvement, and potential development of appropriate health care pathways.

Building on these findings, a coordinated national strategy could foster the harmonization of practices, strengthen the link between genotyping and clinical decision-making, and ultimately enhance patient outcomes through personalized therapy.

DATA AVAILABILITY

Data supporting the findings of this study are available within the article. Additional information is available from the corresponding author upon reasonable request.

REFERENCES

- Minikel EV, Painter JL, Dong CC, Nelson MR. Refining the impact of genetic evidence on clinical success. *Nature*. 2024;629:624–9.
- Swen JJ, van der Wouden CH, Manson LE, Abdullah-Koolmees H, Blagec K, Blagus T, et al. A 12-gene pharmacogenetic panel to prevent adverse drug reactions: an open-label, multicentre, controlled, cluster-randomised crossover implementation study. *Lancet*. 2023;401:347–56.
- Bignucolo A, De Mattia E, Roncato R, Peruzzi E, Scarabel L, D'Andrea M, et al. Ten-year experience with pharmacogenetic testing for DPYD in a national cancer center in Italy: Lessons learned on the path to implementation. *Front Pharmacol*. 2023;14:1199462.
- de With M, Sadlon A, Cecchin E, Haufroid V, Thomas F, Joerger M, et al. Implementation of dihydropyrimidine dehydrogenase deficiency testing in Europe. *ESMO Open*. 2023;8:101197.
- PharmGKB [Internet]. [cited 2024 Feb 9]. Clinical Guideline Annotations. Available from: <https://www.pharmgkb.org/guidelineAnnotations>.
- Peruzzi E, Roncato R, De Mattia E, Bignucolo A, Swen JJ, Guchelaar HJ, et al. Implementation of pre-emptive testing of a pharmacogenomic panel in clinical practice: where do we stand? *Br J Clin Pharmacol*. 2023;91:270–82.
- Floris M, Moschella A, Alcalay M, Montella A, Tirelli M, Fontana L, et al. Pharmacogenetics in Italy: current landscape and future prospects. *Hum Genomics*. 2024;18:78.
- Moschella A, Mourou S, Perfler S, Zoroddu E, Bezzini D, Soru D, et al. Pharmacogenetic information on drug labels of the Italian agency of medicines (AIFA): actionability and comparison across other regulatory agencies. *Clin Transl Sci*. 2025;18:e70138.
- Abou Diwan E, Zeitoun RI, Abou Haidar L, Cascorbi I, Khoueiry Zgheib N. Implementation and obstacles of pharmacogenetics in clinical practice: An international survey. *Br J Clin Pharmacol*. 2019;85:2076–88.
- Roncato R, Perfler S, Pasin D, Peruzzi E, Woodcock M, Thomas H, et al. Uptake of DPYD and UGT1A1 testing in Italy and adherence to pharmacogenetic guidelines: A 5-year perspective from an EQA provider. *Br J Clin Pharmacol*. 2025;92:445–56.
- Roncato R, Dal Cin L, Mezzalana S, Comello F, De Mattia E, Bignucolo A, et al. FARMAPRICE: a pharmacogenetic clinical decision support system for precise and cost-effective therapy. *Genes (Basel)*. 2019;10:276.
- Shaunak N, Keen J, Kim A, Pirmohamed M, Newman WG, Ross PJ. Implementation of mass pharmacogenetic testing: dihydropyrimidine dehydrogenase testing prior to fluoropyrimidine treatment for patients. *Br J Clin Pharmacol*. 2025;91:2534–42.
- EMA & fluoropyrimidines [Internet]. Available from: www.ema.europa.eu/en/documents/press-release/ema-recommendations-dpd-testing-prior-treatment-fluorouracil-capecitabine-tegafur-and-flucytosine_en.pdf.
- AIFA & fluoropyrimidines [Internet]. Available from: www.aifa.gov.it/documents/20142/1097058/2020.05.25_NII_5-FU-capecitabina-tegafur.pdf.
- Roncato R, Bignucolo A, Peruzzi E, Montico M, De Mattia E, Foltran L, et al. Clinical benefits and utility of pretherapeutic DPYD and UGT1A1 testing in gastrointestinal cancer: a secondary analysis of the PREPARE randomized clinical trial. *JAMA Netw Open*. 2024;7:e2449441.
- Wittwer NL, Meier CR, Huber CA, Meyer Zu Schwabedissen HE, Allemann S, Schneider C. Pharmacogenetic testing in the outpatient setting in Switzerland: A descriptive study using Swiss claims data. *Drug Saf*. 2025;48:689–96.
- Klein CJ, Gong L, Caudle KE, Naik H, Empey PE, Hoffman JM, et al. Toward an integrated resource for pharmacogenomics (PGx): Survey findings from the genomic medicine communities. *Genet Med*. 2025;27:101529.
- De Mattia E, Milan N, Assaraf YG, Toffoli G, Cecchin E. Clinical implementation of rare and novel DPYD variants for personalizing fluoropyrimidine treatment: challenges and opportunities. *Int J Biol Sci*. 2024;20:3742–59.
- De Mattia E, Silvestri M, Polesel J, Ecça F, Mezzalana S, Scarabel L, et al. Rare genetic variant burden in DPYD predicts severe fluoropyrimidine-related toxicity risk. *Biomed Pharmacother*. 2022;154:113644.
- De Mattia E, Polesel J, Silvestri M, Roncato R, Scarabel L, Calza S, et al. The burden of rare variants in DPYD gene is a novel predictor of the risk of developing severe fluoropyrimidine-related toxicity. *Hum Genomics*. 2023;17:99.
- Lauschke VM, Ingelman-Sundberg M. Emerging strategies to bridge the gap between pharmacogenomic research and its clinical implementation. *NPJ Genom Med*. 2020;5:9.
- Koo H, Smith TB, Callaghan JT, Osei W, Bray SM, Tillman EM, et al. Return of clinically actionable pharmacogenetic results from molecular tumor board DNA sequencing data: workflow and estimated costs. *Clin Pharm Ther*. 2025;117:1017–20.
- Pratt VM, Cavallari LH, Fulmer ML, Gaedigk A, Hachad H, Ji Y, et al. DPYD Genotyping Recommendations. *J Mol Diagn*. 2024;26:851–63.
- Pratt VM, Cavallari LH, Del Tredici AL, Hachad H, Ji Y, Moyer AM, et al. Recommendations for clinical CYP2C9 genotyping allele selection. *J Mol Diagnostics*. 2019;21:746–55.
- Pratt VM, Del Tredici AL, Hachad H, Ji Y, Kalman LV, Scott SA, et al. Recommendations for clinical CYP2C19 genotyping allele selection. *J Mol Diagn*. 2018;20:269–76.
- Pratt VM, Cavallari LH, Del Tredici AL, Gaedigk A, Hachad H, Ji Y, et al. Recommendations for clinical CYP2D6 genotyping allele selection. *J Mol Diagn*. 2021;23:1047–64.
- ClinPGx [Internet]. [cited 2025 Oct 27]. ClinPGx. Available from: <https://www.clinpgx.org/>.

AUTHOR CONTRIBUTIONS

RR, MF and EC conceptualized and designed the study. RR, MF, SP, MG, EZ and EC carried out interpretation of data. RR, MF, SP, MG, EZ and EC designed the questionnaire and wrote the manuscript. SP, MG, EZ, MRM, AS, SA, PB, MGE, GN, VC, AF, GS, MF and EC edited and revised the questionnaire and the manuscript. All authors read and approved the final manuscript.

FUNDING

The author(s) declare that financial support was received for the research, authorship, and/or publication of this article. EC, SP, MG were supported by the Italian Ministry of Health (Ricerca Corrente). Open access funding provided by Università degli Studi di Sassari within the CRUI-CARE Agreement.

COMPETING INTERESTS

The authors declare no competing interests.

ETHICAL APPROVAL AND INFORMED CONSENT

In accordance with national and institutional guidelines, ethical approval was not required for this study, as it did not involve human participants, patients, clinical data, or identifiable personal information. The survey collected only laboratory-level organizational data from laboratory directors.

ADDITIONAL INFORMATION

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41431-026-02063-4>.

Correspondence and requests for materials should be addressed to Matteo Floris or Erika Cecchin.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2026

THE JOINT ITALIAN WORKING GROUP ON PHARMACOGENETICS IMPLEMENTATION

Myriam Alcalay^{13,14}, Manuela Badioli¹⁵, Marco Benati^{16,17}, Francesca Daniela Bertola¹⁸, Gabriella Bianchino¹⁹, Elisa Boldrin²⁰, Elisa Bonomi²¹, Beatrice Bortesi²², Concetta Cafiero²³, Daniele Calistri²⁴, Mario Capasso^{25,26}, Anna Paola Capra^{27,28}, Rita Congiu¹⁵, Francesca Crosti¹⁸, Matteo Curtarello²⁰, Jessica Cusato²⁹, Giulia D'Addario³⁰, Alessandra D'Urso³¹, Emanuela De Bellis³², Dario De Biase^{5,33}, Giovanna De Matteis^{16,17}, Maria Dono³⁴, Matteo Fabbri³⁵, Antonella Fabretto⁸, Mosè Favarato³⁶, Stefano Fogli³⁷, Alessandra Franzoni³⁸, Luca Gallelli³⁹, Sabrina Rita Giglio^{40,41,42,43}, Mara Giordano⁴⁴, Giorgia Girotto⁴⁵, Paola Grammatico⁴⁶, Cristina Lapucci⁴⁷, Sonia Lastraioli³⁴, Barbara Mancini⁴⁸, Irene Mancini⁴⁹, Simona Natalina Mellone⁴⁴, Pierantonio Menna⁵⁰, Roberta Minari²², Rossana Mineri⁵¹, Giuseppe Miscio⁵², Lorena Mosca⁵³, Antonino Moschella⁵⁴, Stefania Murru¹⁵, Davide Nicolì⁵⁵, Sara Orecchia⁵⁶, Gloria Pessina⁵⁷, Claudia Rengucci²⁴, Nicoletta Sacchi⁵⁸, Federica Sangiuolo¹¹, Maria Scatolini⁵⁹, Lisa Simi⁴⁹, Sabine Stioui⁶⁰, Silvia Tabano^{61,62}, Luisa Toffolatti⁶³, Stefania Tommasi⁶⁴, Davide Trerè^{65,66}, Alessandro Ubiali⁶⁷, Sheila Ulivi⁴⁵ and Monica Zanussi⁶⁸

¹³Department of Experimental Oncology, IEO, European Institute of Oncology IRCCS, Milano, Italy. ¹⁴Department of Oncology and Hemato-Oncology, University of Milano, Milan, Italy. ¹⁵Laboratorio di Genetica e Genomica P.O. Microcitemico A.Cao ASL Cagliari, Cagliari, Italy. ¹⁶Section of Clinical Biochemistry, University of Verona, Verona, Italy. ¹⁷UOC Laboratorio Analisi Azienda Ospedaliera universitaria integrata Verona, Verona, Italy. ¹⁸IRCCS Fondazione San Gerardo dei Tintori, SC Genetica Medica, Monza, Italy. ¹⁹Unit of Clinical Pathology IRCCS-CROB Referral Cancer Center of Basilicata, Rionero in Vulture (PZ), Italy. ²⁰Laboratorio Farmacogenetica—UOC Immunologia e Diagnostica Molecolare Oncologica, Istituto Oncologico Veneto (IOV), Padova, Italy. ²¹S.S.D. Laboratorio Genetica Medica ASST Spedali Civili di Brescia, Brescia, Italy. ²²Medical Oncology Unit, University Hospital of Parma, Parma, Italy. ²³Medical Oncology, S.G. Moscati, Statte (TA), Italy. ²⁴Biosciences Laboratory, IRCCS Istituto Romagnolo Per Lo Studio Dei Tumori “Dino Amadori”, Meldola (FC), Italy. ²⁵Dipartimento di Medicina Molecolare e Biotecnologie Mediche, Università degli Studi di Napoli Federico II, Napoli, Italy. ²⁶CEINGE Biotecnologie Avanzate Franco Salvatore, Napoli, Italy. ²⁷Department of Chemical, Biological, Pharmaceutical and Environmental Sciences, University of Messina, Messina, Italy. ²⁸Genetics and Pharmacogenetics Unit, “Gaetano Martino” University Hospital, Messina, Italy. ²⁹Laboratory of Clinical Pharmacology and Pharmacogenetics, Department of Medical Sciences, University of Turin, Turin, Italy. ³⁰Centro Ricerca Clinica Ospedale Isola Tiberina Gemelli Isola Roma, Rome, Italy. ³¹Pathology Unit, AULSS1 Dolomiti, Feltre (BL), Italy. ³²PhD School “Clinical and Translational Oncology (CTO)”, Scuola Superiore Meridionale, University of Naples “Federico II”, Napoli, Italy. ³³Solid Tumor Molecular Pathology Laboratory, IRCCS Azienda Ospedaliero-Universitaria di Bologna, Bologna, Italy. ³⁴S.S. Diagnostica Molecolare, IRCCS Ospedale Policlinico San Martino Genova, Genova, Italy. ³⁵Università degli Studi di Ferrara, Dipartimento di Medicina Traslazionale, Sezione dipartimentale di medicina legale, Ferrara, Italy. ³⁶UOSD Genetica e Citogenetica Aulss3 Serenissima, Mestre (VE), Italy. ³⁷Unit of Clinical Pharmacology and Pharmacogenetics, Department of Clinical and Experimental Medicine, University of Pisa, Pisa, Italy. ³⁸Institute of Human Genetics, Santa Maria della Misericordia University Hospital, Azienda Sanitaria Universitaria Friuli Centrale (ASUFC), Udine, Italy. ³⁹Department of Health Science, University of Catanzaro, Catanzaro, Italy. ⁴⁰Medical Genetics, R. Binaghi Hospital, ASSL Cagliari, ATS Sardegna, Cagliari, Italy. ⁴¹Centre for Research University Services, University of Cagliari, Monserrato (CA), Italy. ⁴²AART-ODV (Association for the Advancement of Research on Transplantation), Cagliari, Italy. ⁴³Medical Genetics, Department of Medical Sciences and Public Health, University of Cagliari, Cagliari, Italy. ⁴⁴Unit of Genetics, Clinical Biochemistry, AOU Maggiore della Carità, Novara, Italy. ⁴⁵Genetica Medica, Institute for maternal and child health IRCCS Burlo Garofalo, Trieste, Italy. ⁴⁶Laboratory of Medical Genetics, Department of Experimental Medicine, Sapienza University, San Camillo-Forlanini Hospital, Roma, Italy. ⁴⁷Bianalisi, Carate Brianza (MB), Roma, Italy. ⁴⁸U.O.S.D. Genetica Medica e Genomica—AULSS8 Berica, Vicenza, Italy. ⁴⁹SOD Lab. Biochimica Clinica Molecolare Azienda Ospedaliero-Universitaria Careggi, Firenze, Italy. ⁵⁰Department of Science and Technology for Sustainable Development and One Health, Campus Bio-Medico University, and Unit of Clinical Pharmacology, Fondazione Policlinico Universitario Campus Bio-Medico, Roma, Italy. ⁵¹Department of Laboratory Medicine, IRCCS Humanitas Research Hospital, Rozzano (MI), Italy. ⁵²Fondazione IRCCS Casa Sollievo della Sofferenza, San Giovanni Rotondo (FG), Italy. ⁵³Medical Genetics Unit, Department of Medical Services, ASST Grande Ospedale Metropolitano Niguarda, Milano, Italy. ⁵⁴UOSD Genetica Medica, Grande Ospedale Metropolitano “Bianchi-Melacrino-Morelli”, Reggio Calabria, Italy. ⁵⁵Molecular Pathology, Azienda USL-IRCCS di Reggio Emilia, Reggio Emilia, Italy. ⁵⁶Molecular Pathology Unit, SS Antonio and Biagio Hospital, Alessandria, Italy. ⁵⁷UOSD Laboratorio di Genetica Medica, Viterbo, Italy. ⁵⁸Italian Bone Marrow Donor Registry, EO Ospedali Galliera di Genova, Genova, Italy. ⁵⁹Molecular Oncology Laboratory, Fondazione Edo ed Elvo Tempia, Ponderano (BI), Italy. ⁶⁰Laboratorio di Genetica Medica Centro Diagnostico Italiano Milano Italia, Milano, Italy. ⁶¹Dipartimento di Fisiopatologia Medico-Chirurgica e dei Trapianti, Università degli Studi di Milano, Milano, Italy. ⁶²Laboratorio di Genetica Medica, Fondazione IRCCS Ca’ Granda Ospedale Maggiore Policlinico, Milano, Italy. ⁶³Laboratorio di Patologia Molecolare, Ospedale Ca’ Foncello Treviso, Italia, Treviso, Italy. ⁶⁴Molecular Diagnostics and Pharmacogenetics Unit, IRCCS Istituto Tumori “Giovanni Paolo II”, Bari, Italy. ⁶⁵IRCCS Azienda Ospedaliero-Universitaria di Bologna, Bologna, Italy. ⁶⁶Department of Medical and Surgical Sciences (DIMEC), University of Bologna, Bologna, Italy. ⁶⁷Pathology Unit, AUSL Piacenza, Piacenza, Italy. ⁶⁸Laboratory of Clinical Genomics, IRCCS San Raffaele Scientific Institute, Milano, Italy.