



**UNIVERSITÀ DEGLI STUDI DI ROMA
"TOR VERGATA"**

FACOLTA' DI MEDICINA E CHIRURGIA

DOTTORATO DI RICERCA IN
MICROBIOLOGIA MEDICA E IMMUNOLOGIA

- XXI CICLO -

*“Activated T lymphocytes instruct monocytes to differentiate into
inflammatory dendritic cells in a feedback control of human
Th1/Th2 response”*

Dott.^{ssa} Valeria Sargentini

A.A. 2008/2009

Tutor: Dott. Roberto Nisini

Coordinatore: Prof. Enrico Garaci

TABLE OF CONTENTS

ABSTRACT	Pag. 5
1. INTRODUCTION	Pag. 6
1.1 Immune system	Pag. 7
1.1.1 Innate immune system	Pag. 7
1.1.2 Adaptive immune system	Pag. 9
1.2 T lymphocytes	Pag. 10
1.3 Dendritic cells (DCs)	Pag. 13
1.3.1 Activation of DCs and launching of protective immunity	Pag. 13
1.3.2 DC types and subtypes	Pag. 18
1.3.3 DCs in the health/disease balance	Pag. 20
1.4 The origin of DCs	Pag. 22
1.4.1 The origin of steady-state DCs	Pag. 23
1.4.2 Differentiation of conventional DCs locally in the peripheral tissues	Pag. 27
1.5 Monocytes as DCs precursors	Pag. 29
1.5.1 Monocyte subsets	Pag. 29
1.5.2 Monocyte-derived inflammatory DCs in the mouse models	Pag. 30
1.5.3 Differentiation of human monocytes into DCs	Pag. 33
1.6 Cytokines in the generation and maturation of monocytes derived DCs	Pag. 34
2. AIM OF THE STUDY	Pag. 39
3. MATERIALS AND METHODS	Pag. 42
3.1 Reagents	Pag. 43
3.2 Generation and maintenance of antigen specific human T cell line and T cell clones (TCC)	Pag. 43
3.3 Monocytes differentiation experiments	Pag. 45
3.3.1 Monocytes isolation and differentiation into “reference”	

<i>DCs</i>	Pag.	45
3.3.2 <i>DCs differentiation from monocytes co-cultured with TCC</i>	Pag.	46
3.3.3 <i>Transwell culture system</i>	Pag.	47
3.3.4 <i>DCs differentiation from monocytes cultured with activated TCC supernatants</i>	Pag.	48
3.4 Flow cytometry and immunostaining	Pag.	48
3.5 Microscopic evaluation of DCs	Pag.	49
3.6 Phagocytosis assay	Pag.	49
3.7 Endocytosis assay	Pag.	50
3.8 Isolation of naïve CD4 ⁺ T cells	Pag.	50
3.9 Priming of naïve T cells	Pag.	51
3.10 Antigen presentation assay	Pag.	51
3.11 Enzyme Linked ImmunoSorbent Assay (ELISA)	Pag.	51
3.12 Cytotoxic T-cell line (CTL2) assay	Pag.	52
3.13 Real time Polymerase Chain Reaction (PCR)	Pag.	53
3.14 Statistical analysis	Pag.	53
4. RESULTS	Pag.	54
4.1 Characterization of antigen-specific TCC with different functional polarization	Pag.	55
4.2 Specific T cell activation induces antigen-presenting monocytes to differentiate into DCs	Pag.	59
4.3 T cell-dependent monocytes differentiation into DCs does not require cell-to-cell contact	Pag.	61
4.4 Different TCC subpopulations induce monocytes to differentiate into DCs with distinct phenotypes	Pag.	61
4.4.1 <i>Morphologic characteristics of different DC subtypes</i>	Pag.	61
4.4.2 <i>Surface expression of group I CD1 molecules</i>	Pag.	64
4.4.3 <i>DC-SIGN expression</i>	Pag.	66
4.4.4 <i>Larger phenotypic characterization of DCs derived from monocytes cultured with activated Th2, Th1 or Th0 PPD specific TCC</i>	Pag.	67
4.4.5 <i>CD1a-reduced expression on DCh1 and their semi-mature</i>		

<i>phenotype is related to IL-4 absence and/or IFN-γ secretion</i>	Pag.	67
4.4.6 <i>Freshly isolated T lymphocytes ex-vivo induce monocytes to differentiate into DCs</i>	Pag.	70
4.5 DCs differentiated upon T cell activation show a differential cytokine secretion pattern	Pag.	71
4.6 DCs induced by activated T lymphocytes show different degree of bacterial phagocytosis and soluble antigen uptake	Pag.	72
4.7 DCh1 and DCh0 have a reduced capacity to prime naïve CD4 ⁺ T lymphocytes	Pag.	73
4.8 DCh1 and DCh0 have a reduced capacity to activate antigen-specific memory T-cell response	Pag.	78
5. DISCUSSION	Pag.	80
6. ABBREVIATIONS	Pag.	87
7. REFERENCES	Pag.	91
8. ACKNOWLEDGEMENTS	Pag.	109