


This article was downloaded by: [Flamini, Flaminio]

On: 3 December 2008

Access details: Access Details: [subscription number 906309347]

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Communications in Algebra

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597239>

Nodal Curves with General Moduli on K3 Surfaces

Flaminio Flamini ^a; Andreas Leopold Knutsen ^b; Gianluca Pacienza ^c; Edoardo Sernesi ^d

^a Department of Mathematics, University of Studies "Tor Vergata" of Rome, Rome, Italy ^b Department of Mathematics, University of Bergen, Bergen, Norway ^c Institute of Advanced Mathematical Research, University L. Pasteur and CNRS, Strasbourg, France ^d Department of Mathematics, University of Studies Roma Tre, Rome, Italy

Online Publication Date: 01 November 2008

To cite this Article Flamini, Flaminio, Knutsen, Andreas Leopold, Pacienza, Gianluca and Sernesi, Edoardo(2008)'Nodal Curves with General Moduli on K3 Surfaces', Communications in Algebra, 36:11, 3955 — 3971

To link to this Article: DOI: 10.1080/00927870802174082

URL: <http://dx.doi.org/10.1080/00927870802174082>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

NODAL CURVES WITH GENERAL MODULI ON K3 SURFACES

Flaminio Flamini¹, Andreas Leopold Knutsen²,
Gianluca Pacienza³, and Edoardo Sernesi⁴

¹Department of Mathematics, University of Studies
“Tor Vergata” of Rome, Rome, Italy

²Department of Mathematics, University of Bergen, Bergen, Norway

³Institute of Advanced Mathematical Research,
University L. Pasteur and CNRS, Strasbourg, France

⁴Department of Mathematics, University of Studies Roma Tre, Rome, Italy

We investigate the modular properties of nodal curves on a low genus K3 surface. We prove that a general genus g curve C is the normalization of a δ -nodal curve X sitting on a primitively polarized K3 surface S of degree $2p - 2$, for $2 \leq g = p - \delta < p \leq 11$. The proof is based on a local deformation-theoretic analysis of the map from the stack of pairs (S, X) to the moduli stack of curves \mathcal{M}_g that associates to X the isomorphism class $[C]$ of its normalization.

Key Words: K3 surfaces; Moduli spaces; Nodal curves.

2000 Mathematics Subject Classification: Primary 14H10, 14H51, 14J28; Secondary 14C05, 14D15.

1. INTRODUCTION

Nonsingular curves of low genus on a K3 surface have interesting modular properties, related to the existence of Fano 3-folds of index one of the corresponding sectional genus. These properties have been investigated by Mukai who settled, in particular, a problem raised in Mayer (1972). He showed that a general curve of genus $g \leq 9$ or $g = 11$ can be embedded as a nonsingular curve in a K3 surface, and that this is not possible for curves of genus $g = 10$, despite an obvious count of constants indicating the opposite. These facts have been considered again by Beauville in the last section of Beauville (2004) from a different point of view, by means of a local deformation-theoretic analysis.

In the present article, we take a point of view similar to Beauville's with the purpose of studying the corresponding questions about moduli of *singular (nodal) curves* of low genus on a K3 surface. To this end we consider the following algebraic stacks:

Received June 10, 2007. Communicated by L. Ein.

Address correspondence to Flaminio Flamini, Dipartimento di Matematica, Università di Roma “Tor Vergata”, Viale della Ricerca Scientifica, I-00133 Roma, Italy; Fax: +39-06-72594699; E-mail: flamini@mat.uniroma2.it

- \mathcal{B}_p : The stack of smooth $K3$ surfaces S marked by a globally generated, primitive line bundle H of sectional genus $p \geq 2$; it is smooth and irreducible of dimension 19;
- $\mathcal{V}_{p,\delta}$: The stack of pairs (S, X) such that $(S, H) \in \mathcal{B}_p$ and $X \in |H|$ is an irreducible curve with δ nodes and no other singularities, for given $0 \leq \delta \leq p$; it is smooth of dimension $19 + g$, where $g = p - \delta$.

We also consider an étale atlas $V_{p,\delta} \rightarrow \mathcal{V}_{p,\delta}$ and the morphisms

$$\begin{array}{ccc} V_{p,\delta} & \xrightarrow{c_{p,\delta}} & \mathcal{M}_g \\ \downarrow \pi_\delta & & \\ \mathcal{B}_p & & \end{array}$$

where \mathcal{M}_g is the moduli stack of nonsingular curves of genus $g = p - \delta$; $c_{p,\delta}$ and π_δ are induced by associating to a point parametrizing a pair (S, X) the isomorphism class of the normalization of X and $[S]$, respectively.

We study this configuration when $3 \leq p \leq 11$. Our main result is the following theorem.

Theorem 1.1. *Let $3 \leq p \leq 11$ and $0 \leq \delta \leq p - 2$, so that $2 \leq g = p - \delta \leq p$. Let $V \subset V_{p,\delta}$ be an irreducible component, and let*

$$c_{p,\delta|V} : V \longrightarrow \mathcal{M}_g$$

be the restriction to V of the morphism $c_{p,\delta}$. Then, for any $2 \leq g < p \leq 11$, $c_{p,\delta|V}$ is dominant. In particular, for the general fibre $F_{p,\delta}$ of $c_{p,\delta|V}$ we have

$$\dim(F_{p,\delta}) = 22 - 2g = \exp \dim(F_{p,\delta}).$$

The theorem is proved by studying deformations of a pair (S, X) in $\mathcal{V}_{p,\delta}$. The locally trivial deformation theory of such a pair is controlled by a locally free sheaf of rank 2, namely, the sheaf $\mathcal{T}_S\langle X \rangle$ of tangent vectors to S that are tangent to X . Specifically, $H^1(S, \mathcal{T}_S\langle X \rangle)$ is the tangent space to $\mathcal{V}_{p,\delta}$ at (S, X) and $H^2(S, \mathcal{T}_S\langle X \rangle)$ is an obstruction space. These cohomology groups are studied by pulling back $\mathcal{T}_S\langle X \rangle$ to the blowup \tilde{S} of S at the singular points of X . Then the local study of the morphism $c_{p,\delta}$ is carried out on \tilde{S} , and the theorem is reduced to proving the vanishing of an appropriate cohomology group.

The case $p = 11, \delta = 1$ of the theorem appears to be somehow unexpected. Note that, in fact, the theorem says that every irreducible component V of $V_{11,1}$ dominates \mathcal{M}_{10} . This contrasts the fact that, according to Mukai, $V_{10,0}$ does not dominate \mathcal{M}_{10} .

Another interesting case is $p = 10, \delta = 1$. Again the theorem says that every irreducible component of $V_{10,1}$ dominates \mathcal{M}_9 . But, since $V_{10,0}$ is mapped to a divisor of \mathcal{M}_{10} by $c_{10} := c_{10,0}$, it follows that the nodal curves in $V_{10,1}$ only fill a divisor on the boundary $\partial \overline{\mathcal{M}}_{10}$ of $\overline{\mathcal{M}}_{10}$, despite the fact that their normalizations are general curves of genus 9. This means that on a general curve C of genus 9, the effective divisors $P + Q$, with $P \neq Q$, such that the nodal curve $X = C/(P = Q)$ can be embedded in

a K3 surface, belong to a 1-dimensional cycle $\Gamma \subset C^{(2)}$. It would be interesting to compute the numerical class of Γ .

The article consists of 5 sections including the introduction. After recalling the relevant deformation theory in Section 2, we survey the known results about moduli of smooth curves on marked K3 surfaces in Section 3. In Section 4, we develop our approach for the case of nodal curves, and in Section 5, we discuss the existence of nodal curves having normalizations with general moduli. In the end we raise some related open questions.

2. SOME BASIC RESULTS OF DEFORMATION THEORY

In this section, we will review some results on deformation theory that are needed for our aims. For complete details, we refer the reader to, e.g., Sernesi (2006, §3.4.4).

Let Y be a smooth variety, and let $j : X \hookrightarrow Y$ be a closed embedding of a Cartier divisor X . The *locally trivial deformations* of j are studied by means of suitable sheaves on Y .

Let $\mathcal{N}_{X/Y}$ be the normal sheaf of X in Y , and $\mathcal{N}'_{X/Y} \subseteq \mathcal{N}_{X/Y}$ the *equisingular normal sheaf* of X in Y (cf. Sernesi, 2006, Proposition 1.1.9). One can define a coherent sheaf $\mathcal{T}_Y\langle X \rangle$ of rank $\dim(Y)$ on Y via the exact sequence

$$0 \longrightarrow \mathcal{T}_Y\langle X \rangle \longrightarrow \mathcal{T}_Y \longrightarrow \mathcal{N}'_{X/Y} \longrightarrow 0, \tag{2.1}$$

which is called the *sheaf of germs of tangent vectors to Y that are tangent to X* (cf. Sernesi, 2006, §3.4.4). Of course, when X is smooth, then $\mathcal{N}'_{X/Y}$ in (2.1) is nothing but the normal bundle $\mathcal{N}_{X/Y}$.

One has a natural surjective restriction map

$$r : \mathcal{T}_Y\langle X \rangle \longrightarrow \mathcal{T}_X, \tag{2.2}$$

giving the exact sequence

$$0 \longrightarrow \mathcal{T}_Y(-X) \longrightarrow \mathcal{T}_Y\langle X \rangle \longrightarrow \mathcal{T}_X \longrightarrow 0, \tag{2.3}$$

where $\mathcal{T}_Y(-X)$ is the vector bundle of tangent vectors of Y vanishing along X , and where \mathcal{T}_X is the *tangent sheaf* of X , i.e., the dual sheaf of the sheaf of Kähler differentials of X (cf. Sernesi, 2006, §3.4.4).

Observe that, when X is a divisor with simple normal crossings (see Itaka, 1982), $\mathcal{T}_Y\langle X \rangle$ is a locally free subsheaf of the holomorphic tangent bundle \mathcal{T}_Y , whose restriction to X is \mathcal{T}_X , and whose localization at any point $x \in X$ is given by

$$\mathcal{T}_{Y,x} = \sum_{i=1}^l \mathcal{O}_{Y,x} \cdot z_i \frac{\partial}{\partial z_i} + \sum_{j=1+1}^n \mathcal{O}_{Y,x} \cdot \frac{\partial}{\partial z_j},$$

where the local coordinates z_1, z_2, \dots, z_n around x are chosen in such a way that $X = \{z_1 z_2 \cdots z_l = 0\}$. In fact, $\mathcal{T}_Y\langle X \rangle = (\Omega_Y^1(\log X))^\vee$, where $\Omega_Y^1(\log X)$ denotes the sheaf of meromorphic 1-forms on Y that have at most logarithmic poles along X .

Also, when X is an integral curve sitting on a smooth surface Y , by (2.1) the sheaf $\mathcal{T}_Y\langle X \rangle$ is an elementary transformation of the locally free sheaf \mathcal{T}_Y , and then it is locally free (see, e.g., Gómez, 2001, Lemma 2.2).

Recall the following basic result.

Proposition 2.4 (see Sernesi, 2006, Proposition 3.4.17). *The locally trivial deformations of the pair (Y, X) (equivalently of the closed embedding j) are controlled by the sheaf $\mathcal{T}_Y\langle X \rangle$; namely:*

- (i) *The obstructions lie in $H^2(Y, \mathcal{T}_Y\langle X \rangle)$;*
- (ii) *First-order, locally trivial deformations are parametrized by $H^1(Y, \mathcal{T}_Y\langle X \rangle)$;*
- (iii) *Infinitesimal automorphisms are parametrized by $H^0(Y, \mathcal{T}_Y\langle X \rangle)$.*

The map that associates to a first-order, locally trivial deformation of (Y, X) the corresponding first-order deformation of X is the map

$$H^1(r) : H^1(Y, \mathcal{T}_Y\langle X \rangle) \longrightarrow H^1(X, \mathcal{T}_X), \quad (2.5)$$

induced in cohomology by (2.2).

In the rest of the article, we will focus on the case of nodal curves on a surface.

3. MUKAI'S RESULTS ON SMOOTH, CANONICAL CURVES ON GENERAL, MARKED K3 SURFACES, AND BEAUVILLE'S INFINITESIMAL APPROACH

In this section, we shall briefly recall some results of Mukai (1988, 1992) and the infinitesimal approach considered by Beauville (2004, §5).

Let $p \geq 2$ be an integer. Let \mathcal{B}_p be the moduli stack of smooth K3 surfaces marked by a globally generated, primitive line bundle of sectional genus p . That is, the elements of \mathcal{B}_p are pairs (S, H) , where S is a smooth K3 surface, and H is a globally generated line bundle on S with $H^2 = 2p - 2$ and such that H is nondivisible in $\text{Pic}(S)$. It is well known that \mathcal{B}_p is smooth, irreducible, and of dimension 19 (cf., e.g., Barth et al., 2004, Thm. VIII 7.3 and p. 366 for the scheme structure; the same conclusions hold also for the stack structure of \mathcal{B}_p).

Definition 3.1. Let $\mathcal{K}C_p$ be the algebraic stack of pairs (S, C) , where $(S, H) \in \mathcal{B}_p$, $p \geq 2$, and $C \in |H|$ is a smooth irreducible curve.

Observe that there is an induced, surjective morphism of stacks

$$\pi : \mathcal{K}C_p \longrightarrow \mathcal{B}_p \quad (3.2)$$

given by the natural projection. From Beauville (2004, §(5.2)), for any $(S, C) \in \mathcal{K}C_p$, by the Serre duality one has

$$H^2(S, \mathcal{T}_S\langle C \rangle) = H^0(S, \Omega_S^1(\log C))^\vee = (0). \quad (3.3)$$

Furthermore, since C is a smooth curve of genus $p \geq 2$ and since $\mathcal{T}_S \cong \Omega_S^1$, S being a K3 surface and \mathcal{T}_S a rank-two vector bundle on it, by (2.3), we have

$$H^0(S, \mathcal{T}_S\langle C \rangle) = (0).$$

In particular, from Proposition 2.4, $\mathcal{H}C_p$ is a smooth stack of dimension

$$\dim(\mathcal{H}C_p) = h^1(S, \mathcal{T}_S\langle C \rangle) = 19 + p.$$

Since the fibers of π are connected, $\mathcal{H}C_p$ is also irreducible.

Let \mathcal{M}_p be the moduli stack of smooth curves of genus p , which is irreducible and of dimension $3p - 3$, since $p \geq 2$ by assumption. One has a natural morphism of stacks

$$c_p : \mathcal{H}C_p \longrightarrow \mathcal{M}_p \tag{3.4}$$

defined as

$$c_p((S, C)) = [C] \in \mathcal{M}_p,$$

where $[C]$ denotes the isomorphism class of $C \subset S$.

Observe that

$$\begin{aligned} \dim(\mathcal{H}C_p) &> \dim(\mathcal{M}_p), & \text{for } p \leq 10, \\ \dim(\mathcal{H}C_p) &= \dim(\mathcal{M}_p), & \text{for } p = 11, \\ \dim(\mathcal{H}C_p) &< \dim(\mathcal{M}_p), & \text{for } p \geq 12. \end{aligned} \tag{3.5}$$

If we denote by F_p the general fibre of c_p , then the *expected dimension* of F_p is

$$\exp \dim(F_p) = \begin{cases} 22 - 2p & \text{for } p \leq 10, \\ 0 & \text{for } p \geq 11. \end{cases} \tag{3.6}$$

The main results concerning the morphism c_p are contained in the following theorem.

Theorem 3.7 (Mukai). *With notation as above:*

- (i) c_p is dominant for $p \leq 9$ and $p = 11$ (cf. Mukai, 1988);
- (ii) c_p is not dominant for $p = 10$ (cf. Mukai, 1988). More precisely, its image is a hypersurface in \mathcal{M}_{10} (cf. Cukierman and Ulmer, 1993);
- (iii) c_p is generically finite onto its image, for $p = 11$ and for $p \geq 13$, but not for $p = 12$ (cf. Mukai, 1992).

Remark 3.8. (1) In particular, from (3.6) and from Theorem 3.7, one has

$$\dim(F_p) = \exp \dim(F_p),$$

unless either

- (i) $p = 10$, in which case $\dim(F_p) = \exp \dim(F_p) + 1 = 3$; or
- (ii) $p = 12$, in which case $\dim(F_p) \geq 1$.

(2) When the map c_p is not dominant, one can look at it as a way to produce hopefully interesting cycles in the moduli space of curves. The case $p = 10$ is particularly relevant, as the divisor in \mathcal{M}_{10} parametrizing curves lying on a $K3$ surface was the first counterexample to the slope conjecture (see Farkas and Popa, 2005).

Beauville (2004, §(5.2)) considered the morphism c_p from a differential point of view. Let $(S, C) \in \mathcal{H}C_p$ be any point. From Proposition 2.4, the differential of c_p at the point (S, C) can be identified with the map

$$H^1(r) : H^1(\mathcal{T}_S\langle C \rangle) \longrightarrow H^1(\mathcal{T}_C),$$

as in (2.5). From (2.3) and (3.3), it follows that $H^1(r)$ fits in the exact sequence

$$0 \longrightarrow H^1(S, \mathcal{T}_S(-C)) \longrightarrow H^1(S, \mathcal{T}_S\langle C \rangle) \xrightarrow{H^1(r)} H^1(C, \mathcal{T}_C) \longrightarrow H^2(S, \mathcal{T}_S(-C)) \longrightarrow 0.$$

Using Serre duality and the fact that ω_S is trivial, we get

$$H^j(S, \mathcal{T}_S(-C)) \cong H^{2-j}(S, \Omega_S^1(C))^\vee, \quad 0 \leq j \leq 2. \tag{3.9}$$

From (3.9), we obtain that the morphism c_p is:

- (i) *Smooth* at $(S, C) \in \mathcal{H}C_p$ (i.e., the differential $(c_p)_*$ at the point (S, C) is surjective) if and only if $H^0(S, \Omega_S^1(C)) = (0)$;
- (ii) *Unramified* at $(S, C) \in \mathcal{H}C_p$ (i.e., the differential $(c_p)_*$ at the point (S, C) is injective) if and only if $H^1(S, \Omega_S^1(C)) = (0)$.

Note that the above conditions depend only on the marking $H = \mathcal{O}_S(C)$ and not on the particular curve C in $|H|$.

With this approach, Theorem 3.7 is equivalent to the following proposition.

Proposition 3.10 (see Beauville, 2004, §(5.2)). *Let (S, H) be a general primitively polarized $K3$ surface of genus $p \geq 2$. We have:*

- (i) $h^0(S, \Omega_S^1(H)) = 0$, for $p \leq 9$ and $p = 11$;
- (ii) $h^0(S, \Omega_S^1(H)) = 1$, for $p = 10$;
- (iii) $h^1(S, \Omega_S^1(H)) = 0$, for $p = 11$ and $p \geq 13$;
- (iv) $h^1(S, \Omega_S^1(H)) \geq 1$, for $p = 12$.

Remark 3.11. Since $h^2(\Omega_S^1(H)) = h^0(\mathcal{T}_S(-H)) = 0$, $c_1(\Omega_S^1(H)) = 2H$, and $c_2(\Omega_S^1(H)) = H^2 + 24 = 2p + 22$, we have, by Riemann–Roch

$$h^0(\Omega_S^1(H)) = \chi(\Omega_S^1(H)) + h^1(\Omega_S^1(H))$$

$$\begin{aligned}
 &= \frac{c_1(\Omega_S^1(H))^2}{2} - c_2(\Omega_S^1(H)) + 2\text{rk}(\Omega_S^1(H)) + h^1(\Omega_S^1(H)) \\
 &= 2p - 22 + h^1(\Omega_S^1(H)) \geq 2p - 22.
 \end{aligned}$$

In particular, $h^0(\Omega_S^1(H)) \geq 3$ if $p \geq 12$ (cf. also Question 5.7).

4. THE APPROACH TO THE NODAL CASE

By using Proposition 2.4 and a similar approach as in Section 3, we want to deduce some extensions of Theorem 3.7 to irreducible, nodal curves in the primitive linear system $|H|$ on a general primitively polarized K3 surface of genus $p \geq 3$. In particular, we are interested in determining when the normalization of such a singular curve is an (abstract) smooth curve with general moduli.

To do this, we have to fix some notation and to prove some results that will be used in what follows. First we recall that, for any smooth surface S and any line bundle H on S , such that $|H|$ contains smooth, irreducible curves of genus $p := p_a(H)$, and any positive integer $\delta \leq p$, one denotes by

$$V_{|H|,\delta}(S) \quad \text{or simply } V_{|H|,\delta}$$

the locally closed and functorially defined subscheme of $|H|$ parametrizing the universal family of irreducible curves in $|H|$ having δ nodes as the only singularities and, consequently, geometric genus $g := p - \delta$. These are classically called *Severi varieties* of irreducible, δ -nodal curves on S in $|H|$.

It is well known, as a direct consequence of Mumford’s theorem on the existence of nodal rational curves on K3 surfaces (see, e.g., Barth et al., 2004, pp. 365–367) and standard results on Severi varieties (see, e.g., Chiantini and Sernesi, 1997; Flamini, 2001; Tannenbaum, 1982), that if $(S, H) \in \mathcal{B}_p$ is general, $p \geq 2$, then

$$V_{|H|,\delta} \text{ is nonempty and regular,} \tag{4.1}$$

i.e., it is smooth and (each of its irreducible components is) of the expected dimension $g = p - \delta$, for each $\delta \leq p$. (In fact, the regularity holds whenever $V_{|H|,\delta}$ is nonempty.)

From now on, we shall always consider

$$p \geq 3 \quad \text{and} \quad 0 \leq \delta \leq p - 2, \quad \text{so that } g \geq 2. \tag{4.2}$$

Similarly as in Definition 3.1, we have the following definition.

Definition 4.3. For any p and δ as in (4.2), let $\mathcal{V}_{p,\delta}$ be the stack of pairs (S, X) , such that $(S, H) \in \mathcal{B}_p$ and $[X] \in V_{|H|,\delta}(S)$.

Of course, $\mathcal{V}_{p,0} = \mathcal{H}C_p$ as in Definition 3.1.

For any fixed p and any δ as in (4.2), the stacks $\mathcal{V}_{p,\delta}$ are locally closed substacks of a natural enlargement $\overline{\mathcal{H}C}_p$ of $\mathcal{H}C_p$, which is defined as the stack

Downloaded By: [Flamini, Flamini] At: 13:11 3 December 2008

of pairs (S, C) , where $(S, H) \in \mathcal{B}_p$ and $C \in |H|$. It follows that the stacks $\mathcal{V}_{p,\delta}$ are algebraic because $\overline{\mathcal{H}C}_p$ is.

Consider $\mathcal{B}_p^0 \subset \mathcal{B}_p$ the open dense substack parametrizing elements (S, H) in \mathcal{B}_p that verify (4.1).

For any δ as above, let

$$\pi_\delta : \mathcal{V}_{p,\delta} \longrightarrow \mathcal{B}_p^0 \tag{4.4}$$

be the surjective morphism given by the projection.

Let $(S, X) \in \mathcal{V}_{p,\delta}$. From (2.1), we can consider the exact sequence

$$0 \longrightarrow \mathcal{F}_S \langle X \rangle \longrightarrow \mathcal{F}_S \longrightarrow \mathcal{N}'_{X/S} \longrightarrow 0. \tag{4.5}$$

Since S is a $K3$ surface

$$H^2(\mathcal{F}_S) \cong H^0(\Omega_S^1) \cong H^0(\mathcal{F}_S) = (0).$$

Therefore, passing to cohomology in (4.5), we get

$$0 \longrightarrow H^0(\mathcal{N}'_{X/S}) \longrightarrow H^1(\mathcal{F}_S \langle X \rangle) \xrightarrow{H^1(r)} H^1(\mathcal{F}_S) \xrightarrow{\alpha} H^1(\mathcal{N}'_{X/S}) \longrightarrow H^2(\mathcal{F}_S \langle X \rangle) \longrightarrow 0, \tag{4.6}$$

where $H^1(r)$ is as in (2.5).

From Tannenbaum (1982, Lemma 2.4 and Theorem 2.6), we have

$$H^1(\mathcal{N}'_{X/S}) \cong \mathbb{C}, \quad H^0(\mathcal{N}'_{X/S}) \cong T_{[X]}(V_{|H|,\delta}(S)). \tag{4.7}$$

Moreover, recall that $V_{|H|,\delta}(S)$ is regular at $[X]$ (i.e., it is smooth at $[X]$ and of the expected dimension g). This means that, despite the fact that $h^1(\mathcal{N}'_{X/S}) = 1$, the infinitesimal, locally trivial deformations of the closed embedding $X \xrightarrow{j} S$, with S fixed, are unobstructed, and the nodes impose independent conditions to locally trivial deformations (cf. Tannenbaum, 1982, Remark 2.7).

We want to show that, for $\delta > 0$, the properties of the stack $\mathcal{V}_{p,\delta}$ are similar to those of the stack $\mathcal{H}C_p$.

Proposition 4.8. *Let p, δ and g be positive integers as in (4.2).*

Then, for any $(S, X) \in \mathcal{V}_{p,\delta}$, we have

$$h^0(\mathcal{F}_S \langle X \rangle) = 0, \quad h^2(\mathcal{F}_S \langle X \rangle) = 0. \tag{4.9}$$

In particular:

- (i) *The stack $\mathcal{V}_{p,\delta}$ is smooth;*
- (ii) *Any irreducible component $\mathcal{V} \subseteq \mathcal{V}_{p,\delta}$ has dimension $h^1(\mathcal{F}_S \langle X \rangle) = 19 + g$;*
- (iii) *The morphism π_δ is smooth and any irreducible component $\mathcal{V} \subseteq \mathcal{V}_{p,\delta}$ smoothly dominates \mathcal{B}_p^0 .*

Proof. The first equality in (4.9) directly follows from (4.5).

For what concerns the second equality in (4.9), we consider (4.6) above: for any point $(S, X) \in \mathcal{V}_{p,\delta}$, we have that

$$0 \longrightarrow H^0(\mathcal{N}'_{X/S}) \longrightarrow H^1(\mathcal{T}_S\langle X \rangle) \longrightarrow \text{Ker}(\alpha) \longrightarrow 0 \tag{4.10}$$

can be read as the natural differential sequence

$$0 \longrightarrow T_{[X]}(V_{|H|,\delta}(S)) \longrightarrow T_{(S,X)}(\mathcal{V}_{p,\delta}) \longrightarrow T_{[S]}(\mathcal{B}_p^0) \longrightarrow 0; \tag{4.11}$$

indeed, \mathcal{B}_p^0 is smooth of dimension 19, whereas $h^1(\mathcal{T}_S) = 20$ and $h^1(\mathcal{N}'_{X/S}) = 1$ by (4.7), thus we have $\text{Im}(\alpha) = H^1(\mathcal{N}'_{X/S}) \cong H^1(\mathcal{N}_{X/S})$, i.e., the elements of $\text{Ker}(\alpha)$ can be identified with the first-order deformations of S preserving the genus p marking.

Moreover, it follows that $H^2(\mathcal{T}_S\langle X \rangle) = (0)$, i.e., the infinitesimal deformations of the closed embedding $X \xrightarrow{j} S$, with S not fixed, are unobstructed.

Now, (i) directly follows from Proposition 2.4, whereas (ii) and (iii) follow from Proposition 2.4, (4.11) and from what recalled above on Severi varieties on general $K3$ surfaces. \square

In particular, we have the following corollary.

Corollary 4.12. *There exists an open, dense substack $\mathcal{U}_p \subseteq \mathcal{B}_p^0$ such that the number of irreducible components of $V_{|H|,\delta}(S)$ is constant, for (S, H) varying in \mathcal{U}_p .*

Proof. This directly follows from Proposition 4.8(iii) and from the fact that, \mathcal{V} being irreducible, the general fibre of $\pi_\delta|_{\mathcal{V}}$ has to be irreducible. \square

Remark 4.13. Notice that it is not known whether, for a general primitively polarized $K3$ surface (S, H) , the Severi varieties $V_{|kH|,\delta}(S)$, for any integer $k \geq 1$, are irreducible as soon as they have positive dimension. Interesting work on this topic has been done recently in Dedieu (Preprint).

We need a reformulation of Proposition 2.4 in this specific situation, which will be particularly useful for our aims.

First we have to recall some general facts. Let $(S, X) \in \mathcal{V}_{p,\delta}$ be any pair as above. Let

$$\varphi : C \longrightarrow X \subset S$$

be the normalization morphism. By the fact that X is a curve on S , one can consider the exact sequence

$$0 \longrightarrow \mathcal{T}_C \longrightarrow \varphi^*(\mathcal{T}_S) \longrightarrow \mathcal{N}_\varphi \longrightarrow 0, \tag{4.14}$$

defining \mathcal{N}_φ as the *normal sheaf* to φ . Since X is a nodal curve, it is well known that \mathcal{N}_φ is a line bundle on C (cf., e.g., Sernesi, 2006).

Remark 4.15. Denote as above by $\varphi : C \rightarrow X \subset S$ the normalization morphism of a curve sitting on a surface S . Suppose moreover that S is a $K3$ surface. Then

$\mathcal{N}_\varphi = \omega_C$. Therefore, unless C is hyperelliptic, we always have the surjectivity of the map

$$H^0(C, \mathcal{N}_\varphi) \otimes H^0(C, \omega_C^{\otimes 2}) \longrightarrow H^0(C, \mathcal{N}_\varphi \otimes \omega_C^{\otimes 2}).$$

Arguing as in Beauville and M erindol (1987, Proposition 1), we get that the splitting of the exact sequence (4.14) is *equivalent* to the triviality, as abstract deformations of C , of the infinitesimal deformations parametrized by $H^0(C, \mathcal{N}_\varphi)$, i.e., the coboundary map $H^0(C, \mathcal{N}_\varphi) \longrightarrow H^1(C, \mathcal{T}_C)$ is zero.

Lemma 4.16. *With notation as above, one has*

$$\varphi_*(\mathcal{N}_\varphi) \cong \mathcal{N}'_{X/S}.$$

In particular,

$$H^i(\mathcal{N}'_{X/S}) \cong H^i(\mathcal{N}_\varphi), \quad 0 \leq i \leq 1 \tag{4.17}$$

Proof. The reader is referred to Tannenbaum (1980, p. 111) where, with the notation therein, since X is nodal, the *Jacobian ideal* \mathfrak{J} coincides with the *conductor ideal* \mathfrak{C} , i.e., $\mathcal{N}'_{X/S} = \mathfrak{J}\mathcal{N}_{X/S} = \mathfrak{C}\mathcal{N}_{X/S} \cong \varphi_*(\mathcal{N}_\varphi)$. □

On the other hand, if $N = \text{Sing}(X)$, let

$$\mu_N : \tilde{S} \longrightarrow S$$

be the blowup of S along N . Thus, μ_N induces the embedded resolution of X in S , i.e., we have the commutative diagram

$$\begin{array}{ccc} C & \subset & \tilde{S} \\ \downarrow \varphi & & \downarrow \mu_N \\ X & \subset & S, \end{array}$$

where $\mu_N|_C = \varphi$.

Since

$$\varphi^*(\mathcal{T}_S) \cong \mu_N^*(\mathcal{T}_S) \otimes \mathcal{O}_C$$

and since $\mu_N|_C = \varphi$, we also have the exact sequence on \tilde{S}

$$0 \longrightarrow \mu_N^*(\mathcal{T}_S)(-C) \longrightarrow \mu_N^*(\mathcal{T}_S) \longrightarrow \varphi^*(\mathcal{T}_S) \longrightarrow 0. \tag{4.18}$$

We denote by

$$\mu_N^*(\mathcal{T}_S) \xrightarrow{\lambda} \mathcal{N}_\varphi \longrightarrow 0 \tag{4.19}$$

the composition of the two surjections in (4.18) and (4.14).

Definition 4.20. With notation as above, let

$$\mu_N^*(\mathcal{T}_S)\langle C \rangle := \text{Ker}(\lambda).$$

From (4.14), (4.18), and (4.19), the sheaf $\mu_N^*(\mathcal{T}_S)\langle C \rangle$ sits in the following natural exact diagram:

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & \mu_N^*(\mathcal{T}_S)(-C) & \xrightarrow{=} & \mu_N^*(\mathcal{T}_S)(-C) & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & \mu_N^*(\mathcal{T}_S)\langle C \rangle & \longrightarrow & \mu_N^*(\mathcal{T}_S) & \xrightarrow{\lambda} & \mathcal{N}_\varphi \longrightarrow 0 \\
 & & \downarrow \tau & & \downarrow & & \parallel \\
 0 & \longrightarrow & \mathcal{T}_C & \longrightarrow & \varphi^*(\mathcal{T}_S) & \longrightarrow & \mathcal{N}_\varphi \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0.
 \end{array} \tag{4.21}$$

We have the following proposition.

Proposition 4.22. *Let $(S, X) \in \mathcal{V}_{p,\delta}$. Then*

$$H^i(S, \mathcal{T}_S\langle X \rangle) \cong H^i(\tilde{S}, \mu_N^*(\mathcal{T}_S)\langle C \rangle), \quad 0 \leq i \leq 2. \tag{4.23}$$

In particular, the locally trivial deformations of the pair (S, X) are also governed by the sheaf $\mu_N^(\mathcal{T}_S)\langle C \rangle$, i.e.,:*

- (i) *The obstructions lie in $H^2(\tilde{S}, \mu_N^*(\mathcal{T}_S)\langle C \rangle)$;*
- (ii) *First-order, locally trivial deformations are parametrized by $H^1(\tilde{S}, \mu_N^*(\mathcal{T}_S)\langle C \rangle)$;*
- (iii) *Infinitesimal automorphisms are parametrized by $H^0(\tilde{S}, \mu_N^*(\mathcal{T}_S)\langle C \rangle)$.*

Proof. Observe that, in this situation, we have the exact sequence (4.5), i.e.,

$$0 \longrightarrow \mathcal{T}_S\langle X \rangle \longrightarrow \mathcal{T}_S \longrightarrow \mathcal{N}'_{X/S} \longrightarrow 0. \tag{4.24}$$

On the other hand, as above, let $\mu_N : \tilde{S} \rightarrow S$ be the blowup of S along N , and let $C \subset \tilde{S}$ be the proper transform of $X \subset S$. From the second row of (4.21), we get

$$0 \longrightarrow \mu_N^*(\mathcal{T}_S)\langle C \rangle \longrightarrow \mu_N^*(\mathcal{T}_S) \longrightarrow \mathcal{N}_\varphi \longrightarrow 0. \tag{4.25}$$

Since S is smooth then, by the projection formula, we have

$$\mu_{N*}(\mu_N^*(\mathcal{T}_S)) \cong \mathcal{T}_S \otimes \mu_{N*}(\mathcal{O}_{\tilde{S}}) \cong \mathcal{T}_S.$$

Now we apply μ_{N^*} to (4.25), getting

$$0 \longrightarrow \mu_{N^*}(\mu_N^*(\mathcal{T}_S)\langle C \rangle) \longrightarrow \mu_{N^*}(\mu_N^*(\mathcal{T}_S)) \longrightarrow \varphi_*(\mathcal{N}_\varphi), \tag{4.26}$$

where the equality $\mu_{N^*}(\mathcal{N}_\varphi) = \varphi_*(\mathcal{N}_\varphi)$ directly follows from the facts that \mathcal{N}_φ is a line-bundle on C and that, as above, $\mu_N|_C = \varphi$.

By using the exact sequences (4.24) and (4.26), together with Lemma 4.16, we get

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{T}_S\langle X \rangle & \longrightarrow & \mathcal{T}_S & \longrightarrow & \mathcal{N}'_{X/S} \longrightarrow 0 \\ & & \downarrow & & \downarrow \cong & & \downarrow \cong \\ 0 & \longrightarrow & \mu_{N^*}(\mu_N^*(\mathcal{T}_S)\langle C \rangle) & \longrightarrow & \mu_{N^*}(\mu_N^*(\mathcal{T}_S)) & \longrightarrow & \varphi_*(\mathcal{N}_\varphi) \longrightarrow 0. \end{array} \tag{4.27}$$

This implies $\mathcal{T}_S\langle X \rangle \cong \mu_{N^*}(\mu_N^*(\mathcal{T}_S)\langle C \rangle)$. Since μ_N is birational, by Leray’s isomorphism, we obtain (4.23).

The last part of the statement directly follows from Proposition 2.4. □

Let now $V_{p,\delta} \rightarrow \mathcal{V}_{p,\delta}$ be an étale atlas, and let

$$\begin{array}{ccc} \mathcal{X}^C & \longrightarrow & \mathcal{S} \\ & \searrow \rho & \downarrow \\ & & V_{p,\delta} \end{array} \tag{4.28}$$

be the family induced by the universal one. Since $V_{p,\delta}$ is a smooth, in particular normal, scheme we are in position to apply the results in Tossier (1980, p.80), i.e., there exists a commutative diagram

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{\Phi} & \mathcal{X} \\ & \searrow \tilde{\rho} & \downarrow \rho \\ & & V_{p,\delta} \end{array} \tag{4.29}$$

where

- (i) Φ is the normalization morphism,
- (ii) $\tilde{\rho}$ is smooth,
- (iii) $\tilde{\rho}$ gives the simultaneous desingularization of the fibres of the family ρ ; namely, for each $v \in V_{p,\delta}$, $\mathcal{C}(v)$ is a smooth curve of genus $g = p - \delta$, which is the normalization of the irreducible, nodal curve $\mathcal{X}(v)$.

Thus, as in (3.4), the family $\tilde{\rho}$ defines the natural morphism

$$V_{p,\delta} \xrightarrow{c_{p,\delta}} \mathcal{M}_g, \tag{4.30}$$

by sending v to the birational isomorphism class $[\mathcal{C}(v)] \in \mathcal{M}_g$.

As discussed in Section 3 for smooth curves, by Propositions 2.4 and 4.22, and by passing to cohomology in the left-hand-side column of diagram (4.21), the differential of the morphism $c_{p,\delta}$ at a point $v \in V_{p,\delta}$ parametrizing a pair (S, X) can be identified with the cohomology map

$$H^1(\mu_N^*(\mathcal{T}_S)\langle C \rangle) \xrightarrow{H^1(\tau)} H^1(\mathcal{T}_C). \tag{4.31}$$

As in (3.5), we have

$$\begin{aligned} \dim(V_{p,\delta}) &> \dim(\mathcal{M}_g), & \text{for } g \leq 10, \\ \dim(V_{p,\delta}) &= \dim(\mathcal{M}_g), & \text{for } g = 11, \\ \dim(V_{p,\delta}) &< \dim(\mathcal{M}_g), & \text{for } g \geq 12. \end{aligned} \tag{4.32}$$

In particular, independently from $p \geq 3$, for any $g \leq 11$ the morphism $c_{p,\delta}$ is expected to be dominant. Moreover, as in the smooth case of Section 3, if we denote by $F_{p,\delta}$ the general fibre of $c_{p,\delta}$, then, for any $p \geq 3$, the *expected dimension* of $F_{p,\delta}$ is

$$\exp \dim(F_{p,\delta}) = \begin{cases} 22 - 2g & \text{for } g \leq 10, p \geq g + 1, \\ 0 & \text{for } g \geq 11, p \geq g + 1. \end{cases} \tag{4.33}$$

5. GENERAL MODULI FOR $g < p \leq 11$

The aim of this section is to give some partial affirmative answers to the above expectations. Namely, we show that on a general, primitively polarized K3 surface of genus $3 \leq p \leq 11$, the normalizations of δ -nodal curves in $|H|$, with $\delta > 0$ as in (4.2), define families of smooth curves with general moduli.

Precisely, by recalling Proposition 4.8, we have the following proposition.

Theorem 5.1. *Let $3 \leq p \leq 11$ be an integer. Let δ and $g = p - \delta$ be positive integers as in (4.2).*

Let $V \subseteq V_{p,\delta}$ be any irreducible component, and let

$$c_{p,\delta}|_V : V \longrightarrow \mathcal{M}_g$$

be the restriction to V of the morphism $c_{p,\delta}$ as in (4.30).

Then, for any $2 \leq g < p \leq 11$, $c_{p,\delta}|_V$ is dominant. In particular, for the general fibre $F_{p,\delta}$ of $c_{p,\delta}|_V$, we have

$$\dim(F_{p,\delta}) = 22 - 2g = \exp \dim(F_{p,\delta}).$$

Remark 5.2. (1) Recall that, if (S, H) is general in \mathcal{B}_{10} , smooth curves in $|H|$ are not with general moduli (cf. Theorem 3.7(ii)). On the contrary, from Theorem 5.1, if (S, H) is general in \mathcal{B}_{11} , then nodal curves in $V_{|H|,1}$ have normalizations of geometric genus $g = 10$ that are curves with general moduli.

Downloaded By: [Flaminio, Flaminio] At: 13:11 3 December 2008

(2) At the same time, if (S, H) is general in \mathcal{B}_{10} , from Theorem 5.1, it follows that the general, irreducible, δ -nodal curve in the linear system $|H|$ has a normalization of genus $g = 10 - \delta$ with general moduli.

In particular, we have the following situation: consider the rational map

$$\overline{\mathcal{H}C}_{10} \dashrightarrow \overline{\mathcal{M}}_{10},$$

which is defined on the open substack $\overline{\mathcal{H}C}_{10}^0$ of pairs (S, X) s.t. $S = (S, H)$ is in \mathcal{B}_{10}^0 , and X is nodal and irreducible. Let

$$\overline{c}_{10} : \overline{\mathcal{H}C}_{10}^0 \longrightarrow \overline{\mathcal{M}}_{10}$$

be the induced morphism. Then $\text{Im}(\overline{c}_{10})$ is a divisor in $\overline{\mathcal{M}}_{10}$ such that

$$\text{Im}(\overline{c}_{10}) \cap \partial \overline{\mathcal{M}}_{10} \subset \Delta_0$$

is a divisor whose general element has normalization a general curve of genus 9.

Since this divisor has dimension

$$\dim(\Delta_0) - 1 = \dim(\overline{\mathcal{M}}_{10}) - 2 = 25,$$

the universal curve

$$\begin{array}{c} \mathcal{C} \\ \downarrow \\ \text{Im}(\overline{c}_{10}) \cap \Delta_0 \end{array}$$

induces a rational map

$$(\text{Im}(\overline{c}_{10}) \cap \Delta_0) \dashrightarrow \mathcal{M}_9$$

that is dominant by Theorem 5.1, and whose general fibre has dimension 1. Precisely, this fibre determines a 1-dimensional subscheme in the second symmetric product of the curve of genus 9 parametrized by the image point in \mathcal{M}_9 .

Proof of Theorem 5.1. From (4.31) and the left vertical column in (4.21), a sufficient condition for the surjectivity of the differential of $c_{p,\delta}|_V$ at a point $v \in V$ parametrizing a pair (S, X) is that

$$H^2(\mu_N^*(\mathcal{T}_S)(-C)) = (0). \tag{5.3}$$

If $E = \sum_{i=1}^{\delta} E_i$ denotes the μ_N -exceptional divisor on \tilde{S} , by Serre duality,

$$H^2(\mu_N^*(\mathcal{T}_S)(-C)) \cong H^0(\mu_N^*(\Omega_S^1)(C + E)) \cong H^0(\mu_N^*(\Omega_S^1(H))(-E)),$$

since $\mu_N^*(X) = C + 2E$ and since $X \sim H$ on S .

By the Leray isomorphism,

$$H^0(\mu_N^*(\Omega_S^1(H))(-E)) \cong H^0(\mathcal{F}_{N/S} \otimes \Omega_S^1(H)),$$

where $\mathcal{F}_{N/S}$ denotes the ideal sheaf of N in S . One has the exact sequence

$$0 \longrightarrow \mathcal{F}_{N/S} \otimes \Omega_S^1(H) \longrightarrow \Omega_S^1(H) \longrightarrow \Omega_S^1(H)|_N \longrightarrow 0. \tag{5.4}$$

For what concerns the cases either $3 \leq p \leq 9$ or $p = 11$, from (5.4) and from Proposition 3.10(i), we get that also $H^0(\mathcal{F}_{N/S} \otimes \Omega_S^1(H)) = (0)$, which concludes the proof in this case.

For what concerns the case $p = 10$, from Proposition 3.10(ii) we know that, if (S, H) is general in \mathcal{B}_{10} , then $H^0(\Omega_S^1(H)) \cong \mathbb{C}$. For any δ as above, let $W \subseteq V_{|H|, \delta}$ be any irreducible component of the Severi variety of irreducible, δ -nodal curves in $|H|$ on S . Let $[X] \in W$ be a general point and let $N = \text{Sing}(X)$.

From the exact sequence (5.4), we know that

$$H^0(\mathcal{F}_{N/S} \otimes \Omega_S^1(H)) \hookrightarrow H^0(\Omega_S^1(H)) \cong \mathbb{C}.$$

We claim that this implies $h^0(\mathcal{F}_{N/S} \otimes \Omega_S^1(H)) = 0$, so the statement follows.

Indeed, if $h^0(\mathcal{F}_{N/S} \otimes \Omega_S^1(H)) \neq 0$, then we would have

$$H^0(\mathcal{F}_{N/S} \otimes \Omega_S^1(H)) \cong H^0(\Omega_S^1(H)),$$

which means that the unique (up to scalar multiplication) nonzero global section of $H^0(\Omega_S^1(H))$ would pass through $N = \text{Sing}(X)$. Since X is general in a $g \geq 2$ dimensional family of curves, this is a contradiction. \square

Our result naturally leads to some questions.

Question 5.5. When X is smooth, the surjectivity of the Wahl map

$$\phi_{\omega_X} : \wedge^2 H^0(X, \omega_X) \longrightarrow H^0(X, \omega_X^{\otimes 3})$$

is an obstruction to embed X in a $K3$ surface (see Wahl, 1987 and also Beauville and M erindol, 1987). It is natural to ask whether a similar result holds in the singular case. Precisely it would be interesting to understand whether there exists a Wahl-type obstruction for a smooth curve to have a nodal model lying on a $K3$ surface.

Question 5.6. Does the image of the map $c_{p, \delta}$ meet the hyperelliptic locus \mathcal{H}_g in \mathcal{M}_g ? If yes, and (S_0, X_0) is a pair mapped into the hyperelliptic locus, does the dimension of the locus of curves in $V_{|X_0|, \delta}(S_0)$ having hyperelliptic normalizations coincide with the expected one, which is two, cf. Flamini et al. (2007b, Lemma 5.1)? In fact, in Flamini et al. (2007b, Lemma 5.1) we show that the dimension is two if $\text{Pic}(S) \simeq \mathbb{Z}[X_0]$. Answers to these questions, as explained in Flamini et al. (2007b, §6), would yield a better understanding of rational curves in the Hilbert square $S^{[2]}$ of the general $K3$ surface S , and then of its Mori cone $\text{NE}(S^{[2]})$. The answers seem to be subtle, as they do not depend only on the geometric genus of X_0 but also on

the number of nodes. For instance, for a general $(S, H) \in \mathcal{B}_p$, the locus of curves in $V_{|H|, \delta}(S)$ having hyperelliptic normalizations is *empty* when $\delta \leq (p-3)/2$ (Flamini et al., 2007a, Theorem 1), and *nonempty*, and two-dimensional, for $p-3 \leq \delta \leq p-2$ (for $\delta = p-2$ is obvious; for $\delta = p-3$ cf. Flamini et al., 2007b, Theorem 5.2). Of course, similar questions may be asked for other gonality strata in \mathcal{M}_g .

Question 5.7. What about the cases $p \geq 12$? From (4.32) and Theorem 5.1, one could in principle expect the following situation:

Let $p \geq 12$ be an integer. Let δ and $g = p - \delta$ be positive integers as in (4.2). For any irreducible component $V \subseteq V_{p, \delta}$, consider

$$c_{p, \delta}|_V : V \longrightarrow \mathcal{M}_g.$$

Then:

- (i) For any $2 \leq g \leq 11$, the morphism $c_{p, \delta}|_V$ is dominant;
- (ii) For any $12 \leq g < p$, $c_{p, \delta}|_V$ is generically finite.

Possible approaches to investigate the above expectations are the following:

(a) For (i), as in the proof of Theorem 5.1, a sufficient condition for $c_{p, \delta}|_V$ to be smooth at a point $v \in V$ parametrizing a pair (S, X) is $H^0(\mathcal{F}_{N/S} \otimes \Omega_S^1(H)) = (0)$, where $N = \text{Sing}(X)$. By Remark 3.11, one can say that

$$h^0(\mathcal{F}_{N/S} \otimes \Omega_S^1(H)) \geq h^0(\Omega_S^1(H)) - 2\delta \geq 2(p - \delta) - 22 = 2g - 22$$

(note that $2g - 22 \leq 0$ by assumption). On the other hand,

$$h^0(\mathcal{F}_{N/S} \otimes \Omega_S^1(H)) > 0 \quad \text{if } g \geq 12,$$

as it must be (cf. (4.32));

(b) For (ii) above, a sufficient condition for $c_{p, \delta}|_V$ to be unramified at a point $v \in V$ parametrizing a pair (S, X) is $H^1(\mathcal{F}_{N/S} \otimes \Omega_S^1(H)) = (0)$, where $N = \text{Sing}(X)$. Observe that, for $p \geq 13$, from (5.4) we have the exact sequence

$$\dots \longrightarrow H^0(\Omega_S^1(H)) \xrightarrow{ev_N^2} H^0(\Omega_S^1(H)|_N) \longrightarrow H^1(\mathcal{F}_{N/S} \otimes \Omega_S^1(H)) \longrightarrow 0,$$

since $H^1(\Omega_S^1(H)) = (0)$ from Proposition 3.10(iii). Thus, the surjectivity of the evaluation morphism ev_N^2 as above would imply that the differential of $c_{p, \delta}|_V$ at v is unramified.

ACKNOWLEDGMENTS

We warmly thank B. Fantechi and M. Roth for useful conversations, and C. Voisin for her comments.

F. Flamini and E. Sernesi are members of MIUR-GNSAGA at INdAM ‘‘F. Severi’’. The research of A. L. Knutsen was supported by Marie Curie Intra-European Fellowship within the 6th Framework Programme. During the

preparation of the article, G. Pacienza benefited from an “accueil en délégation au CNRS.”

REFERENCES

- Barth, W., Hulek, K., Peters, C., Van de Ven, A. (2004). *Compact Complex Surfaces*. 2nd ed. Berlin: Springer-Verlag.
- Beauville, A. (2004). *Fano Threefolds and $K3$ Surfaces*. The Fano Conference. Univ. Torino, Turin, pp. 175–184.
- Beauville, A., Mériandol, J.-Y. (1987). Sections hyperplanes des surfaces $K3$. *Duke Math. J.* 55:873–878.
- Chiantini, L., Sernesi, E. (1997). Nodal curves on surfaces of general type. *Math. Ann.* 307:41–56.
- Chiantini, L., Ciliberto, C. (1999). On the Severi varieties of surfaces in \mathbb{P}^3 . *J. Alg. Geom.* 8:67–83.
- Cukierman, F., Ulmer, D. (1993). Curves of genus ten on $K3$ surfaces. *Compos. Math.* 89:81–90.
- Dedieu, T. Severi varieties and self rational maps of $K3$ surfaces. Preprint arXiv:0704.3163.
- Farkas, G., Popa, M. (2005). Effective divisors on M_g , curves on $K3$ surfaces and the slope conjecture. *J. Alg. Geom.* 14:241–267.
- Flamini, F. (2001). Some results of regularity for Severi varieties of projective surfaces. *Comm. Algebra* 29(6):2297–2311.
- Flamini, F., Knutsen, A. L., Pacienza, G. (2007a). Singular curves on $K3$ surfaces and linear series on their normalizations. *Internat. J. Math.* 18:1–23.
- Flamini, F., Knutsen, A. L., Pacienza, G. (2007b). On families of rational curves in the Hilbert square of a surface. With an Appendix of E. Sernesi, Partial desingularizations of families of nodal curves. Preprint 2007, arXiv:0704.1367. To appear in *Michigan Math. J.*
- Gómez, T. L. (2001). Brill–Noether theory on singular curves and torsion-free sheaves on surfaces. *Comm. Anal. Geom.* 9:725–756.
- Iitaka, S. (1982). *Algebraic Geometry*. Graduate Texts in Math., 76. New York: Springer-Verlag.
- Mayer, A. (1972). Families of $K3$ surfaces. *Nagoya Math. J.* 48:1–17.
- Mukai, S. (1988). Curves, $K3$ surfaces and Fano 3-folds of genus ≤ 10 . In: *Algebraic Geometry and Commutative Algebra*. Vol. I. Tokyo: Kinokuniya, pp. 357–377.
- Mukai, S. (1992). Fano 3-folds. In: *Complex Projective Geometry* (Trieste–Bergen, 1989). London Math. Soc. Lecture Note Ser. Vol. 179. Cambridge: Cambridge Univ. Press. pp. 255–263.
- Sernesi, E. (2006). *Deformations of Algebraic Schemes*. Grundlehren der Mathematischen Wissenschaften, 334. Berlin: Springer-Verlag.
- Tannenbaum, A. (1980). Families of algebraic curves with nodes. *Compos. Math.* 41:107–126.
- Tannenbaum, A. (1982). Families of curves with nodes on $K - 3$ surfaces. *Math. Ann.* 260:239–253.
- Tessier, B. (1980). *Résolution Simultanée I, II*. Lecture Notes in Math. Vol. 777. Berlin: Springer, pp. 71–146.
- Wahl, J. (1987). The Jacobian algebra of a graded Gorenstein singularity. *Duke Math. J.* 55:843–872.