

CENTRAL LIMITS OF SQUEEZING OPERATORS

L. Accardi

Dipartimento di Matematica
Centro Matematico V. Volterra
Università di Roma II, Roma, Italy

A. Bach

Institut für Theoretische Physik I
Universität Münster, Münster, FRG

Contents

1	Introduction	3
2	The De Moivre-Laplace theorem in the 2-dimensional case	3
3	d-level systems: 1 copy	5
4	d-level systems: N copies	8
5	Inclusion of the number operator	10
6	Squeezing states	14

1 Introduction

In [1] we have proved a quantum De Moivre-Laplace theorem based on a modification of the Giri-von Waldenfels quantum central limit theorem. In [2] P.A. Meyer outlined a method based on direct calculations which, taking advantage of the explicit structure of the algebra of 2×2 matrices, allows a drastic simplification of the proof of the main result of the first part of our paper and relates it with a similar result obtained, independently and simultaneously, by Parthasarathy [5]. In the first part of the present note we simplify the Parthasarathy-Meyer method and extend it to deal with arbitrary d -dimensional Bernoulli processes, where d is a natural integer (cfr. Sections (3),(4). We also prove another statement in Meyer's note (cf. Theorem (5.1)). Finally (Section (6)) we show that the method of proof used in [1] allows, with minor modifications, to solve the problem of the central limit approximation of the squeezing states - a problem left open in [1] and to which, due to the nonlinearity of the coupling, Parthasarathy-Meyer direct computational method cannot be applied.

Acknowledgments.

Part of this paper was written while L.A. was visiting the University of Strasbourg, continued while L.A. was visiting the Center for Mathematical system theory in Gainesville and completed while A.B. was visiting the University of Rome II. The authors are grateful to P.A.Meyer for several useful comments. L.A. also acknowledges support from Grant AFOSR 870249 and ONR N00014-86-K-0538 through the Center for Mathematical System Theory, University of Florida.

2 The De Moivre-Laplace theorem in the 2-dimensional case

Throughout this paper we adopt the notations of [1] with the only exception that, in the first four sections, we use Meyer's normalization for the N -coherent vectors and the N -Weyl operators. Thus, in particular, $\varphi_N(0)$ denotes the vacuum state in $\otimes^n \mathbf{C}^2$; $W_0(z) = \exp(zs^+ - \bar{z}s)$ the Weyl operator on \mathbf{C}^2 ; $W_N(z) = \otimes^N W_0(z/\sqrt{N})$ the N -Weyl operator on $\otimes^N \mathbf{C}^2$; while $\Phi(0)$, $W(z) = \exp(za^+ - \bar{z}a)$ denotes the corresponding objects for the

harmonic oscillator. As usual

$$s_+ = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}; \quad s_- = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}; \quad n_+ = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}; \quad (1)$$

$$\varphi_N(0) = \otimes^N e_1 \quad ; \quad e_1 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

The main result of the first part of [1] is:

Theorem 1 *For every natural integer k and for every $k \in \mathbb{C}, z_1, \dots, z_R \in \mathbb{C}$*

$$\lim_{N \rightarrow \infty} \langle \varphi_N(0), w_N(z_1) \dots w_N(z_R) \varphi_N(0) \rangle = \langle \Phi(0), W(z_1) \dots W(z_R) \Phi(0) \rangle \quad (2)$$

uniformly for z_1, \dots, z_R in a bounded set.

Proof. For $z_j \in \mathbb{C}$ one has

$$\begin{aligned} & \langle \varphi_N(0), W_N(z_1) \dots W_N(z_k) \varphi_N(0) \rangle = \quad (3) \\ & = \langle \otimes^N e_i, \otimes^N W_o \left(\frac{z_1}{\sqrt{N}} \right) \dots \otimes^N W_o \left(\frac{z_k}{\sqrt{N}} \right) \otimes^N e_i \rangle = \langle e_i, W_o \left(\frac{z_i}{\sqrt{N}} \right) e \dots W_o \left(\frac{z_k}{\sqrt{N}} \right) e \rangle^N \\ & = \left\{ \sum_{\substack{(j_1, \dots, j_k) \\ \in \mathbb{N}^k}} \frac{1}{j_1! \dots j_k!} \langle e_i, \prod_{i=1, \dots, k} \left(\frac{z_i}{\sqrt{N}} s^+ - \frac{\bar{z}_i}{\sqrt{N}} s^- \right)^{j_i} e_1 \rangle \right\}^N \\ & = \left\{ \sum_{n=0}^{\infty} N^{-n/2} \sum_{\substack{(j_1, \dots, j_k) \in \mathbb{N}^k \\ \sum_{i=1}^k j_i = n}} \frac{1}{j_1! \dots j_k!} \langle e_1, \prod_{i=1, \dots, k} (z_i s^+ - \bar{z}_i s^-)^{j_i} e_1 \rangle \right\}^N \end{aligned}$$

Denote R_N the right hand side of (3) with the sum in n starting from 3 rather than 0. Then

$$\begin{aligned} |R_N| & \leq \sum_{n=3}^{\infty} N^{-n/2} \sum_{\substack{j_1, \dots, j_k \\ \sum_i j_i = n}} \frac{1}{j_1! \dots j_k!} \left\| \prod_{i=1, \dots, k} (z_i s^+ - \bar{z}_i s^-)^{j_i} \right\| \leq \quad (4) \\ & \leq \sum_{n=3}^{\infty} N^{-n/2} \sum \frac{1}{j_i! \dots j_k!} |z_1|^{j_1} \dots |z_k|^{j_k} \leq \frac{2e^{|z_1|} \dots e^{|z_k|}}{N^{3/2}} = o(N^{-1/2}) \end{aligned}$$

For $n = 0$ there is only one term equal to 1; for $n = 1$ the contribution is zero, since $\langle e_1, S^\pm e_i \rangle = 0$. The term $n = 2$ is equal to

$$\begin{aligned} & \frac{1}{N} \sum_{\substack{(j_1, \dots, j_k) \in \mathbf{N}^k \\ \sum_i j_i = 2}} \frac{1}{j_1! \dots j_k!} \langle e_1, \prod_{i=1}^{\rightarrow} (z_i s^+ - \bar{z}_i s^-)^{j_i} e_1 \rangle = \\ & = \frac{1}{N} \left\{ \sum_{i=1}^k \frac{1}{2} \langle e_1, (z_i s^+ - \bar{z}_i s^-)^2 e_1 \rangle + \sum_{i < j} \langle e_1, (z_i s^+ - \bar{z}_i s^-) (z_j s^+ - \bar{z}_j s^-) e_1 \rangle \right\} = \\ & = \frac{1}{N} \left(\frac{1}{2} \sum_{i=1}^k -\bar{z}_i z_i + \sum_{i < j} -z_j \bar{z}_i \right) \end{aligned}$$

In conclusion we obtain

$$\begin{aligned} & \langle \varphi_N(0), W_N(z_1), \dots, W_n(z_k) \varphi_N(0) \rangle \\ & = \left\{ 1 + \frac{1}{N} \left[-\frac{1}{2} \sum_{i=1}^k |z_i|^2 - \sum_{i < j} \bar{z}_i z_j \right] + o(N^{-1/2}) \right\}^N \end{aligned} \quad (5)$$

and the algebraic identity

$$-\frac{1}{2} \sum_i |z_i|^2 - \sum_{i < j} \bar{z}_i z_j = -\frac{1}{2} \left| \sum_i z_i \right|^2 - i \operatorname{Im} \sum_{i < j} z_i \bar{z}_j$$

shows that the right hand (side of (5)) converges to the right hand side of (2). The uniformity of the convergence is obvious in view of (4).

3 d-level systems: 1 copy

Our strategy to deal with d-level systems is to exploit the isomorphism between the symmetric tensor product $(\otimes^{d-1} \mathbf{C})_+$ of $d-1$ copies of \mathbf{C}^2 and \mathbf{C}^d , which intertwines the *Lie* algebra representation of $SO(3; R)$ given by

$$S^\pm = \sum_{k=1}^{d-1} j_k(s^\pm) \quad , \quad \sum_{k=1}^{d-1} j_k(s_3) = S_3$$

with the one of the $\frac{d-1}{2} - th$ irreducible representation. This allows to reduce the d -dimensional case to the 2-dimensional one. This isomorphism is described as follows: let

$$e_1 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}; \quad e_2 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

then the symmetric tensor product $(\otimes^{d-1} \mathbf{C}^2)_+$ is a d -dimensional complex vector space with basis given by the $(d-1)$ -number vectors:

$$\varphi_{(d-1)}(k) = \binom{d-1}{k}^{1/2} \text{symm} (\otimes^{d-1-k} e_1 \otimes^k e_2); \quad k = 0, \dots, d-1 \quad (6)$$

where $\text{symm}(\cdot)$ denotes the projection onto the symmetric tensor product. The action of the operators S^\pm, S_3 on this basis is given by:

$$S^+ \varphi_{d-1}(j) = \sqrt{(j+1)(d-1-j)} \varphi_{d-1}(j+1) \quad (7)$$

$$S^- \varphi_{d-1}(j) = \sqrt{j(d-j)} \varphi_{d-1}(j-1) \quad (8)$$

$$S_3 \varphi_{d-1}(j) = (d-1-2j) \varphi_{d-1}(j) \quad (9)$$

Since, by obvious dimensional reasons, the representation is irreducible, it follows that it is isomorphic to the $(d-1)/2$ -th representation of $SO(3)$ (cf. [2],[3]).

Now, for $W_0(z)$ as in section (2) ($z = |z| e^{i\alpha} \in \mathbf{C}$), we denote

$$W_{d-1}(z) = \otimes^{d-1} W_0(z)$$

and

$$\begin{aligned} \psi(z) &= W_{d-1}(z) \varphi_{d-1}(0) = \otimes_{j=1}^{d-1} [\exp(zs^+ - \bar{z}s^-) e_1] = \quad (10) \\ &= \otimes^{d-1} (e_1 \cos |z| + e_2 e^{i\alpha} \sin |z|) = \\ &= \sum_{j=0}^{d-1} \binom{d-1}{j}^{1/2} (\sin |z| e^{i\alpha})^j (\cos |z|)^{d-j-1} \varphi_{d-1}(j) \end{aligned}$$

From this one easily deduces that, uniformly for z, z' in a bounded set :

$$\left\langle \psi \left(\frac{z}{\sqrt{n}} \right), \psi \left(\frac{z'}{\sqrt{n}} \right) \right\rangle = \left(1 + o \left(\frac{1}{n} \right) \right)^{d-1} \quad (11)$$

Lemma 1 *Uniformly for $z, z' \in$ a bounded subset of \mathbf{C} :*

$$\sqrt{N} \langle \psi \left(\frac{z}{\sqrt{N}} \right), S^+ \psi \left(\frac{z'}{\sqrt{N}} \right) \rangle \rightarrow (d-1) \bar{z} \quad (12)$$

$$\sqrt{N} \langle \psi(z/\sqrt{N}), S^- \psi(z'/\sqrt{N}) \rangle \rightarrow (d-1)z' \quad (13)$$

Proof. From (7), (8) one deduces that the left hand side of (12) is equal to:

$$\begin{aligned} & \sqrt{N} \sum_{j,j'} \binom{d-1}{j}^{1/2} \binom{d-1}{j'}^{1/2} \left(\sin \frac{|z|}{\sqrt{N}} e^{i\alpha} \right)^j \left(\cos \frac{|z|}{\sqrt{N}} \right)^{d-1-j} \\ & \left(\sin \frac{|z'|}{\sqrt{N}} e^{i\alpha'} \right)^{j'} \left(\frac{\cos |z'|}{\sqrt{N}} \right)^{d-1-j'} \sqrt{(j'+1)(d-1-j')} \delta_{j,j'+1} \end{aligned} \quad (14)$$

and the left hand side of (13) is equal to:

$$\begin{aligned} & \sqrt{N} \sum_{j,j'} \binom{d-1}{j}^{1/2} \binom{d-1}{j'}^{1/2} \left(\sin \frac{|z|}{\sqrt{N}} e^{i\alpha} \right)^j \left(\cos \frac{|z|}{\sqrt{N}} \right)^{d-1-j} \\ & \left(\sin \frac{|z'|}{\sqrt{N}} e^{i\alpha'} \right)^{j'} \left(\frac{\cos |z'|}{\sqrt{N}} \right)^{d-1-j'} \sqrt{j'(d-j)} \delta_{j,j'-1} \end{aligned} \quad (15)$$

therefore, in the limit $N \rightarrow \infty$, the only surviving term in (??) is the one with $j = 1$, i.e.

$$\binom{d-1}{1}^{1/2} \sqrt{d-1} |z| e^{-i\alpha} = (d-1) \bar{z}$$

while in (??) the only surviving term is the one with $j = 0$ and this gives

$$\binom{d-1}{1}^{1/2} \sqrt{d-1} |z'| e^{j\alpha'} = (d-1)z$$

4 d-level systems: N copies

Define the normalized N -coherent vectors by:

$$W_N(z) = \otimes^N W_o \left(\frac{z}{\sqrt{N}} \right)$$

$$\psi_N(z) = [\otimes^N W_o \left(\frac{z}{\sqrt{N}} \right)] \otimes^N \varphi_{d-1}(0) = \otimes^N \psi \left(\frac{z}{\sqrt{N}} \right)$$

Lemma 2

$$\langle \psi_N(z), \psi_N(z') \rangle \rightarrow \langle \otimes^{d-1} \psi(z), \otimes^{d-1} \psi(z') \rangle \quad (16)$$

Proof.

$$\begin{aligned} \langle \psi_N(z), \psi_N(z') \rangle &= \langle \psi \left(\frac{z}{\sqrt{N}} \right), \psi \left(\frac{z'}{\sqrt{N}} \right) \rangle^N = \\ &= \left\{ \sum_{j=0}^{d-1} \binom{d-1}{j} \left(\sin \frac{|z|}{\sqrt{N}} e^{-i\alpha} \sin \frac{|z'|}{\sqrt{N}} e^{-ia'} \right)^j \left(\cos \frac{|z|}{\sqrt{N}} \cos \frac{|z'|}{\sqrt{N}} \right)^{d-1-j} \right\}^N = \\ &= \left\{ \sum_{j=0}^{d-1} \binom{d-1}{j} \left(\frac{\bar{z}z}{N} + o(N) \right)^j \left(1 - \frac{|z|^2 + |z'|^2}{2N} + o(N) \right)^{d-1-j} \right\}^N = \\ &= \left\{ 1 - \frac{|z|^2 + |z'|^2}{2N} + \frac{\bar{z}z}{N} + o(N) \right\}^{N(d-1)} \rightarrow \langle \psi(z), \psi(z') \rangle^{d-1} \end{aligned}$$

For $u, v \in \mathbf{C}$, we set

$$E_N : M(d; \mathbf{C}) \rightarrow \mathbf{C}$$

$$E_N(b) = \langle \psi \left(\frac{v}{\sqrt{N}} \right), \psi \left(\frac{u}{\sqrt{N}} \right) \rangle^{-1} \langle \psi \left(\frac{u}{\sqrt{N}} \right), b \psi \left(\frac{v}{\sqrt{N}} \right) \rangle \quad (17)$$

From (11) we deduce that

$$E_N(b) \rightarrow \langle \varphi_{d-1}(0), b \varphi_{d-1}(0) \rangle$$

Theorem 2 For any $u, v, z \in \mathbf{C}$

$$\begin{aligned} &\langle \psi_N(u), \exp \left(z \frac{S_N(S^+)}{\sqrt{N}} - z \frac{S_N(S^-)}{\sqrt{N}} \right) \psi_N(v) \rangle \\ &\quad \rightarrow \langle \psi(u), W(z) \psi(v) \rangle^{d-1} \\ &= \left\{ \exp \left(-\frac{|z|^2}{2} \right) \exp(z\bar{u} - \bar{z}v) \right\}^{d-1} \cdot \left(\exp \left(-\frac{|u|^2 + |v|^2}{2} + \bar{u}v \right) \right)^{d-1} \end{aligned}$$

Proof

$$\begin{aligned}
& \langle \psi_N(u), \exp \left(z \frac{S_N(S^+)}{\sqrt{N}} - z \frac{S_N(S^-)}{\sqrt{N}} \right) \psi_N(v) \rangle \\
&= \langle \psi_N(u), \exp \left[z \frac{1}{\sqrt{N}} (\otimes^N E_N) (S_N(S^+)) - z \frac{1}{\sqrt{N}} (\otimes^N E_N) (S_N(S^-)) \right] \\
&\quad \exp \left[z \frac{\hat{S}(S^+)}{\sqrt{N}} - z \frac{\hat{S}(S^-)}{\sqrt{N}} \right] \psi_N(v) \rangle \\
&\quad = \langle \psi_N(u), \psi_N(v) \rangle \cdot \\
&\quad \cdot \exp \left\{ \frac{z}{\sqrt{N}} (\otimes^N E_N) (S_N(S^+)) - \frac{\bar{z}}{\sqrt{N}} (\otimes^N E_N) (S_N(S^-)) \right\} \\
&\quad \cdot (\otimes^N E_N) \left\{ \exp \left(z \frac{\hat{S}_N(S^+)}{\sqrt{N}} - z \frac{\hat{S}_N(S^-)}{\sqrt{N}} \right) \right\} = I \cdot II \cdot III
\end{aligned}$$

We know from Lemma 2:

$$I = \langle \psi_N(u), \psi_N(v) \rangle \rightarrow \langle \psi(u), \psi(v) \rangle^{d-1}$$

Moreover

$$\begin{aligned}
& \frac{z}{\sqrt{N}} (\otimes^N E_N) (S_N(S^+)) - \frac{\bar{z}}{\sqrt{N}} (\otimes^N E_N) (S_N(S^-)) \\
&= z\sqrt{N} \langle \psi \left(\frac{u}{\sqrt{N}} \right), \psi \left(\frac{v}{\sqrt{N}} \right) \rangle^{-1} \langle \psi \left(\frac{u}{\sqrt{N}} \right), S^+ \psi \left(\frac{v}{\sqrt{N}} \right) \rangle \\
&\quad - \bar{z}\sqrt{N} \langle \psi \left(\frac{u}{\sqrt{N}} \right), \psi \left(\frac{v}{\sqrt{N}} \right) \rangle^{-1} \langle \psi \left(\frac{u}{\sqrt{N}} \right), S^- \psi \left(\frac{v}{\sqrt{N}} \right) \rangle \\
&\quad \rightarrow (z\bar{u} - \bar{z}v)(d-1)
\end{aligned}$$

and, because of Lemma 1 and 2

$$II \rightarrow (\exp(z\bar{u} - \bar{z}v))^{d-1}$$

For III we have (cf. Theorem (5.1) of [1])

$$(\otimes^N E_N) \exp \left(z \frac{\hat{S}_N(S^+)}{\sqrt{N}} - \bar{z} \frac{\hat{S}_N(S^-)}{\sqrt{N}} \right)$$

$$= \sum_k \frac{1}{k!} \sum_\varepsilon z^{\varepsilon(1)} \dots z^{\varepsilon(k)} (\otimes^N E_N) \left(\frac{\hat{S}_N(S^{\varepsilon(1)})}{\sqrt{N}} \dots \frac{\hat{S}_N(\varepsilon^{(k)})}{\sqrt{N}} \right)$$

and as,

$$\langle \varphi_{d-1}(0), S^+ \varphi_{d-1}(0) \rangle = 0$$

$$\langle \varphi_{d-1}(0), S^- \varphi_{d-1}(0) \rangle = 0$$

we have:

$$\langle \varphi_{d-1}(0), \hat{S}^+ \hat{S}^+ \varphi_{d-1} \rangle = 0$$

$$\langle \varphi_{d-1}(0), \hat{S}^+ \hat{S}^- \varphi_{d-1} \rangle = 0$$

$$\langle \varphi_{d-1}(0), \hat{S}^- \hat{S}^- \varphi_{d-1} \rangle = 0$$

$$\begin{aligned} \langle \varphi_{d-1}(0), \hat{S}^- \hat{S}^+ \varphi_{d-1} \rangle &= \langle \varphi_{d-1}(0), \hat{S}^- \hat{S}^+ \varphi_{d-1}(0) \rangle = |S^+ \varphi_{d-1}(0)|^2 \\ &= |\sqrt{(d-1)} \varphi_{d-1}(1)|^2 = (d-1) \end{aligned}$$

Therefore:

$$III \rightarrow \sum_{p=0} \frac{1}{(2p)!} \frac{(2p)!}{p!2^p} (-|z|^2 (d-1))^p = \left(\exp \left(-\frac{|z|^2}{2} \right) \right)^{d-1} \quad (18)$$

5 Inclusion of the number operator

Theorem 3 For any $u, v, z, \in \mathbf{C}$ and any $\lambda \in \mathbf{R}$

$$\begin{aligned} \lim_{N \rightarrow \infty} \langle \psi_N \left(\frac{u}{\sqrt{N}} \right), \exp \left\{ z \frac{S_N(S^+)}{\sqrt{N}} - \bar{z} \frac{S_N(S^-)}{\sqrt{N}} + i2\lambda S_N(n^+) \right\} \psi_N \left(\frac{u}{\sqrt{N}} \right) \rangle \\ = \langle \psi(u), \exp(za^+ - \bar{z}a + i2\lambda a^+ a) \psi(v) \rangle \end{aligned} \quad (19)$$

Proof. We first evaluate the right hand side of equation (??). For $\lambda = 0$ we have

$$\begin{aligned} \langle \psi(u), W(z), \psi(v) \rangle &= \exp \left(-\frac{\bar{z}v}{z} + \frac{z\bar{v}}{z} \right) \langle \psi(u), \psi(z+u) \rangle \\ &= \exp \left(-\frac{|u|^2}{2} - \frac{|v|^2}{2} - \frac{|z|^2}{2} - \frac{\bar{z}v}{2} - \frac{z\bar{v}}{2} + \bar{u}z + \bar{u}v - \frac{\bar{z}v}{2} - \frac{z\bar{v}}{2} \right) \\ &= \exp \left(-\frac{|u|^2}{2} - \frac{|v|^2}{2} - \frac{|z|^2}{2} - \bar{z}v + \bar{u}z + \bar{u}v \right) \end{aligned} \quad (20)$$

Assume now $\lambda \neq 0$, then

$$\begin{aligned}
& \exp(z a^+ - \bar{z} a + i2\lambda a^+ a) \\
&= \exp\left(i2\lambda\left(aa^+ + \frac{z}{2i\lambda}a^+ - \frac{\bar{z}}{2i\lambda}a\right)\right) \\
&= \exp\left\{i2\lambda\left(a^+ - \frac{\bar{z}}{2i\lambda}\right)\left(a + \frac{z}{2i\lambda}\right) - i2\lambda\frac{|z|^2}{4\lambda^2}\right\} \\
&= \exp(-i2\lambda|\zeta|^2) \exp\{i2\lambda(a^+ + \bar{\zeta})(a + \zeta)\} \tag{21}
\end{aligned}$$

where

$$\zeta := \frac{z}{i2\lambda}, \quad \bar{\zeta} = -\frac{\bar{z}}{i2\lambda}, \quad |\zeta|^2 = \frac{|z|^2}{4\lambda^2} \tag{22}$$

therefore,

$$\exp(z a^+ - \bar{z} a + i2\lambda a^+ a) = \exp(-i2\lambda|\zeta|^2) W^+(\zeta) \exp(i2\lambda a^+ a) W(\zeta) \tag{23}$$

and for the matrix elements we obtain

$$\begin{aligned}
& \langle \psi(u), \exp(z a^+ - \bar{z} a + i2\lambda a^+ a) \psi(v) \rangle = \tag{24} \\
&= \exp(-i2\lambda|\zeta|^2) \exp\left(-\frac{\zeta\bar{u}}{2} + \frac{\bar{\zeta}u}{2} - \frac{\bar{\zeta}v}{2} + \frac{\zeta\bar{v}}{2}\right) \langle \psi(u+S), \exp(i2\lambda a^+ a) \psi(v+\zeta) \rangle
\end{aligned}$$

Using the identity

$$\exp(\alpha a^+ a) = \sum_k \frac{(e^\alpha - 1)^k}{k!} a^{+k} a^k \tag{25}$$

we obtain

$$\begin{aligned}
& \langle \psi(u), \exp(z a + i2\lambda a^+ a) \psi(v) \rangle \tag{26} \\
& \exp\left(-i2\lambda|\zeta|^2 - \zeta\bar{u} - \bar{\zeta}v - \frac{|v|^2}{2} - \frac{|u|^2}{2} - |\zeta|^2 + e^{2i\lambda}(\bar{u} + \bar{\zeta})(v + \zeta)\right) \\
&= \exp\left(-i2\lambda|\zeta|^2 - |\zeta|^2 + e^{-i2\lambda}|\zeta|^2 - \frac{|u|^2}{2} - \frac{|v|^2}{2} - \zeta\bar{u} - \bar{\zeta}v + e^{2i\lambda}\bar{u}\zeta + e^{2i\lambda}\bar{\zeta}v\right)
\end{aligned}$$

Introducing

$$w = v e^{2i\lambda}$$

we have

$$\begin{aligned} & \langle \psi(u), \exp(z a + -\bar{z} a + i2\lambda a^+ a) \psi(v) \rangle = \quad (27) \\ & = \exp\left\{ (e^{2i\lambda} - 2i\lambda - 1) |\zeta|^2 - \frac{|u|^2}{2} - |w|^2 + \bar{u}w + (e^{2i\lambda} - 1) \bar{u}\zeta + (1 - e^{-2i\lambda}) \bar{\zeta}w \right\} \end{aligned}$$

Transforming back to z, \bar{z} we obtain

$$\begin{aligned} & \langle \psi(u), \exp(z a^+ - \bar{z} a + 2i\lambda a^+ a) \psi(v) \rangle = \quad (28) \\ & = \exp\left\{ \frac{e^{2i\lambda} - 2i\lambda - 1}{4\lambda^2} |z|^2 - \frac{|u|^2}{2} - \frac{|w|^2}{2} + \bar{u}w + \frac{e^{2i\lambda} - 1}{i2\lambda} \bar{u}z - \frac{e^{2i\lambda} - 1}{i2\lambda} \bar{z}w \right\} \end{aligned}$$

In the limit $\lambda \rightarrow 0$ we have

$$w = v e^{i2\lambda} \rightarrow v ; \quad \frac{e^{2i\lambda} - 1}{i2\lambda} \rightarrow 1 , \quad \frac{e^{-2i\lambda} - 1}{-i2\lambda} \rightarrow 1 , \quad \frac{e^{2i\lambda} - 2i\lambda - 1}{4\lambda^2} \rightarrow \frac{1}{2}$$

so that (??) is valid for all values of $\lambda \in \mathbf{R}$. In order to evaluate the left hand side of (??), we notice that

$$\begin{aligned} & \langle \psi_N\left(\frac{u}{\sqrt{N}}\right), \exp\left(z \frac{S_N(s^+)}{\sqrt{N}} - \bar{z} \frac{S_N(s^-)}{\sqrt{N}} + 2i\lambda S_N(n^+)\right) \psi_N\left(\frac{v}{\sqrt{N}}\right) \rangle = \\ & = \langle \psi\left(\frac{u}{\sqrt{N}}\right), \exp\left(\frac{z}{\sqrt{N}} s^+ - \frac{\bar{z}}{\sqrt{N}} s^- + 2i\lambda n^+\right) \psi\left(\frac{v}{\sqrt{N}}\right) \rangle^N \quad (29) \end{aligned}$$

We may use $n^+ = \frac{1}{2}(\sigma^3 + 1)$ and, denoting

$$\underline{t} = \left(\operatorname{Im} \frac{z}{\sqrt{N}}, \operatorname{Re} \frac{z}{\sqrt{N}}, \lambda \right) \in \mathbf{R}^3$$

we obtain

$$\exp(i \underline{t} \times \underline{\sigma}) = \cos(|t|) \underline{u} + i \frac{\underline{t} \cdot \underline{\sigma}}{|t|} \sin(|t|) \quad (30)$$

$$\exp\left(\frac{z}{\sqrt{N}} s^+ - \frac{\bar{z}}{\sqrt{N}} s^- + 2i\lambda n^+\right) = \exp(i\lambda) \exp(i \underline{t} \cdot \underline{\sigma}) \quad (31)$$

Since

$$|t|^2 = \frac{|z|^2}{N} + \lambda^2 \quad , \quad |t| = \lambda \sqrt{\frac{|z|^2}{\lambda^2 N} + 1} \quad (32)$$

We can collect the leading terms in (??) finding (in obvious notations)

$$\underline{1} : \quad \cos(|t|) = \cos \lambda - \frac{|z|^2}{2\lambda N} \sin \lambda + O\left(\frac{1}{N^2}\right) \quad (33)$$

$$\sigma_3 : \quad \lambda \frac{\sin(|t|)}{|t|} = \sin \lambda + \frac{|z|^2}{2\lambda N} \cos \lambda - \frac{|z|^2}{2\lambda^2 N} \sin \lambda + O\left(\frac{1}{N^2}\right) \quad (34)$$

$$\sigma_1 : \quad \Im \frac{z}{\sqrt{N}} \cdot \frac{\sin(|t|)}{|t|} = \left(\Im \frac{z}{\sqrt{N}} \right) \frac{\sin \lambda}{\lambda} + O(N^{-3/2}) \quad (35)$$

$$\sigma_2 : \quad \operatorname{Re} \frac{z}{\sqrt{N}} \cdot \frac{\sin(|t|)}{|t|} = \left(\operatorname{Re} \frac{z}{\sqrt{N}} \right) \frac{\sin \lambda}{\lambda} + O(N^{-3/2}) \quad (36)$$

Therefore,

$$\begin{aligned} & \exp\left(\frac{z}{\sqrt{N}}s^+ - \frac{\bar{z}}{\sqrt{N}}s^- + 2i\lambda n^+\right) \cong \\ & \cong e^{i\lambda} \begin{pmatrix} e^{i\lambda} + i\frac{|z|^2}{2\lambda N} \left(e^{i\lambda} - \frac{\sin \lambda}{\lambda}\right) & \frac{\sin \lambda}{\lambda} \frac{z}{\sqrt{N}} \\ -\frac{\sin \lambda}{\lambda} \frac{\bar{z}}{\sqrt{N}} & e^{i\lambda} - i\frac{|z|^2}{2\lambda N} \left(e^{i\lambda} - \frac{\sin \lambda}{\lambda}\right) \end{pmatrix} \end{aligned} \quad (37)$$

Moreover,

$$\psi\left(\frac{v}{\sqrt{N}}\right) = \begin{pmatrix} \frac{v}{\sqrt{N}} \\ \sqrt{1 - \frac{|v|^2}{N}} \end{pmatrix} \cong \begin{pmatrix} \frac{v}{\sqrt{N}} \\ 1 - \frac{1}{2} \frac{|v|^2}{N} \end{pmatrix} \quad (38)$$

and analogously, for $\psi(v/\sqrt{N})$. In the limit $\lambda \rightarrow 0$ we have

$$\frac{e^{\pm i\lambda} - \frac{\sin \lambda}{\lambda}}{\lambda} \rightarrow \pm i \quad (39)$$

so that the matrix (??) is reduced (up to terms of order $o(1/N)$) to

$$\begin{pmatrix} 1 - \frac{|z|^2}{2N} & \frac{z}{\sqrt{N}} \\ -\frac{\bar{z}}{\sqrt{N}} & 1 - \frac{|z|^2}{2N} \end{pmatrix} \cong \exp\left(\frac{z}{\sqrt{N}}s^+ - \frac{\bar{z}}{\sqrt{N}}s^-\right) \quad (40)$$

For the matrix elements we obtain in the limit $\lambda \rightarrow 0$

$$\begin{aligned} & \langle \psi\left(\frac{u}{\sqrt{N}}\right), \exp\left(\frac{z}{\sqrt{N}}S^+ - \frac{\bar{z}}{\sqrt{N}}S^-\right) \psi\left(\frac{v}{\sqrt{N}}\right) \rangle \\ & 1 - \frac{|z|^2}{2N} - \frac{|v|^2}{2N} - \frac{|u|^2}{2N} + \frac{\bar{u}v}{N} + \frac{\bar{u}z}{N} - \frac{\bar{z}v}{N} \end{aligned} \quad (41)$$

which, as $N \rightarrow \infty$, converges to

$$\langle \psi(u), \exp(za^+ - \bar{z}a)\psi(v) \rangle = \exp\left(-\frac{|z|^2}{2} - \frac{|u|^2}{2} - \frac{|v|^2}{2} + \bar{u}v + \bar{u}z - \bar{z}v\right) \quad (42)$$

We now evaluate the matrix elements of (??). This gives

$$\begin{aligned}
& \frac{\bar{u}v}{N}e^{2i\lambda} + \frac{\sin \lambda}{\lambda} \frac{z\bar{u}}{N}e^{i\lambda} - \frac{\sin \lambda}{\lambda} \frac{\bar{z}v}{N}e^{i\lambda} + 1 \\
& -i\frac{|z|^2}{2N\lambda}e^{i\lambda} \left(e^{-i\lambda} - \frac{\sin \lambda}{\lambda} \right) - \frac{|v|^2}{2N} - \frac{|u|^2}{2N} \\
& \cong \langle \psi \left(\frac{u}{\sqrt{N}} \right) \exp \left(\frac{z}{\sqrt{N}}s^+ - \frac{\bar{z}}{\sqrt{N}}s^- + 2i\lambda n^+ \right) \psi \left(\frac{v}{\sqrt{N}} \right) \rangle
\end{aligned} \tag{43}$$

Therefore

$$\langle \psi \left(\frac{u}{\sqrt{N}} \right) \exp \left(z\frac{S_N(s^+)}{\sqrt{N}} - \bar{z}\frac{S_N(s^-)}{\sqrt{N}} + 2i\lambda S_N(n^+) \right) \psi \left(\frac{v}{\sqrt{N}} \right) \rangle$$

converges to

$$\exp \left\{ -\frac{|u|^2}{2} - \frac{|v|^2}{2} - i\frac{|z|^2}{2} e^{i\lambda} \frac{e^{i\lambda} - \frac{\sin \lambda}{\lambda}}{\lambda} + \bar{u}v e^{2i\lambda} + \frac{\sin \lambda}{\lambda} z + \bar{u}e^{i\lambda} - \frac{\sin \lambda}{\lambda} \bar{z}v e^{i\lambda} \right\} = \tag{44}$$

$$= \exp \left\{ -\frac{|u|^2}{2} - \frac{|w|^2}{2} - i\frac{|z|^2}{2} \frac{e^{i\lambda}}{\lambda} \left(e^{-i\lambda} - \frac{\sin \lambda}{\lambda} \right) + \bar{u}w + \bar{u}z e^{i\lambda} \frac{\sin \lambda}{\lambda} - \bar{z}e^{-i\lambda} \frac{\sin \lambda}{\lambda} \right\} \tag{45}$$

Finally comparing

$$\begin{aligned}
\frac{e^{2i\lambda} - 1}{2i\lambda} &= e^{i\lambda} \frac{e^{i\lambda} - e^{-i\lambda}}{2i\lambda} = e^{i\lambda} \frac{\sin \lambda}{\lambda} \\
\frac{e^{-2i\lambda} - 1}{-2i\lambda} &= e^{-i\lambda} \frac{e^{-i\lambda} - e^{i\lambda}}{-2i\lambda} = e^{-i\lambda} \frac{\sin \lambda}{\lambda} \\
\frac{e^{2i\lambda} - 2i\lambda - 1}{4\lambda^2} &= \frac{e^{i\lambda}}{2i\lambda} \left\{ e^{i\lambda} - 2i\lambda e^{-i\lambda} - \frac{e^{-i\lambda}}{-2i\lambda} \right\} = -i\frac{e^{i\lambda}}{2\lambda} \left(e^{i\lambda} - \frac{\sin \lambda}{\lambda} \right)
\end{aligned}$$

with (??), we obtain the thesis.

6 Squeezing states

Let a, a^+ be a representation of the CCR on a certain invariant domain D in a Hilbert space H and vacuum vector Φ .

Definition 1 A vector χ is called a squeezing vector for the representation $\{a^\pm, D \subseteq H, \Phi\}$ if it can be represented in the form

$$\chi = e^{(za^{+2} - \bar{z}a^2)} W(z') \Phi \quad (46)$$

The unitary operator $\exp(za^{+2} - \bar{z}a^2)$ is called the squeezing operator. The exponential in (46) is defined by its power series on the number vectors, which are analytic for $(a^\pm)^2$.

Theorem 4 For any u, v, z, \in, \mathbf{C}

$$\begin{aligned} \lim \left\langle \psi_N \left(\frac{u}{\sqrt{N}} \right), \exp \left\{ z \left(\frac{S_N(s^+)}{\sqrt{N}} \right) - \bar{z} \left(\frac{S_N(s^-)}{\sqrt{N}} \right)^2 \right\} \psi_N \left(\frac{v}{\sqrt{N}} \right) \right\rangle \\ = \langle \psi(u), \exp(za^{+2} - \bar{z}a^2) \psi(v) \rangle \end{aligned} \quad (47)$$

The proof of the theorem consists of two steps first to compute explicitly the matrix elements of the right hand side of (47) in the coherent vectors; then to prove that the limit on the left hand side exists and the equality holds. We shall only outline the first step, while we give the full proof of the second one.

The explicit computation of the right hand side of (47) is based on the formula

$$e^{\frac{1}{2}(za^{+2} - \bar{z}a^2)} = e^{\frac{1}{2}e^{i\alpha} \tanh |z| a^{+2}} \cdot e^{-\log(\cosh |z|)(a^+ a + 1/2)} \cdot e^{-\frac{1}{2}e^{-i\alpha} \tanh |z| a^2} \quad (48)$$

which implies:

$$\begin{aligned} \langle \psi(u), e^{\frac{1}{2}(za^{+2} - \bar{z}a^2)} \psi(v) \rangle = \\ = \exp \left(-\frac{|u|^2 + |v|^2}{2} + \frac{1}{2}e^{i\alpha} \bar{u}^2 \tanh |z| - \frac{1}{2}e^{-i\alpha} \bar{v}^2 \tanh |z| - \left(\frac{1}{2} + \bar{u}v \right) \log \cosh |z| \right) \end{aligned} \quad (49)$$

To control the limit on the left hand side of (47), we expand the exponential and show, in the following lemma, that the limit and the summation sign can be interchanged. To this goal we introduce the following notation: for any $u, v, z, \in, \mathbf{C}, N \in \mathbf{N}$ we set

$$a_{k,N} = \frac{1}{k!} \left\langle \psi_N \left(\frac{u}{\sqrt{N}} \right), \left\{ z \frac{S_N^2(s^+)}{N} - \bar{z} \frac{S_N^2(s^-)}{N} \right\}^k \psi_N \left(\frac{v}{\sqrt{N}} \right) \right\rangle$$

Lemma 3 For any $\varepsilon > 0$ there exists N_ε and h_ε such that for any $N > N_\varepsilon$ and any $h > h_\varepsilon$

$$\sum_{k=h}^{\infty} |a_{k,N}| < \varepsilon$$

Proof. We set

$$f_N : b \in M(2; \mathbf{C}) \rightarrow f_N(b) = \langle \psi \left(\frac{u}{\sqrt{N}} \right), b \psi \left(\frac{v}{\sqrt{N}} \right) \rangle \in \mathbf{C}$$

and

$$E_N : b \in \otimes^N M(2; \mathbf{C}) \rightarrow E_N(b) = \frac{(\otimes^N f_N)(b)}{(\otimes^N f_N)(1_N)} \in \mathbf{C}$$

With this notation

$$\begin{aligned} a_{k,N} &= \frac{1}{k!} (\otimes^N f_N)(1_N) E_N \left(z \frac{S_N^2(s^+)}{N} - \bar{z} \frac{S_N^2(s^-)}{N} \right)^k \\ &= \frac{1}{k!} (\otimes^N f_N)(1_N) E_N \left(\sum_{\varepsilon=1}^6 \gamma_N^{(\varepsilon)} J_N^{(\varepsilon)} \right)^k \end{aligned}$$

where

$$\begin{aligned} \gamma_N^{(1)} &= z \frac{E_N^2(S_N(s^+))}{N}, \quad J_N^{(1)} = 1_N \\ \gamma_N^{(2)} &= -\bar{z} \frac{E_N^2(S_N(s^-))}{N}, \quad J_N^{(2)} = 1_N \\ \gamma_N^{(3)} &= 2z \frac{E_N(S_N(s^+))}{\sqrt{N}}, \quad J_N^{(3)} = \frac{S_N^\wedge(s^+)}{\sqrt{N}} \\ \gamma_N^{(4)} &= -2\bar{z} \frac{E_N(S_N(s^-))}{\sqrt{N}}, \quad J_N^{(4)} = \frac{S_N^\wedge(s^-)}{\sqrt{N}} \\ \gamma_N^{(5)} &= z, \quad J_N^{(5)} = \frac{S_N^{\wedge 2}(s^+)}{N} \\ \gamma_N^{(6)} &= -\bar{z}, \quad J_N^{(6)} = \frac{S_N^{\wedge 2}(s^-)}{N} \end{aligned}$$

and where we have used the notation (cf [1])

$$\hat{X} = X - E_N(X)$$

With these notations:

$$a_{k,n} = \frac{1}{k!} (\otimes^N f_N) (1_N) \\ \sum_{(\varepsilon_1, \dots, \varepsilon_k) \in \{1, \dots, 6\}^k} \gamma_N^{(\varepsilon_1)} \dots \gamma_N^{\varepsilon_k} E_N \left(J_N^{(\varepsilon_1)} \dots J_N^{(\varepsilon_k)} \right)$$

We know that

$$|(\otimes^N f_N) (1_N)| \leq 1 \quad ; \quad \left| \frac{E_N(S_N(s^\pm))}{\sqrt{N}} \right| \leq C_1 + \frac{C_2}{N} \quad (50)$$

Therefore, for some $(\varepsilon_1, \dots, \varepsilon_k) \in \{1, \dots, 6\}^k$ one has

$$|a_{k,n}| \leq \frac{1}{k!} 6^k C_N^k |E_N \left(J_N^{(\varepsilon_1)} \dots J_N^{(\varepsilon_k)} \right)|$$

where, because of (50)

$$C_N = C_3 + \frac{C_4}{N}$$

and C_3 depends upon \bar{u}, v, z, \bar{z} . The products $J_N^{(\varepsilon_1)} \dots J_N^{(\varepsilon_k)}$ consist of a scalar times j ($0 \leq j \in 2k$), terms of the form $\frac{\hat{S}_N(s^\pm)}{\sqrt{N}}$. From the central limit theorem estimate of [1] we know that for any $j \in \mathbf{N}$

$$|E_N \left(\frac{\hat{S}_N(b_i)}{\sqrt{N}} \dots \frac{\hat{S}_N(b_j)}{\sqrt{N}} \right)| \leq C_5 \binom{j}{2} + C_6^j \frac{1}{\sqrt{N}} \leq C_5 \binom{2k}{2} + C_6^{2k} \frac{1}{\sqrt{N}}$$

In conclusion we obtain

$$|a_{k,n}| \leq \frac{1}{k!} 6^k \left(C_3 + \frac{C_4}{N} \right)^k \left\{ C_5 \binom{2k}{2} + C_6^{2k} \frac{1}{\sqrt{N}} \right\} \\ \leq \frac{1}{k!} \left(C_7 + \frac{C_8}{N} \right)^k \frac{1}{\sqrt{N}} + \frac{C_5}{k!} \left(C_9 + \frac{C_{10}}{N} \right)^k \frac{(2k)!}{2(2k-2)!} \\ \leq \frac{1}{k!} \left(C_7 + \frac{C_8}{N} \right)^k \frac{1}{\sqrt{N}} + \frac{(2k)(2k-1)}{k!} C_{11} \left(C_9 + \frac{C_{10}}{N} \right)^k$$

from wih the thesis follows.

References

- [1] Accardi L., Bach A. *The harmonic oscillator as quantum central limit of Bernoulli processes*. to appear in: Theory of Probability and related fields
- [2] Biedenharn L.C., Louck J.D. *Angular Momentum in Quantum Physics*, Addison-Wesley (1981).
- [3] Biedenharn L.C., Louck J.D. *The Racah-Wigner algebra in Quantum theory*, Addison-Wesley (1981).
- [4] Meyer P.A. *Approximation de l'oscillateur harmonique*, Seminaire de Probabilités, Strasbourg 1988/89.
- [5] Parthasarathy K.R. *The passage from random walk to diffusion in quantum probability*. in : Celebration Volume in Applied Probability (ed. J.Gani), Applied Probability Trust, Sheffield, U.K. 231-245, to appear