

Provided for non-commercial research and education use.
Not for reproduction, distribution or commercial use.



(This is a sample cover image for this issue. The actual cover is not yet available at this time.)

This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



ELSEVIER

Contents lists available at SciVerse ScienceDirect

Journal of Algebra

www.elsevier.com/locate/jalgebra



Kazhdan–Lusztig combinatorics in the moment graph setting

Martina Lanini

Department Mathematik, Friedrich-Alexander-Universität Erlangen-Nürnberg, Cauerstr. 11, 91058 Erlangen, Germany

ARTICLE INFO

Article history:

Received 23 August 2011

Available online xxxx

Communicated by Peter Littelmann

Keywords:

Representation theory

Algebraic combinatorics

Kazhdan–Lusztig polynomials

Moment graphs

ABSTRACT

Motivated by a question on the graded rank of the stalks of the canonical sheaf on a Bruhat graph, we lift some equalities concerning (parabolic) Kazhdan–Lusztig polynomials to this moment graph setting. Our proofs hold also in positive characteristic, under some technical assumptions.

© 2012 Elsevier Inc. All rights reserved.

1. Introduction

In 1979 Kazhdan and Lusztig [20] associated to a given Coxeter group \mathcal{W} a family of polynomials $\{P_{y,w}(q)\}$ indexed by pairs of elements in \mathcal{W} . In the case \mathcal{W} was a Weyl group, then $P_{y,w}(q)$ was related to the local intersection cohomology of the corresponding Schubert variety in a partial flag variety (cf. Appendix A of [20] and [21]). Some years later, Deodhar in [7] introduced the parabolic analogue of Kazhdan–Lusztig polynomials. Namely, if $(\mathcal{W}, \mathcal{S})$ is a Coxeter system, $J \subseteq \mathcal{S}$, and \mathcal{W}^J is the set of minimal coset representatives of $\mathcal{W}/\langle J \rangle$, he defined two families of polynomials $\{P_{y,w}^{J,-1}(q)\}$ and $\{P_{y,w}^{J,q}(q)\}$, where $y, w \in \mathcal{W}^J$. The $\{P_{y,w}^{J,-1}(q)\}$ are a generalization of the polynomials defined by Kazhdan and Lusztig and for $J = \emptyset$ they coincide. As in the regular case, if \mathcal{W} is a Weyl group, then these polynomials have a geometrical meaning, and, in particular, they are related to the intersection cohomology of the corresponding Schubert variety in a partial flag variety.

Kazhdan and Lusztig [20], resp. Lusztig [25], conjectured that the Kazhdan–Lusztig polynomials played a very important role in the representation theory of complex Lie algebras, resp. of semisimple, simply connected, reductive algebraic groups over a field of positive characteristic. The characteristic zero setting is now well understood (cf. [21,2,6]), while the positive characteristic analogue is

E-mail address: lanini@mi.uni-erlangen.de.

not. Actually Lusztig's conjecture was *almost* proved in the 90s via the joint work of Kazhdan and Lusztig [22], Kashiwara and Tanisaki [23] and Andersen, Jantzen and Soergel [1]. Here *almost* means that it was possible to prove the conjecture only if the characteristic of the base field is big enough, since it was obtained as a limit of the characteristic zero case. A new approach to Lusztig's conjecture is due to Fiebig [9,13] and it is based on the theory of sheaves on moment graphs.

Moment graphs were introduced by Goresky, Kottwitz and MacPherson [16], in order to study the equivariant cohomology of a complex algebraic variety equipped with a torus action and having some nice properties. In 2001 Braden and MacPherson in [3] were able to describe the equivariant intersection cohomology of such a variety via sheaves on the moment graph. In particular, if \mathcal{W} is a Weyl group with \mathcal{S} , the set of simple reflections, and $J \subseteq \mathcal{S}$, Braden and MacPherson associated to $w \in \mathcal{W}^J$ a sheaf \mathcal{B}_w^J : the *canonical*–or *BMP*–sheaf. This object describes the local intersection cohomology of the corresponding Schubert variety in a partial flag variety. Braden–MacPherson's construction was performed in characteristic zero, but it is possible to develop this theory in any characteristic. Fiebig and Williamson proved in [15] that, with certain technical assumptions, in positive characteristic \mathcal{B}_w^J computes the stalks of indecomposable parity sheaves (introduced in [18]). It is now natural to ask whether it is possible to connect the canonical sheaf to Kazhdan–Lusztig polynomials.

Question 4.1. (Cf. [11], Conjecture 4.4.) Under which assumptions on the characteristic of the base field, do we have $\underline{\text{rk}}(\mathcal{B}_w^J)^y = P_{y,w}^{J,-1}$ for $y \leq w$ and y, w varying in some relevant subset of \mathcal{W}^J ?

This equality is true in characteristic zero for any pair y, w and in this case it is equivalent to Kazhdan–Lusztig's conjecture (cf. [9]). In characteristic p , if \mathcal{W} is affine and if we only consider w restricted (cf. [13]), it is proved for p bigger than a huge (but explicit) bound (cf. [14]), and, for p bigger than the Coxeter number, it implies Lusztig's conjecture (cf. [11,13]). From a recent result of Polo (private communication, 7 May 2012), it follows that if $\mathcal{W} = S_{4p}$ the stalks of the BMP-sheaf are definitively not given by these polynomials (see Section 4.1 for more details). Anyway, this question motivates our work, since it now makes sense to interpret some equalities concerning (parabolic) Kazhdan–Lusztig polynomials in terms of stalks of the canonical sheaves. In order to lift properties of KL-polynomials to the level of canonical sheaves, we will use two different techniques: the pullback of canonical sheaves (see Section 5) and an action of the Weyl group on the set of global sections of the BMP-sheaf (see Section 6).

Let k be a local ring with $2 \in k^\times$. We define the notion of k -homomorphism between two moment graphs and of pullback of sheaves. These will provide a useful tool, namely, under some assumptions on k ,

Lemma 5.1. *Let \mathcal{G} and \mathcal{G}' be two moment graphs, both with a unique maximal vertex, w resp. w' , and let f be a k -isomorphism between them. If \mathcal{B}_w and $\mathcal{B}_{w'}$ are the corresponding canonical sheaves, then $\mathcal{B}_w \cong f^* \mathcal{B}_{w'}$ as k -sheaves on \mathcal{G} .*

Thanks to this result, in some good situations it will be enough to study the combinatorics of the underlying moment graphs that in our case are just labeled, oriented Bruhat graphs (see Section 2.2). This is the case in the following theorem:

Theorem 5.1. *Let $y, w \in \mathcal{W}$ be such that $y \leq w$, then*

(i) $\mathcal{B}_w^y \cong \mathcal{B}_{w^{-1}}^{y^{-1}}$.

Let $s \in \mathcal{S}$ be such that $ws < w$, then

(ii) *if $y \not\leq ws$, $\mathcal{B}_w^y \cong \mathcal{B}_{ws}^{ys}$,*

where we write \mathcal{B}_w instead of \mathcal{B}_w^\emptyset .

The last part of the paper is devoted to the study of an action of a certain subgroup of the Weyl group \mathcal{W} on the space of global sections of the canonical sheaf and, in particular, to the proof that the data we need to build the canonical sheaf is contained in the invariants with respect to this action. This result, together with some combinatorics of the corresponding Bruhat graph, gives us the categorical analogue of a result due to Kazhdan and Lusztig (cf. [21]):

Theorem 6.1. *Under some assumptions on k , if $y, w \in \mathcal{W}$ and $s \in \mathcal{S}$ are such that $y \leq w$ and $ws < w$, then $\mathcal{B}_w^y \cong \mathcal{B}_w^{ys}$.*

Inspired by a theorem of Deodhar [7] we prove a relation between the canonical sheaf on a regular Bruhat graph \mathcal{G} and the ones on the corresponding parabolic Bruhat graphs \mathcal{G}^J , for J such that the subgroup $\langle J \rangle$ is finite. Let w_J be the longest element of $\langle J \rangle$ then, under some assumptions on k , we have

Theorem 6.2. *If $y, w \in \mathcal{W}^J$, then $(\mathcal{B}_w^J)^y \cong (\mathcal{B}_{ww_J}^\emptyset)^{yw_J}$.*

In order to prove this result we consider again the space of invariants with respect to the above action. The claim follows from the connection between this space and the parabolic canonical sheaf.

Structure of the paper. Sections 2 and 3 are about moment graphs and sheaves on them. In Section 4 we introduce Braden–MacPherson sheaves and recall some of their properties. We develop and apply the technique of pullbacks in Section 5, while the one of invariants is used in the last section.

2. Moment graphs

In this section we recall the definition of moment graphs on a lattice and we define the notion of k -homomorphism between two moment graphs.

Let k be from now on a local ring inside which 2 is an invertible element. Let $Y \cong \mathbb{Z}^r$ be a lattice of finite rank and denote by $Y_k := Y \otimes_{\mathbb{Z}} k$.

Definition 2.1. A moment graph \mathcal{G} on Y is given by $(\mathcal{V}, \mathcal{E}, \leq, l)$, where:

- (i) $(\mathcal{V}, \mathcal{E})$ is a directed graph without directed cycles nor multiple edges,
- (ii) \leq is a partial order on \mathcal{V} such that if $x, y \in \mathcal{V}$ and $E : x \rightarrow y \in \mathcal{E}$ then $x \leq y$,
- (iii) $l : \mathcal{E} \rightarrow Y \setminus \{0\}$ is a map called the *label function*.

Definition 2.2. Let \mathcal{G} be a moment graph on the lattice Y , then

- \mathcal{G} is a k -moment graph on Y if all labels are non-zero in Y_k .
- (\mathcal{G}, k) is a *GKM-pair* if all pairs of distinct edges containing a common vertex have labels k -linearly independent in Y_k .

Observe that if (\mathcal{G}, k) is a *GKM-pair*, then \mathcal{G} is a k -moment graph. These properties are very important and in the sequel they will give a restriction on the ring k .

2.1. k -Homomorphisms of moment graphs

Let $\mathcal{G} = (\mathcal{V}, \mathcal{E}, \leq, l)$ and $\mathcal{G}' = (\mathcal{V}', \mathcal{E}', \leq', l')$ be two moment graphs on Y . Since a moment graph is given by an oriented and ordered graph plus some other data coming from Y , we define a k -homomorphism as a map of graphs plus a collection of automorphisms of the k -module Y_k satisfying certain requirements. More precisely,

Definition 2.3. A k -homomorphism between two moment graphs on Y

$$f : (\mathcal{V}, \mathcal{E}, \triangleleft, l) \rightarrow (\mathcal{V}', \mathcal{E}', \triangleleft', l')$$

is given by $(f_{\mathcal{V}}, (f_{l,x})_{x \in \mathcal{V}})$, where

- (MORPH1) $f_{\mathcal{V}} : \mathcal{V} \rightarrow \mathcal{V}'$ is any (order preserving) map of posets such that, if $x \longrightarrow y \in \mathcal{E}$, then either $f_{\mathcal{V}}(x) \longrightarrow f_{\mathcal{V}}(y) \in \mathcal{E}'$, or $f_{\mathcal{V}}(x) = f_{\mathcal{V}}(y)$.
For an edge $E : x \longrightarrow y \in \mathcal{E}$ such that $f_{\mathcal{V}}(x) \neq f_{\mathcal{V}}(y)$, we will denote $f_{\mathcal{E}}(E) := f_{\mathcal{V}}(x) \longrightarrow f_{\mathcal{V}}(y)$.
- (MORPH2) For all $x \in \mathcal{V}$, $f_{l,x} : Y_k \rightarrow Y_k \in \text{Aut}_k(Y_k)$ is such that, if $E : x \longrightarrow y \in \mathcal{E}$ and $f_{\mathcal{V}}(x) \neq f_{\mathcal{V}}(y)$, the following two conditions are verified:
- (MORPH2a) $f_{l,x}(l(E)) = h \cdot l'(f_{\mathcal{E}}(E))$, for some $h \in k^\times$.
- (MORPH2b) $\pi \circ f_{l,x} = \pi \circ f_{l,y}$, where π is the canonical quotient map $\pi : Y_k \rightarrow Y_k/l'(f_{\mathcal{E}}(E))Y_k$.

Definition 2.4. $f = (f_{\mathcal{V}}, (f_{l,x})_{x \in \mathcal{V}}) : \mathcal{G} \rightarrow \mathcal{G}'$ is a k -isomorphism of moment graphs if $f : \mathcal{G} \rightarrow \mathcal{G}'$ is a k -homomorphism and the following two conditions hold:

- (ISO1) $f_{\mathcal{V}}$ is an isomorphism of posets;
- (ISO2) for all $u \rightarrow w \in \mathcal{E}'$, there exists exactly one $x \rightarrow y \in \mathcal{E}$ such that $f_{\mathcal{V}}(x) = u$ and $f_{\mathcal{V}}(y) = w$.

If f is a k -isomorphism from the moment graph \mathcal{G} to itself, we say that it is a k -automorphism of \mathcal{G} .

2.2. Bruhat graphs

Here we describe a class of moment graphs, that is for our purposes the most important.

We start by recalling some notation from [19]. Let \mathfrak{g} be a symmetrizable Kac–Moody algebra, that is the Lie algebra $\mathfrak{g}(A)$ associated to a symmetrizable generalized Cartan matrix A , and \mathfrak{h} its Cartan subalgebra. Let $\Pi = \{\alpha_i\}_{i=1,\dots,n} \subset \mathfrak{h}^*$, resp. $\Pi^\vee = \{\alpha_i^\vee\}_{i=1,\dots,n} \subset \mathfrak{h}$, be the set of simple roots, resp. coroots; let Δ , resp. Δ_+ , resp. Δ_+^{re} be the root system, resp. the set of positive roots, resp. the set of positive real roots; and let $Q = \sum_{i=1}^n \mathbb{Z}\alpha_i$, resp. $Q^\vee = \sum_{i=1}^n \mathbb{Z}\alpha_i^\vee$, be the root lattice, resp. the coroot lattice. For any $\alpha \in \Delta$, we denote by $s_\alpha \in GL(\mathfrak{h}^*)$ the reflection, whose action on $v \in \mathfrak{h}^*$ is given by

$$s_\alpha(v) = v - \langle v, \alpha^\vee \rangle \alpha. \tag{1}$$

Let $\mathcal{W} = \mathcal{W}(A)$ be the Weyl group associated to A , that is the subgroup of $GL(\mathfrak{h}^*)$ generated by the set of simple reflections $\mathcal{S} = \{s_\alpha \mid \alpha \in \Pi\}$. Recall that $(\mathcal{W}, \mathcal{S})$ is a Coxeter system (cf. [19], Section 3.10).

However, \mathcal{W} can be seen also as subgroup of $GL(\mathfrak{h})$, by setting, for any $\lambda \in \mathfrak{h}$

$$s_\alpha(\lambda) := \lambda - \langle \alpha, \lambda \rangle \alpha^\vee. \tag{2}$$

We will denote by $\mathcal{T} \subset \mathcal{W}$ the set of reflections, that is

$$\mathcal{T} = \{s_\alpha \mid \alpha \in \Delta_+^{\text{re}}\} = \{wsw^{-1} \mid w \in \mathcal{W}, s \in \mathcal{S}\}. \tag{3}$$

Hereafter we will write α_t to denote the positive real root corresponding to the reflection $t \in \mathcal{T}$. Finally, denote by $\ell : \mathcal{W} \rightarrow \mathbb{Z}_{\geq 0}$ the length function and by \leq the Bruhat order on \mathcal{W} .

For any $J \subseteq \mathcal{S}$, denote by $\mathcal{W}_J := \langle J \rangle$ and by \mathcal{W}^J the set of minimal representatives of the equivalence classes of $\mathcal{W}/\mathcal{W}_J$.

Definition 2.5. Let \mathcal{W} , \mathcal{S} and J be as above. Then the Bruhat (moment) graph $\mathcal{G}^J = \mathcal{G}(\mathcal{W}^J) = (\mathcal{V}, \mathcal{E}, \leq, l)$ associated to \mathcal{W}^J is a moment graph on Q^\vee and it is given by

- (i) $\mathcal{V} = \mathcal{W}^J$;
- (ii) $\mathcal{E} = \{x \rightarrow y \mid x < y, \exists \alpha \in \Delta_+^{\text{re}}, \exists w \in \mathcal{W}_J \text{ such that } ywx^{-1} = s_\alpha\}$;
- (iii) $l(x \rightarrow s_\alpha x w^{-1}) := \alpha^\vee$.

Such a moment graph has an important geometric meaning. Indeed, there is a partial flag variety Y corresponding to \mathcal{W} and J as above (see [24]) and it carries an action of a torus T , whose Lie algebra is \mathfrak{h} , and a (T -invariant) stratification with certain good properties (see [3]). The Bruhat graph encodes the action of this torus, in particular, the vertices are the 0-dimensional orbits, while the edges represent the 1-dimensional orbits (cf. Section 2.1 of [15]). The partial order on the set of vertices is induced by the stratification coming from the decomposition $Y = \bigsqcup_{w \in \mathcal{W}^J} Y_w$, where, indeed, $\overline{Y}_w = \bigsqcup_{\substack{y \leq w \\ y \in \mathcal{W}^J}} Y_y$.

3. Category of k -sheaves on a moment graph

Consider a moment graph $\mathcal{G} = (\mathcal{V}, \mathcal{E}, \leq, l)$ on a lattice Y . Recall that for any local ring k with $2 \in k^\times$ we denoted by $Y_k := Y \otimes_{\mathbb{Z}} k$.

Let $S_k := \text{Sym}(Y_k)$ be the symmetric algebra of Y_k . We provide S_k with a \mathbb{Z} -grading such that $(S_k)_{\{2\}} = Y_k$. From now on every S_k -module will be finitely generated and \mathbb{Z} -graded and every morphism between S_k -modules will be of degree zero.

Definition 3.1. A k -sheaf \mathcal{F} on \mathcal{G} is given by $(\{\mathcal{F}^x\}, \{\mathcal{F}^E\}, \{\rho_{x,E}\})$, where:

- (i) for all $x \in \mathcal{V}$, \mathcal{F}^x is an S_k -module;
- (ii) for all $E \in \mathcal{E}$, \mathcal{F}^E is an S_k -module such that $l(E) \cdot \mathcal{F}^E = \{0\}$;
- (iii) for $x \in \mathcal{V}$ and $E \in \mathcal{E}$ such that x is in the boundary of the edge E , the map $\rho_{x,E} : \mathcal{F}^x \rightarrow \mathcal{F}^E$ is a homomorphism of S_k -modules.

Definition 3.2. A homomorphism $\varphi : \mathcal{F} \rightarrow \mathcal{F}'$ between k -sheaves on the moment graph \mathcal{G} is given by the following data:

- (i) for all $x \in \mathcal{V}$, $\varphi^x : \mathcal{F}^x \rightarrow \mathcal{F}'^x$ is a homomorphism of S_k -modules;
- (ii) for all $E \in \mathcal{E}$, $\varphi^E : \mathcal{F}^E \rightarrow \mathcal{F}'^E$ is a homomorphism of S_k -modules such that for any $x \in \mathcal{V}$ on the border of $E \in \mathcal{E}$ the following diagram commutes

$$\begin{array}{ccc}
 \mathcal{F}^x & \xrightarrow{\varphi^x} & \mathcal{F}'^x \\
 \rho_{x,E} \downarrow & & \downarrow \rho'_{x,E} \\
 \mathcal{F}^E & \xrightarrow{\varphi^E} & \mathcal{F}'^E
 \end{array}$$

We denote by $\mathfrak{Sh}_{\mathcal{G}}^k$ the category of k -sheaves on \mathcal{G} having as objects the k -sheaves on \mathcal{G} and as morphisms the homomorphisms between them.

3.1. Pullback sheaves

Let $\mathcal{G} = (\mathcal{V}, \mathcal{E}, \leq, l)$ and $\mathcal{G}' = (\mathcal{V}', \mathcal{E}', \leq', l')$ be two moment graphs on Y and fix $f : \mathcal{G} \rightarrow \mathcal{G}'$ a k -homomorphism of moment graphs on Y (cf. Section 2.1).

Definition 3.3. Let $\mathcal{F} \in \text{Ob}(\mathfrak{Sh}_{\mathcal{G}}^k)$, then $f^*\mathcal{F} \in \text{Ob}(\mathfrak{Sh}_{\mathcal{G}'}^k)$ is defined as follows:

(PULL1) for all $x \in \mathcal{V}$, $(f^*\mathcal{F})^x := \mathcal{F}^{f_{\mathcal{V}}(x)}$ and $s \in S_k$ acts on it via $f_{l,x}(s)$;

(PULL2) for all $E : x \rightarrow y \in \mathcal{E}$

$$(f^*\mathcal{F})^E = \begin{cases} \mathcal{F}^{f_{\mathcal{V}}(x)}/l(E)\mathcal{F}^{f_{\mathcal{V}}(x)} & \text{if } f_{\mathcal{V}}(x) = f_{\mathcal{V}}(y) \\ \mathcal{F}^{f_{\mathcal{E}}(E)} & \text{otherwise} \end{cases}$$

and of $s \in S_k$ acts on $(f^*\mathcal{F})^E$ via $f_{l,x}(s)$;

(PULL3) for all $x \in \mathcal{V}$ and $E \in \mathcal{E}'$, such that $E : x \rightarrow y$,

$$(f^*\rho)_{x,E} = \begin{cases} \text{canonical quotient map} & \text{if } f_{\mathcal{V}}(x) = f_{\mathcal{V}}(y) \\ \rho_{f_{\mathcal{V}}(x), f_{\mathcal{E}}(E)} & \text{otherwise.} \end{cases}$$

Remark 3.1. For all $E \in \mathcal{E}$, the action of S_k on $(f^*\mathcal{F})^E$ in (PULL2) is well-defined thanks to conditions (MORPH2a) and (MORPH2b).

We say that $f^*\mathcal{F}$ is the *pullback* of \mathcal{F} . In what follows, the notion of pullback sheaf will allow us to compare k -sheaves on different moment graphs.

3.2. Sections of sheaves

For each $\mathcal{I} \subset \mathcal{V}$ we can consider the set of local sections of a k -sheaf $\mathcal{F} \in \text{Ob}(\mathfrak{Sh}_{\mathcal{G}}^k)$ over \mathcal{I} :

$$\Gamma(\mathcal{I}, \mathcal{F}) := \left\{ (m_x) \in \prod_{x \in \mathcal{I}} \mathcal{F}^x \mid \begin{array}{l} \rho_{x,E}(m_x) = \rho_{y,E}(m_y) \\ \forall E : x \rightarrow y \in \mathcal{E}, x, y \in \mathcal{I} \end{array} \right\}.$$

We denote by $\Gamma(\mathcal{F}) = \Gamma(\mathcal{V}, \mathcal{F})$ the set of global sections of the k -sheaf \mathcal{F} .

We call k -structure algebra of the moment graph \mathcal{G} the set

$$\mathcal{Z}_k = \mathcal{Z}_k(\mathcal{G}) := \left\{ (z_x) \in \prod_{x \in \mathcal{V}} S_k \mid \begin{array}{l} z_x - z_y \in l(E)S_k \\ \forall E : x \rightarrow y \in \mathcal{E} \end{array} \right\}.$$

It is easy to check that for any $\mathcal{F} \in \text{Ob}(\mathfrak{Sh}_{\mathcal{G}}^k)$ the k -structure algebra \mathcal{Z}_k acts on $\Gamma(\mathcal{F})$ via component-wise multiplication.

3.3. Flabby sheaves

We use the order on the set of vertices of a moment graph \mathcal{G} to define a topology on it: the *Alexandrov topology*. We say that \mathcal{I} is open if for any $x \in \mathcal{I}$ and any $y \in \mathcal{V}$ such that $x \triangleleft y$ then $y \in \mathcal{I}$ as well.

A classical question in sheaf theory is to ask whether a sheaf is flabby or not, that is whether any local section over an open set extends to a global one or not.

Let $\mathcal{F} \in \text{Ob}(\mathfrak{Sh}_{\mathcal{G}}^k)$. We fix a vertex $x \in \mathcal{V}$ and we denote

$$\mathcal{E}_{\delta x} := \{E \in \mathcal{E} \mid E : x \rightarrow y\},$$

$$\mathcal{V}_{\delta x} := \{y \in \mathcal{V} \mid \exists E \in \mathcal{E}_{\delta x} \text{ such that } E : x \rightarrow y\}.$$

Now we define $\mathcal{F}^{\delta x}$ as the image of $\Gamma(\{\triangleright x\}, \mathcal{F}) := \Gamma(\{y \in \mathcal{V} \mid y \triangleright x\}, \mathcal{F})$ under the composition of the following functions:

$$u_x : \Gamma(\{\triangleright x\}, \mathcal{F}) \longrightarrow \bigoplus_{y \triangleright x} \mathcal{F}^y \longrightarrow \bigoplus_{y \in \mathcal{V}_{\delta x}} \mathcal{F}^y \xrightarrow{\oplus \rho_{y,E}} \bigoplus_{E \in \mathcal{E}_{\delta x}} \mathcal{F}^E.$$

Denote

$$d_x = \bigoplus_{E \in \mathcal{E}_{\delta x}} \rho_{x,E} : \mathcal{F}^x \longrightarrow \bigoplus_{E \in \mathcal{E}_{\delta x}} \mathcal{F}^E.$$

Observe that $m \in \Gamma(\{\triangleright x\}, \mathcal{F})$ can be extended, via m_x , to a section $\tilde{m} = (m, m_x) \in \Gamma(\{\triangleright x\}, \mathcal{F})$ if and only if $d_x(m_x) = u_x(m)$. This fact motivates the following result, due to Fiebig, that gives a characterization of the flabby objects in $\mathfrak{Sh}_{\mathcal{G}}^k$.

Proposition 3.1. (See [9].) *Let $\mathcal{F} \in \text{Ob}(\mathfrak{Sh}_{\mathcal{G}}^k)$. Then the following are equivalent:*

- (i) \mathcal{F} is flabby with respect to the Alexandrov topology, i.e. for any open $\mathcal{I} \subseteq \mathcal{V}$ the restriction map $\Gamma(\mathcal{F}) \rightarrow \Gamma(\mathcal{I}, \mathcal{F})$ is surjective.
- (ii) For any vertex $x \in \mathcal{V}$ the restriction map $\Gamma(\{\triangleright x\}, \mathcal{F}) \rightarrow \Gamma(\{\triangleright x\}, \mathcal{F})$ is surjective.
- (iii) For any vertex $x \in \mathcal{V}$ the map $d_x : \mathcal{F}^x \rightarrow \bigoplus_{E \in \mathcal{E}_{\delta x}} \mathcal{F}^E$ contains $\mathcal{F}^{\delta x}$ in its image.

4. Braden–MacPherson sheaves

In this section we introduce the most important object of our paper, namely the canonical sheaf. It was first defined by Braden and MacPherson—only in characteristic zero—in order to compute certain intersection cohomology complexes. Despite this, their algorithm makes sense in any characteristic and Fiebig and Williamson proved in [15] that it computes the multiplicities of parity sheaves (see [18]) in positive characteristic if (\mathcal{G}, k) is a GKM-pair. The following theorem allows us to consider this sheaf.

Theorem 4.1. (See [3], char $k = 0$; [9].) *Let \mathcal{G} be a finite k -moment graph over Y with highest vertex w . There exists exactly one (up to isomorphism) indecomposable k -sheaf \mathcal{B}_w on \mathcal{G} with the following properties:*

- (i) $\mathcal{B}_w^w \cong S_k$;
- (ii) If $x, y \in \mathcal{V}$, $E : x \rightarrow y \in \mathcal{E}$, then the map $\rho_{y,E} : \mathcal{B}_w^y \rightarrow \mathcal{B}_w^E$ is surjective with kernel $l(E)\mathcal{B}_w^y$;
- (iii) If $x, y \in \mathcal{V}$, $E : x \rightarrow y \in E$, then $\rho_{\delta x} := \bigoplus_{E \in \mathcal{E}_{\delta x}} \rho_{x,E} : \mathcal{B}_w^x \rightarrow \mathcal{B}_w^{\delta x}$ is a projective cover in the category of graded S_k -modules.

We call \mathcal{B}_w the Braden–MacPherson sheaf or the canonical sheaf. We will also refer to it as the BMP-sheaf.

Remark 4.1. By Theorem 4.1 and Proposition 3.1, the canonical sheaf is flabby for the Alexandrov topology. This property will be crucial in what follows.

4.1. Graded rank of the stalks of a BMP-sheaf

For $j \in \mathbb{Z}$ and M a graded S -module we denote by $M\{j\}$ the graded S -module obtained from M by shifting the grading by j , i.e. $M\{j\}_{\{i\}} = M_{\{j+i\}}$. If $M = \bigoplus_{i=1}^n S_k\{j_i\}$, then its graded rank is $\text{rk } M = \sum_{i=1}^n q^{-\frac{j_i}{2}} \in \mathbb{Z}_{\geq 0}[q^{\frac{1}{2}}, q^{-\frac{1}{2}}]$.

Let \mathcal{G}^J be the Bruhat graph we defined in Section 2.2. Thus for any $w \in \mathcal{W}^J$ we can consider the subgraph $\mathcal{G}_w^J := \mathcal{G}_{\{\leq w\}}^J$. It is a finite k -moment graph (for any k) with highest vertex w , hence we may build the corresponding Braden–MacPherson sheaf $\mathcal{B}_w^J \in \text{Ob}(\mathfrak{Sh}_{\mathcal{G}_w^J}^k)$ and we have:

Question 4.1. Under which assumptions on the characteristic of the base field, do we have $\underline{\text{rk}}(\mathcal{B}_w^J)^y = P_{y,w}^{J,-1}$ for $y \leq w$ and y, w varying in some relevant subset of \mathcal{W}^J ?

$P_{y,w}^{J,-1}$ stand for the parabolic Kazhdan–Lusztig polynomials (corresponding to the parameter $u = -1$) introduced by Deodhar in [7].

If $k = \mathbb{Q}$, then $\underline{\text{rk}}(\mathcal{B}_w^J)^y = P_{y,w}^{J,-1}$ for any pair $y, w \in \mathcal{W}^J$, with $y \leq w$, from [20,21,3]. Moreover, Fiebig proved that in this case the equality is equivalent to a character formula conjectured by Kazhdan and Lusztig in [20] (see [9]). Hence for a fixed pair of elements, from the characteristic zero case we get that the equality holds for p large enough. Observe that the bound depends on the pair and there is no global bound in the infinite case.

Now let \mathcal{W} be an affine Weyl group, h be its Coxeter number, k be a field of characteristic $p \geq h$ and y, w be restricted elements (cf. [13]). A positive answer to Question 4.1 would imply a conjecture by Lusztig (cf. [11,13]). Fiebig was able to prove that the stalks $(\mathcal{B}_w^J)^y$ have the expected graded rank for p bigger than an explicit—but huge!—bound depending on \mathcal{W} (cf. [14]). Motivated by the fact that in the affine case the GKM-condition (see Definition 2.2(ii)) for the Bruhat graph of restricted elements is precisely $p \geq h$ (see [12], Lemma 4.3), Fiebig suggested the answer to Question 4.1 to be yes as soon as the GKM-condition were satisfied (cf. [11], Conjecture 4.4).

Actually, very recently this conjecture has been proven to be false for $\mathcal{W} = S_{4p}$. Indeed, Polo (private communication, 7 May 2012) produced a family of elements w_n in S_{4n} (for each integer $n \geq 2$) such that there is n -torsion in some costalk of the intersection cohomology of the Schubert variety corresponding to w_n . The fact that this provides us with a family of counterexamples to Fiebig’s conjecture is not immediate at all. We have to notice first that the Bruhat graph for sl_r and k constitute a GKM-pair for any r and any field k of characteristic $p > 2$ and then to translate Question 4.1 in terms of intersection cohomology complexes and parity sheaves (cf. [15], Theorem 9.2).

Finally, let us consider an affine Weyl group \mathcal{W} , whose Coxeter number is h and a field k of characteristic $p > h$, but let us make y, w vary in the finite Weyl group $\mathcal{W}^f < \mathcal{W}$. In this case a positive answer to Question 4.1 would imply the Lusztig’s conjecture around the Steinberg weight, which was presented by Soergel in the 90s as “toy model” for the original Lusztig’s conjecture (cf. [26]).

In view of Polo’s result, the bound $p \geq h$ seems to be the right one for expecting Question 4.1 to have a positive answer for y, w restricted, resp. for any pair y, w , if \mathcal{W} is affine, resp. finite. Moreover in this case, this problem would still be related to Lusztig’s conjecture, resp. Lusztig’s conjecture on the Steinberg weight, as discussed above.

Anyway, Question 4.1 proposes us an explicit formula connecting canonical sheaves and parabolic Kazhdan–Lusztig polynomials and motivates our work. We will indeed interpret in terms of stalks of BMP-sheaves some well-known identities concerning these polynomials.

5. Pullback of BMP-sheaves

The following lemma tells us that the pullback functor f^* preserves canonical sheaves if f is a k -isomorphism.

Lemma 5.1. Let \mathcal{G} and \mathcal{G}' be two k -moment graphs on Y , both with a unique maximal vertex, w resp. w' , and let $f : \mathcal{G} \rightarrow \mathcal{G}'$ be a k -isomorphism. If \mathcal{B}_w and $\mathcal{B}'_{w'}$ are the corresponding canonical sheaves, then $\mathcal{B}_w \cong f^* \mathcal{B}'_{w'}$ as k -sheaves on \mathcal{G} .

Proof. Let $\mathcal{G} = (\mathcal{V}, \mathcal{E}, \leq, l)$, $\mathcal{G}' = (\mathcal{V}', \mathcal{E}', \leq', l')$ and $f = (f_{\mathcal{V}}, (f_{l,x}))$.

Notice that $\mathcal{I} \subseteq \mathcal{V}$ is an open subset if and only if $\mathcal{I}' := f_{\mathcal{V}}(\mathcal{I}) \subseteq \mathcal{V}'$ is an open subset. We prove that $\mathcal{B}_w|_{\mathcal{I}} \cong f^* \mathcal{B}'_{w'}|_{\mathcal{I}'}$ by induction on $|\mathcal{I}| = |\mathcal{I}'|$, for \mathcal{I} open.

If $|\mathcal{I}| = |\mathcal{I}'| = 1$, we have $\mathcal{I} = \{w\}$ and $\mathcal{I}' = \{w'\}$. In this case $\mathcal{B}_w^w = S_k$, $\mathcal{B}'_{w'}^{w'} = S_k$ and the isomorphism $\varphi^w : \mathcal{B}_w^w \rightarrow \mathcal{B}'_{w'}^{w'}$ is just given by the twisting of the S_k -action, coming from the automorphism of S_k induced by the automorphism $f_{l,w}$ of Y_k .

Now let $|\mathcal{I}| = |\mathcal{I}'| = n > 1$ and $y \in \mathcal{I}$ be a minimal element. Obviously, $y' := f_{\mathcal{V}}(y)$ is also a minimal element for \mathcal{I}' . Moreover, for any $E \in \mathcal{E}$ we set $E' := f_{\mathcal{E}}(E)$.

First of all, observe that $z \in \mathcal{V}_{\delta y}$ if and only if $z' := f_{\mathcal{V}}(z) \in \mathcal{V}'_{\delta y'}$. By the inductive hypothesis, for all $x \triangleright y$ there exists an isomorphism $\varphi^x : \mathcal{B}_w^x \xrightarrow{\sim} \mathcal{B}'_{w'}^{x'}$ such that $\varphi^x(s \cdot m) = f_{l,x}(s) \cdot \varphi^x(m)$, for $s \in S_k$ and $m \in \mathcal{B}_w^x$. Moreover, if $E \notin \mathcal{E}_{\delta y}$ and x is on the border of E with $x \triangleright y$, by the inductive hypothesis we have an isomorphism $\varphi^E : \mathcal{B}_w^E \xrightarrow{\sim} \mathcal{B}'_{w'}^{E'}$ such that $\varphi^E(s \cdot n) = f_{l,x}(s) \cdot \varphi^E(n)$, for $s \in S_k$ and $n \in \mathcal{B}_w^E$ and such that the following diagram commutes

$$\begin{array}{ccc} \mathcal{B}_w^x & \xrightarrow{\varphi^x} & \mathcal{B}'_{w'}^{x'} \\ \rho_{x,E} \downarrow & & \downarrow \rho'_{x',E'} \\ \mathcal{B}_w^E & \xrightarrow{\varphi^E} & \mathcal{B}'_{w'}^{E'} \end{array}$$

Now, if $E : y \longrightarrow x$ and $E' : y' \longrightarrow x'$, then

$$\mathcal{B}_w^E \cong \mathcal{B}_w^x / l(E) \mathcal{B}_w^x \quad \text{and} \quad \mathcal{B}'_{w'}^{E'} \cong \mathcal{B}'_{w'}^{x'} / l'(E') \mathcal{B}'_{w'}^{x'}.$$

By assumption, $f_{l,x}(l(E)) = h \cdot l'(E')$ for some invertible element $h \in k^\times$ and $\varphi^x(l(E) \mathcal{B}_w^x) = f_{l,x}(l(E)) \mathcal{B}'_{w'}^{x'} = l'(E') \mathcal{B}'_{w'}^{x'}$. Thus the quotients are also isomorphic and so there exists $\varphi^E : \mathcal{B}_w^E \xrightarrow{\sim} \mathcal{B}'_{w'}^{E'}$ such that the following diagram commutes:

$$\begin{array}{ccc} \mathcal{B}_w^x & \xrightarrow{\varphi^x} & \mathcal{B}'_{w'}^{x'} \\ \rho_{x,E} \downarrow & & \downarrow \rho'_{x',E'} \\ \mathcal{B}_w^E & \xrightarrow[\varphi^E]{\sim} & \mathcal{B}'_{w'}^{E'} \end{array}$$

Now we have to construct $\mathcal{B}_w^{\delta y}$ and $\mathcal{B}'_{w'}^{\delta y'}$. Observe that $(\varphi^x)_{x \triangleright y}$ induces an isomorphism of S_k -modules between the sets of sections $\Gamma(\{\triangleright y\}, \mathcal{B}_w) \cong \Gamma(\{\triangleright y'\}, \mathcal{B}'_{w'})$ and, from what we have observed above, the following diagram commutes:

$$\begin{array}{ccccccc} & & & & u_y & & \\ & & & & \curvearrowright & & \\ \Gamma(\{\triangleright y\}, \mathcal{B}_w) & \longrightarrow & \bigoplus_{x \triangleright y} \mathcal{B}_w^x & \longrightarrow & \bigoplus_{x \in \mathcal{V}_{\delta y}} \mathcal{B}_w^x & \xrightarrow{\oplus \rho_{x,E}} & \bigoplus_{E \in \mathcal{E}_{\delta y}} \mathcal{B}_w^E \\ \downarrow \oplus_{x \triangleright y} \varphi^x & & \downarrow \oplus_{x \triangleright y} \varphi^x & & \downarrow \oplus_{x \in \mathcal{V}_{\delta y}} \varphi^x & & \downarrow \oplus_{E \in \mathcal{E}_{\delta y}} \varphi^E \\ \Gamma(\{\triangleright y'\}, \mathcal{B}'_{w'}) & \longrightarrow & \bigoplus_{x' \triangleright y'} \mathcal{B}'_{w'}^{x'} & \longrightarrow & \bigoplus_{x' \in \mathcal{V}_{\delta y'}} \mathcal{B}'_{w'}^{x'} & \xrightarrow{\oplus \rho'_{x',E'}} & \bigoplus_{E \in \mathcal{E}_{\delta y'}} \mathcal{B}'_{w'}^{E'} \\ & & & & \curvearrowleft & & \\ & & & & u'_{y'} & & \end{array}$$

It follows that there exists an isomorphism of S_k -modules $\mathcal{B}_w^{\delta y} \cong \mathcal{B}'_{w'}^{\delta y'}$ and by the unicity of the projective cover we obtain $\mathcal{B}_w^y \cong \mathcal{B}'_{w'}^{y'}$. This proves the lemma. \square

Remark 5.1. Let $y, x, z, w \in \mathcal{W}$ be such that $y \leq w$ and $x \leq z$. If one could show that any isomorphism of posets $[y, w] \cong [x, z]$ induces a k -isomorphism of moment graphs $f : \mathcal{G}_{|[y,w]} \rightarrow \mathcal{G}_{|[x,z]}$ (at least for $k = \mathbb{Q}$), then, by Lemma 5.1, the Lusztig–Dyer *Combinatorial Invariance Conjecture* (stated in [8]) would follow. See [5] for partial results on this conjecture.

5.1. Two KL-properties of the canonical sheaf

Here we apply Lemma 5.1 in order to lift some equalities concerning KL-polynomials to the moment graph setting.

From now on we denote by $\mathcal{G} = (\mathcal{V}, \mathcal{E}, l, \leq)$ the Bruhat graph corresponding to a Weyl group \mathcal{W} and $J = \emptyset$. As in Section 2.2 we denote by \mathcal{S} and \mathcal{T} the set of simple reflections and all reflections, respectively, of \mathcal{W} . Recall that \mathcal{G} is a moment graph on the coroot lattice Q^\vee and that there is a linear \mathcal{W} -action Q^\vee .

5.1.1. Inverses

Kazhdan and Lusztig gave an inductive formula to calculate the KL-polynomials ((2.2.c) of [20]). From such a formula it follows easily by induction (cf. Example 12, Chapter 5 of [4]) that for any pair $y, w \in \mathcal{W}$ such that $y \leq w$ one has

$$P_{y,w} = P_{y^{-1},w^{-1}}. \tag{4}$$

We translate this equality to a k -isomorphism of stalks of canonical sheaves.

Lemma 5.2. *Let \mathcal{W} be a Weyl group. The anti-involution on \mathcal{W} defined by the mapping $x \mapsto x^{-1}$ induces a k -automorphism of the Bruhat graph \mathcal{G} for any k .*

Proof. The map $f_{\mathcal{V}} : \mathcal{V} \rightarrow \mathcal{V}$ defined by $x \mapsto x^{-1}$ is obviously a bijection. Moreover, for each $x, y \in \mathcal{W}$, $x \leq y$ if and only if $x^{-1} \leq y^{-1}$. So $f_{\mathcal{V}} : \mathcal{V} \rightarrow \mathcal{V}$ is an isomorphism of posets.

Observe that there exists a reflection $t \in \mathcal{T}$ such that $y = tx$ if and only if $y^{-1} = rx^{-1}$, where $r = x^{-1}tx \in \mathcal{T}$. So $x \xrightarrow{t} y \in \mathcal{E}$ if and only if $x^{-1} \xrightarrow{r} y^{-1} \in \mathcal{E}$.

Thus, for every $x \in \mathcal{W}$ and any $v \in Y_k$, we set $f_{l,x}(v) := x^{-1}(v)$ and observe that if $E : x \xrightarrow{t} y = tx$, we have

- (a) $f_{l,x}(l(x \xrightarrow{t} tx)) = x^{-1}(\alpha_t^\vee) = x^{-1}(\alpha_t)^\vee = \pm l(x^{-1} \xrightarrow{r} y^{-1})$, where $\pm x^{-1}(\alpha_t) \in \Delta_+^{re}$, because $x^{-1}(\alpha_t) = \pm \alpha_{x^{-1}tx}$.
- (b) $f_{l,y}(v) = y^{-1}(v) = x^{-1}(tv) = x^{-1}(v) - \langle \alpha_t, v \rangle x^{-1}(\alpha_t^\vee) \equiv x^{-1}(v) = f_{l,x}(v) \pmod{x^{-1}(\alpha_t^\vee)}$.

This proves that we have a k -automorphism of the moment graph \mathcal{G} for any k . \square

From this we obtain the following corollary.

Corollary 5.1. *Let $w \in \mathcal{W}$. Then there exists an isomorphism $g : \mathcal{G}_w \rightarrow \mathcal{G}_{w^{-1}}$ of k -moment graphs on Q^\vee and $\mathcal{B}_w \cong g^* \mathcal{B}_{w^{-1}}$ as k -sheaves on \mathcal{G}_w for any k .*

Proof. By Lemma 5.2, $f_{\mathcal{V}} : x \mapsto x^{-1}$ induces a k -isomorphism between the two complete subgraphs \mathcal{G}_w and $\mathcal{G}_{w^{-1}}$. We may then set $g := f_{|\mathcal{G}_w}$ and apply Lemma 5.1; the statement follows. \square

5.1.2. Multiplying by a simple reflection. Part I

Let $y, w \in \mathcal{W}$ and $s \in \mathcal{S}$ such that $y \leq w$, $ws < w$ and $y \not\leq ws$. Under those hypotheses Kazhdan and Lusztig observed (proof of Theorem 4.2 of [20]) that

$$P_{y,w} = P_{ys,ws}. \tag{5}$$

In order to interpret (5) in our moment graph setting we need a standard combinatorial result (that actually holds for any Coxeter group):

Lemma 5.3. (See [17], Lemma 7.4.) *Let $s \in \mathcal{S}$ and $v, u \in \mathcal{W}$ be such that $vs < v$ and $u < v$.*

- (i) *If $us < u$, then $us < vs$.*
- (ii) *If $us > u$, then $us \leq v$ and $u \leq vs$.*

Thus, in both cases, $us \leq v$.

We are now able to define for any k a k -isomorphism of Bruhat (sub)graphs:

Lemma 5.4. *Let $y, w \in \mathcal{W}$ and $s \in \mathcal{S}$ such that $y \leq w$, $ws < w$ and $y \not\leq ws$, then for any k there is a k -isomorphism of moment graphs $\mathcal{G}_{|[y,w]} \xrightarrow{\sim} \mathcal{G}_{|[ys,ws]}$.*

Proof. We show that $f_{\mathcal{V}} : [y, w] \rightarrow [ys, ws]$, $x \mapsto xs$ is a bijection of posets inducing the identity map on the labels.

We verify that if $x \in [y, w]$ then $xs \in [ys, ws]$. We see that $xs < x$; indeed, if it were not the case, then by Lemma 5.3(ii) we would have $x \leq ws$, but this would imply $y \leq ws$. In particular, this holds for y , that is $ys < y$. Now, by Lemma 5.3(i),

$$xs < x, \quad ws < w \quad \Rightarrow \quad xs \leq ws,$$

$$ys < y, \quad xs < x \quad \Rightarrow \quad ys \leq xs.$$

We now show that if $z \in [ys, ws]$ then $zs \in [y, w]$. Observe that $zs > z$; indeed, $ys < z$, $y = (ys)s > ys$ and if $zs < z$, then by Lemma 5.3(ii), with $u = ys$ and $v = z$, we would get $y = (ys)s \leq z \leq ws$. Moreover, $z \leq ws < w$ and, by Lemma 5.3(ii),

$$zs > z, \quad ws < w \quad \Rightarrow \quad zs \leq w,$$

$$y = (ys)s > ys, \quad z = (zs)s < zs \quad \Rightarrow \quad y \leq zs.$$

This completes the proof that $f_{\mathcal{V}}$ maps $[y, w]$ to $[ys, ws]$.

Let $x, z \in [y, w]$, then $x \leq z$ if and only if $xs \leq zs$. Indeed, we have already proved that $xs < x$ and $zs < z$ so, by Lemma 5.3(i), with $u = x$ and $v = z$, we have $xs \leq zs$. On the other hand, $x = (xs)s > xs$ and it follows from Lemma 5.3(ii) with $u = xs$ and $v = z$ that $x = (xs)s \leq z$.

Finally from what we proved above, for each $t \in \mathcal{T}$ we have that $x, tx \in [y, w]$ if and only if $xs, txs \in [ys, ws]$. This means that we have a bijection between sets of edges such that $f_{\mathcal{E}}(x \xrightarrow{\gamma} tx) = xs \xrightarrow{\gamma} txs$.

Therefore $f = (f_{\mathcal{V}}, (Id_{\mathcal{V}})_{x \in \mathcal{V}})$ is a k -isomorphism of moment graphs on Q^{\vee} for any k . \square

So we have:

Corollary 5.2. *Consider $y, w \in \mathcal{W}$ such that $y \leq w$ and $ws < w$, $y \not\leq ws$ for some $s \in \mathcal{S}$. Let f be as in Lemma 5.4, then $\mathcal{B}_w \cong f^* \mathcal{B}_{ws}$ as k -sheaves on $\mathcal{G}_{|[y,w]}$ for any k .*

Proof. The statement follows by combining Lemma 5.4 and Lemma 5.1. \square

We recollect the results of this section:

Theorem 5.1. *Let $y, w \in \mathcal{W}$ be such that $y \leq w$, then*

(i) $\mathcal{B}_w^y \cong \mathcal{B}_{w^{-1}}^{y^{-1}}$.

Let $s \in \mathcal{S}$ be such that $ws < w$, then

(ii) if $y \not\leq ws$, $\mathcal{B}_w^y \cong \mathcal{B}_{ws}^{ys}$.

All isomorphisms are isomorphisms of S_k -modules, for any k .

Proof. (i) This follows from Corollary 5.1, since two k -sheaves are isomorphic only if their stalks are pairwise isomorphic.

(ii) As before, the isomorphism descends from the k -isomorphism of k -sheaves we obtained in Corollary 5.2. \square

6. Invariants

Clearly not all equalities concerning Kazhdan–Lusztig polynomials come from k -isomorphisms of the underlying Bruhat graphs. In this section we develop another technique and, as in the previous section, we apply it in order to categorify two well-known properties of these polynomials.

6.1. Multiplying by a simple reflection. Part II

Another property that Kazhdan and Lusztig in [20] (2.3.g) proved is that if $y, w \in \mathcal{W}$ and $s \in \mathcal{S}$ are such that $y \leq w$ and $ws < w$, then

$$P_{y,w} = P_{ys,w}. \tag{6}$$

It is clear that in this case there is no hope of finding any k -isomorphism of moment graphs, since the two Bruhat intervals $[y, w]$ and $[ys, w]$ obviously have different cardinality.

The goal of this section is to prove the following theorem.

Theorem 6.1. For any pair $y, w \in \mathcal{W}$ and for any $s \in \mathcal{S}$ such that $ws < w$ and $ys, y \leq w$, there exist

- an isomorphism of S_k -modules $\varphi^y : \mathcal{B}_w^y \rightarrow \mathcal{B}_w^{ys}$,
- a family of isomorphisms of S_k -modules $\varphi^E : \mathcal{B}_w^E \rightarrow \mathcal{B}_w^{Es}$, where $E : y \rightarrow x \in \mathcal{E}$ and $Es : ys \rightarrow xs \in \mathcal{E}$

such that the following diagram commutes

$$\begin{array}{ccc}
 \mathcal{B}_w^y & \xrightarrow{\varphi^y} & \mathcal{B}_w^{ys} \\
 \rho_{y,E} \downarrow & & \downarrow \rho_{ys,Es} \\
 \mathcal{B}_w^E & \xrightarrow{\varphi^E} & \mathcal{B}_w^{Es}
 \end{array} \tag{7}$$

and such that $\varphi^{ys} = (\varphi^y)^{-1}$.

6.2. Two preliminary lemmata

In order to prove our claim, we need two combinatorial lemmata.

Recall that

$$\mathcal{T} = \{s_\alpha \mid \alpha \in \Delta_+^{re}\} = \{wsw^{-1} \mid w \in \mathcal{W}, s \in \mathcal{S}\}$$

and, for all $x, y \in \mathcal{W}$, denote

$$G_L(x, y) := \{t \in \mathcal{T} \mid tx \in (x, y)\}.$$

Lemma 6.1. *Let $w, y \in \mathcal{W}$ and $s \in \mathcal{S}$ be such that $y \leq w$, $ws < w$ and $ys < y$, then*

$$G_L(ys, w) = G_L(y, w) \cup \{ysy^{-1}\}.$$

Proof. We show that for all $t \in G_L(y, w)$ we have $ys < tys \leq w$ as well, i.e. $t \in G_L(ys, w)$. Indeed, if $tys > ty$, then $ys < y < ty < tys$ and, by Lemma 5.3(ii) with $u = ty$ and $v = w$, $tys \leq w$. Otherwise, $tys < ty \leq w$, $y < ty$, $ys < y$ and, by Lemma 5.3(i) with $u = y$ and $v = ty$, we obtain $ys < tys$.

Clearly, $ysy^{-1} \in G_L(ys, w)$ and this completes the proof that the set on the right-hand side is a subset of the one on the left.

Now we verify that if $t \in \mathcal{T}$, $tys \in [ys, w]$ and $ty \notin [y, w]$, then $t = ysy^{-1}$. Indeed, by Lemma 5.3 with $u = tys$ and $v = w$, $tys \leq w$ and, if $ty \notin [y, w]$, then $ty < y$. Moreover, $ys < y$ and so, by Lemma 5.3(ii) with $u = ty$ and $v = y$, $tys \leq y$. So $ys < tys \leq y$ and we know that $[ys, y] = \{ys, y\}$. Thus $tys = y$, that is, $t = ysy^{-1}$. \square

Lemma 6.2. *Let $w, y \in \mathcal{W}$ and $s \in \mathcal{S}$ be such that $y \leq w$, $ys < y$ and $ws < w$, then the set $[ys, w] \setminus \{ys, y\}$ is stabilized by the mapping $x \mapsto xs$.*

Proof. Notice that $ys < y \leq w$, so it makes sense to write $[ys, w]$. Let $\mathcal{I} := [ys, w] \setminus \{ys, y\}$ and let $x \in \mathcal{I}$. If $xs > x$, then obviously $ys < xs$ and, by Lemma 5.3(ii) with $u = x$ and $v = w$, $xs \leq w$. On the other hand, if $xs < x$, then $xs < w$ and, by applying Lemma 5.3(ii) with $u = ys$ and $v = x$, $ys \leq xs$. Then, in both cases $xs \in [ys, w]$ and, since $xs \neq y$ and $xs \neq ys$, we get $x \in \mathcal{I}$.

Finally, if $x \in \mathcal{I}$, then $xs \neq y$. Indeed $xs = y$ if and only if $x = ysy^{-1} \notin \mathcal{I}$. \square

6.3. Proof of the main theorem

Let $\ell : \mathcal{W} \rightarrow \mathbb{Z}_{\geq 0}$ denote the length function on \mathcal{W} . We will prove Theorem 6.1 by induction on $n = \ell(w) - \ell(y)$.

If $n = 0$, then $y = w$ and there is nothing to prove. If $n > 0$ and $ys > y$, then $\ell(w) - \ell(ys) = n - 1$ and by induction we get the desired isomorphisms.

Now, we may suppose $n > 0$ and $ys < y$. Let $\mathcal{I} = [ys, w] \setminus \{y, ys\}$. From the inductive hypothesis, for any $x \in \mathcal{I}$ we get

- an isomorphism of S_k -modules $\varphi^x : \mathcal{B}_w^x \rightarrow \mathcal{B}_w^{xs}$,
- a family of isomorphisms of S_k -modules $\varphi^F : \mathcal{B}_w^F \rightarrow \mathcal{B}_w^{Fs}$, where $F : x \rightarrow z \in \mathcal{E}^{\delta y}$ and $Es : xs \rightarrow zs \in \mathcal{E}^{\delta ys}$

such that the following diagram commutes

$$\begin{array}{ccc} \mathcal{B}_w^x & \xrightarrow{\varphi^x} & \mathcal{B}_w^{xs} \\ \rho_{x,F} \downarrow & & \downarrow \rho_{xs,Fs} \\ \mathcal{B}_w^F & \xrightarrow{\varphi^F} & \mathcal{B}_w^{Fs} \end{array} \tag{8}$$

and such that $\varphi^{xs} = (\varphi^x)^{-1}$.

Observe that our claim will follow, once we prove that there is an isomorphism of S_k -modules $\varphi^y : \mathcal{B}_w^y \rightarrow \mathcal{B}_w^{ys}$ compatible with the restriction maps. Indeed, for $E : y \rightarrow x \in \mathcal{E}_{\delta y}$ there exists exactly

one $Es : ys \rightarrow xs \in \mathcal{E}_{\delta_{ys}}$, and φ^E would already have been given. If $E : ys \rightarrow y$, then we could set $\varphi^E = Id$. Finally, for $x \neq ys$, there exists an edge $E : x \rightarrow y \in \mathcal{E}$ if and only if there is $Es : xs \rightarrow ys \in \mathcal{E}$ (cf. Lemma 6.1) and in this case $\mathcal{B}_W^E \cong \mathcal{B}_W^y/l(E)\mathcal{B}_W^y \cong \mathcal{B}_W^{ys}/l(Es)\mathcal{B}_W^{ys}$, since $E = Es$.

We will get φ^y by defining a surjective map from \mathcal{B}_W^y to $\mathcal{B}_W^{\delta_{ys}}$. Since \mathcal{B}_W^{ys} is the projective cover of the S_k -module $\mathcal{B}_W^{\delta_{ys}}$, and, since $\text{rk}_{S_k} \mathcal{B}_W^y \leq \text{rk}_{S_k} \mathcal{B}_W^{ys}$ (cf. Lemma 3.12. of [10]), Theorem 6.1 will follow from the unicity of the projective cover.

6.3.1. Invariants

By Lemma 6.2, \mathcal{I} is invariant with respect to the right multiplication by s and we may define an automorphism σ_s of the set of global sections of the Braden–MacPherson sheaf as follows. Let $m = (m_x) \in \Gamma(\mathcal{I}, \mathcal{B}_W)$, then we set $\sigma_s(m) = (m'_x)$, where $m'_x := \varphi^{xs}(m_{xs})$. Since the φ^{xs} are, by definition, compatible with the restriction maps (see diagram (8)), $\sigma_s(m) \in \Gamma(\mathcal{I}, \mathcal{B}_W)$. Moreover, for any $x \in \mathcal{I}$, $\varphi^{xs} = (\varphi^x)^{-1}$ and so σ_s is an involution.

Let us denote by Γ^s the submodule of σ_s -invariant elements of $\Gamma(\mathcal{I}, \mathcal{B}_W)$, and by Γ^{-s} the elements $m \in \Gamma(\mathcal{I}, \mathcal{B}_W)$ such that $\sigma_s(m) = -m$.

Let us consider $c_s := (c_{s,x}) \in \bigoplus_{x \in \mathcal{W}} S_k$, where $c_{s,x} := x(\alpha_s^\vee)$; then $c_s \in \mathcal{Z}_k$ and so it acts on $\Gamma(\mathcal{I}, \mathcal{B}_W)$ via componentwise multiplication.

Lemma 6.3. *Let $(\mathcal{G}_{|\mathcal{I}}, k)$ be a GKM-pair, then we have $\Gamma(\mathcal{I}, \mathcal{B}_W) = \Gamma^s \oplus c_s \cdot \Gamma^s$.*

Proof. (We follow [13], Lemma 2.3.)

By definition, σ_s is an involution and 2 is an invertible element in k , so we get $\Gamma(\mathcal{I}, \mathcal{B}_W) = \Gamma^s \oplus \Gamma^{-s}$.

Let $m \in \Gamma^s$, then $\sigma_s(c_s \cdot m) = -(c_s \cdot m)$, i.e. $c_s \cdot \Gamma^s \subseteq \Gamma^{-s}$. Indeed, $s(\alpha_s^\vee) = -\alpha_s^\vee$ and so for any $x \in \mathcal{I}$ we have

$$(\sigma_s(c_s \cdot m))_x = \varphi^{xs}(xs(\alpha_s^\vee) \cdot m_{xs}) = -x(\alpha_s^\vee) \cdot m_x = -c_{s,x} \cdot m_x = -(c_s \cdot m)_x.$$

We have to prove the other inclusion, that is, every element $m \in \Gamma^{-s}$ can be divided by $(x(\alpha_s^\vee))_{x \in \mathcal{I}}$ in $\Gamma(\mathcal{I}, \mathcal{B}_W)$.

If $m = (m_x) \in \Gamma^{-s}$ then $m_x = -\varphi^{xs}(m_{xs})$ and so $\rho_{xs,xs \rightarrow x}(m_{xs}) = -\rho_{x,xs \rightarrow x}(m_x)$, since the following diagram commutes:

$$\begin{array}{ccc} \mathcal{B}_W^{xs} & \xrightarrow{\varphi^{xs}} & \mathcal{B}_W^x \\ \downarrow \rho_{xs,xs \rightarrow x} & & \downarrow \rho_{x,xs \rightarrow x} \\ \mathcal{B}_W^{xs \rightarrow x} & \xrightarrow{\varphi^{xs \rightarrow x}} & \mathcal{B}_W^x \end{array} \tag{9}$$

But m is a section so $\rho_{xs,xs \rightarrow x}(m_{xs}) = \rho_{x,xs \rightarrow x}(m_x)$. It follows that $2\rho_{x,xs \rightarrow x}(m_x) = 0$, but, by definition of the canonical sheaf, $\ker \rho_{x,xs \rightarrow x} = \alpha_{xsx^{-1}}^\vee \mathcal{B}_W^x$, that is, $\alpha_{xsx^{-1}}^\vee$ divides m_x in \mathcal{B}_W^x .

Notice that $\alpha_{xsx^{-1}}^\vee = \pm x(\alpha_s^\vee) = \pm c_{s,x}$, i.e. $c_s^{-1} \cdot m \in \bigoplus_{x \in \mathcal{I}} \mathcal{B}_W^x$. We have to verify that $\rho_{x,x \rightarrow tx}(c_{s,x}^{-1} m_x) = \rho_{tx,x \rightarrow tx}(c_{s,tx}^{-1} m_{tx})$ for all $t \in \mathcal{T}$:

$$\begin{aligned} & (c_{s,tx} c_{s,x}) (\rho_{tx,x \rightarrow tx}(c_{s,tx}^{-1} m_{tx}) - \rho_{x,x \rightarrow tx}(c_{s,x}^{-1} m_x)) \\ &= c_{s,x} (\rho_{tx,x \rightarrow tx}(m_{tx})) - c_{s,tx} (\rho_{x,x \rightarrow tx}(m_x)) \\ &= (c_{s,x} - c_{s,tx}) \rho_{tx,x \rightarrow tx}(m_{tx}) + c_{s,tx} (\rho_{tx,x \rightarrow tx}(m_{tx}) - \rho_{x,x \rightarrow tx}(m_x)). \end{aligned}$$

The term on the last line is divisible by α_t^\vee ; indeed,

$$c_{s,x} - c_{s,t} = x(\alpha_s^\vee) - x(\alpha_s^\vee) + \langle x(\alpha_s^\vee), \alpha_t \rangle \alpha_t^\vee \equiv 0 \pmod{\alpha_t}$$

and

$$\rho_{tx,x-tx}(m_{tx}) - \rho_{x,x-tx}(m_x) = 0.$$

Using the GKM-property $c_{s,t}c_{s,x} = tx(\alpha_s^\vee) \cdot x(\alpha_s^\vee)$ is a multiple of α_t^\vee if and only if $xsx^{-1} = t$, that is $xs = tx$. Then, $m_x = -\varphi^{xs}(m_{tx})$, $c_{s,t} = -c_{s,x}$ and, considering that diagram (9) commutes, we obtain

$$\begin{aligned} \rho_{x,x-tx}(c_{s,t}^{-1}m_x) &= -c_{s,t}^{-1}\rho_{x,x-tx}(m_x) \\ &= -c_{s,t}^{-1}(-\rho_{tx,x-tx}(m_{tx})) \\ &= \rho_{tx,x-tx}(c_{s,t}^{-1}m_{tx}). \end{aligned}$$

Otherwise, $xsx^{-1} \neq t$ and α_t^\vee divides $\rho_{tx,x-tx}(c_{s,t}^{-1}m_{tx}) - \rho_{x,x-tx}(c_{s,x}^{-1}m_x)$ and so $\rho_{x,x-tx}(c_{s,x}^{-1}m_x) = \rho_{tx,x-tx}(c_{s,t}^{-1}m_{tx})$. \square

6.3.2. Building $\mathcal{B}_w^{\delta_{ys}}$

Let us denote

$$\Gamma(\mathcal{I}, \mathcal{B}_w) \hookrightarrow \bigoplus_{x \in \mathcal{I}} \mathcal{B}_w^x \longrightarrow \bigoplus_{x \in \mathcal{V}_{\delta y}} \mathcal{B}_w^x \xrightarrow{\oplus \rho_{x,E}} \bigoplus_{E \in \mathcal{E}_{\delta y}} \mathcal{B}^E$$

π_1

Recall that $\mathcal{B}_w^{\delta y} = u_y(\Gamma(\{> y\}, \mathcal{B}_w))$, where u_y was defined as the composition of the following maps

$$\Gamma(\{> y\}, \mathcal{B}_w) \hookrightarrow \bigoplus_{x > y} \mathcal{B}_w^x \longrightarrow \bigoplus_{x \in \mathcal{V}_{\delta y}} \mathcal{B}_w^x \xrightarrow{\oplus \rho_{x,E}} \bigoplus_{E \in \mathcal{E}_{\delta y}} \mathcal{B}_w^E$$

u_y

Remark 6.1. Since \mathcal{B}_w is flabby and \mathcal{I} and $\{> y\}$ are both open sets, we get

$$\pi_1(\Gamma(\mathcal{I}, \mathcal{B}_w)) = u_y(\Gamma(\{> y\}, \mathcal{B}_w)) = \mathcal{B}_w^{\delta y}. \tag{10}$$

Now, let us denote

$$\Gamma(\mathcal{I}, \mathcal{B}_w) \hookrightarrow \bigoplus_{x \in \mathcal{I}} \mathcal{B}_w^x \longrightarrow \bigoplus_{x \in \mathcal{V}_{\delta y}} \mathcal{B}_w^{xs} \xrightarrow{\oplus \rho_{xs,Es}} \bigoplus_{E \in \mathcal{E}_{\delta y}} \mathcal{B}_w^{Es}$$

π_2

and define $\widetilde{\mathcal{B}_w^{\delta_{ys}}} := \pi_2(\Gamma(\mathcal{I}, \mathcal{B}_w))$.

Lemma 6.4.

- (i) $\widetilde{\mathcal{B}}_W^{\delta y} = \pi_1(\Gamma(\mathcal{I}, \mathcal{B}_W)) = \pi_1(\Gamma^S)$.
- (ii) $\widetilde{\mathcal{B}}_W^{\delta ys} = \pi_2(\Gamma(\mathcal{I}, \mathcal{B}_W)) = \pi_2(\Gamma^S)$.

Proof. (i) Let $m \in \Gamma(\mathcal{I}, \mathcal{B}_W)$. Then, by Lemma 6.3, $m = m' + c_s \cdot m''$, with $m', m'' \in \Gamma^S$ and, if $m' = (m'_x)$, $m'' = (m''_x)$,

$$\pi_1(m) = (\rho_{x,E}(m'_x))_{x \in \mathcal{V}: y \rightarrow x \in \mathcal{E}} + (\rho_{x,E}(x(\alpha_s^\vee) \cdot m''_x))_{x \in \mathcal{V}: y \rightarrow x \in \mathcal{E}}.$$

If $E : y \rightarrow x \in \mathcal{E}_{\delta y}$, then there exists a reflection $t \in \mathcal{T}$ such that $x = ty$ and we have

$$x(\alpha_s^\vee) = ty(\alpha_s^\vee) = y(\alpha_s^\vee) - \langle y(\alpha_s^\vee), \alpha_t \rangle \alpha_t^\vee.$$

But, by definition, $\rho_{x,E}$ is a surjective map whose kernel is $l(E)\mathcal{B}_W^x = \alpha_t^\vee \mathcal{B}_W^x$ and


$$\begin{aligned} \rho_{x,E}(x(\alpha_s^\vee) \cdot m''_x) &= \rho_{x,E}(y(\alpha_s^\vee) \cdot m''_x) - \langle y(\alpha_s^\vee), \alpha_t \rangle \rho_{x,E}(\alpha_t^\vee \cdot m''_x) \\ &= \rho_{x,E}(y(\alpha_s^\vee) \cdot m''_x). \end{aligned}$$

We conclude that $\pi_1(m) = \pi_1(m' + \overline{y(\alpha_s^\vee)} \cdot m'')$, where $\overline{y(\alpha_s^\vee)}$ is the element of the structure algebra, whose components are all equal to $y(\alpha_s^\vee)$. Clearly, $m' + \overline{y(\alpha_s^\vee)} \cdot m'' \in \Gamma^S$ and we get the claim.

(ii) As in (i). \square

Lemma 6.5. There is an isomorphism of S_k -modules $\tau : \mathcal{B}_W^{\delta y} \rightarrow \widetilde{\mathcal{B}}_W^{\delta ys}$ given by $(m_E)_{E \in \mathcal{E}_{\delta y}} \mapsto (\varphi^E(m_E))_{E \in \mathcal{E}_{\delta y}}$, that is for all $m \in \Gamma^S$, $\tau \circ \pi_1(m) = \pi_2(m)$.

Proof. The element $(m_E)_{E \in \mathcal{E}_{\delta y}} \in \mathcal{B}_W^{\delta y}$ if and only if there exists an element $m \in \Gamma(\{> y\}, \mathcal{B}_W)$ such that $u_y(m) = (m_E)_{E \in \mathcal{E}_{\delta y}}$. We have already noticed that this is the case if and only if there is an element $m' \in \Gamma(\mathcal{I}, \mathcal{B})$ such that $\pi_1(m') = (m_E)_{E \in \mathcal{E}_{\delta y}}$. From the previous lemma, we know that this is equivalent to the existence of an $\tilde{m} \in \Gamma^S$ such that $\pi_1(\tilde{m}) = (m_E)_{E \in \mathcal{E}_{\delta y}}$. But, since the squares in the following diagram are all commutative,



we get $(\varphi^E(m_E))_{E \in \mathcal{E}_{\delta y}} = (\oplus \varphi^E) \circ \pi_1(\tilde{m}) = \pi_2(\tilde{m}) \in \widetilde{\mathcal{B}}_W^{\delta ys}$.

Analogously, $(m_{Es})_{E \in \mathcal{E}_{\delta y}} \in \widetilde{\mathcal{B}}_W^{\delta ys}$ if and only if $((\varphi^E)^{-1}(m_{Es}))_{E \in \mathcal{E}_{\delta y}} \in \mathcal{B}_W^{\delta y}$. \square

Let us denote by $\rho : \mathcal{B}_W^y \rightarrow \mathcal{B}_W^y / \alpha_s^\vee \cdot \mathcal{B}_W^y$ the canonical quotient map.

Lemma 6.6. *We have*

$$\mathcal{B}_w^{\delta ys} = \{(\tau \circ d_y(m_y), \rho(m_y)) \in \widetilde{\mathcal{B}_w^{\delta ys}} \oplus (\mathcal{B}_w^y / \alpha_s^\vee \cdot \mathcal{B}_w^y)\}. \quad (11)$$

Proof.

$$\begin{aligned} \mathcal{B}_w^{\delta ys} &= u_{ys}(\Gamma(\{> ys\}, \mathcal{B}_w)) \\ &= u_{ys}(\{(m, m_y) \in \Gamma(\mathcal{I}, \mathcal{B}_w) \oplus \mathcal{B}_w^y \mid u_y(m_{\{>y\}}) = d_y(m_y)\}) \end{aligned}$$

by Remark 6.1

$$\begin{aligned} &= u_{ys}(\{(m, m_y) \in \Gamma(\mathcal{I}, \mathcal{B}_w) \oplus \mathcal{B}_w^y \mid \pi_1(m) = d_y(m_y)\}) \\ &= \{(\pi_2(m), \rho(m_y)) \mid m \in \Gamma(\mathcal{I}, \mathcal{B}_w), m_y \in \mathcal{B}_w^y, \pi_1(m) = d_y(m_y)\} \end{aligned}$$

by Lemma 6.4

$$= \{(\pi_2(m), \rho(m_y)) \mid m \in \Gamma^s, m_y \in \mathcal{B}_w^y, \pi_1(m) = d_y(m_y)\}$$

by Lemma 6.5

$$\begin{aligned} &= \{(\tau \circ \pi_1(m), \rho(m_y)) \mid m \in \Gamma^s, m_y \in \mathcal{B}_w^y, \pi_1(m) = d_y(m_y)\} \\ &= \{(\tau \circ d_y(m_y), \rho(m_y)) \mid m_y \in \mathcal{B}_w^y\}. \quad \square \end{aligned}$$

From the lemma above, it follows immediately, that there is a surjective map of S_k -modules $\mathcal{B}_w^y \rightarrow \mathcal{B}_w^{\delta ys}$ given by $m_y \mapsto (\tau \circ d_y(m_y), \rho(m_y))$ and this concludes the proof of Theorem 6.1.

6.4. Rational smoothness and p -smoothness of the flag variety

We have an easy corollary of Theorem 6.1:

Corollary 6.1. *Let \mathcal{W} be a finite Weyl group and w_0 its longest element. Then $\mathcal{B}_{w_0}^y \cong S_k$ for any $y \in \mathcal{W}$ and any k .*

Proof. We proceed by induction on $n = \ell(w_0) - \ell(y)$. If $n = 0$, by definition, $\mathcal{B}_{w_0}^{w_0} \cong S_k$. If $n \geq 1$ then there exists a simple reflection $s \in \mathcal{S}$ such that $ys > y$ (so, $\ell(w_0) - \ell(ys) = n - 1$). Actually, $w_0 s < w_0$ for any $s \in \mathcal{S}$ and, by Theorem 6.1 and inductive hypothesis, we have $\mathcal{B}_{w_0}^y \cong \mathcal{B}_{w_0}^{ys} \cong S_k$. \square

Remark 6.2. If $k = \mathbb{Q}$ the result above corresponds to the (rational) smoothness of flag varieties, while if k is a field of characteristic p it gives their p -smoothness (cf. [15]). Our proof is based only on the definition of canonical sheaf; we do not use Fiebig's multiplicity one results (see [12]), nor the geometry of the corresponding flag varieties.

6.5. Parabolic setting

Let $J \subseteq \mathcal{S}$ be such that $\mathcal{W}_J = \langle J \rangle$ is finite with longest element w_J . Let \mathcal{W}^J be the set of minimal representatives of the equivalence classes $\mathcal{W}/\mathcal{W}_J$. For $w \in \mathcal{W}^J$, denote by \mathcal{B}_{ww_J} , resp. \mathcal{B}_w^J , the canonical sheaf on \mathcal{G}_{ww_J} , resp. on \mathcal{G}_w^J . It is now easy to see that:

Lemma 6.7. *Let \mathcal{W}_J and w_J be as above and consider $x, w \in \mathcal{W}^J$ such that $x \leq w$, then $\mathcal{B}_{ww_J}^x \cong \mathcal{B}_{ww_J}^{xu}$ for any $u \in \mathcal{W}_J$.*

Proof. We proceed by induction on $\ell(u)$. Clearly there is nothing to prove if $\ell(u) = 0$. If $\ell(u) > 0$ then there exists an $s \in \mathcal{S}$ such that $us < u$ and so by the inductive hypothesis, we get $\mathcal{B}_{ww_J}^x \cong \mathcal{B}_{ww_J}^{xus}$. Now for any $s \in J$, $ww_Js < ww_J$ and $xus, xu \leq ww_J$ and by Theorem 6.1 we obtain the claim. \square

Theorem 6.2. Let (\mathcal{G}_{ww_J}, k) be a GKM-pair and let \mathcal{W}^J and w_J be as above. If $y, w \in \mathcal{W}^J$ and $y \leq w$, then there is an isomorphism of S_k -modules

$$(\mathcal{B}_{ww_J})^{y w_J} \cong (\mathcal{B}_w^J)^y.$$

Proof. We proceed by induction on $n = \ell(w) - \ell(y)$. If $n = 0$ the statement is trivial. Suppose we have a collection of isomorphisms of S_k -modules $\eta_x : (\mathcal{B}_w^J)^x \rightarrow (\mathcal{B}_{ww_J})^{x w_J}$ for any x such that $\ell(w) - \ell(x) < n$.

There is a natural injective homomorphism,

$$j : \Gamma(\{> y\}, \mathcal{B}_w^J) \rightarrow \Gamma(\{> y w_J\}, \mathcal{B}_{ww_J}),$$

defined by setting $(m_x)_{x \in (y, w] \subset \mathcal{W}^J} \mapsto (\tilde{m}_z)_{z \in (y w_J, w w_J] \subset \mathcal{W}}$, where $\tilde{m}_z := \psi^z(\eta_x(m_x))$ if $z \in x \mathcal{W}_J$ and $\psi^z : \mathcal{B}_{ww_J}^{x w_J} \rightarrow \mathcal{B}_{ww_J}^z$ is an isomorphism (it exists by Lemma 6.7).

We will show that such a homomorphism induces an isomorphism $(\mathcal{B}_{ww_J})^{\delta y w_J} \cong (\mathcal{B}_w^J)^{\delta y}$. Then, by the unicity of projective cover, the statement will follow.

Let $z \in (y w_J, w w_J]$, $z = xu$, for some $x > y \in \mathcal{W}^J$, $u \in \mathcal{W}_J$ and $u = s_1 \dots s_r$ a reduced expression with $s_i \in J$ for every i . Moreover, let $(n_v) \in \Gamma(\{> y w_J\}, \mathcal{B}_{ww_J})$. We prove by induction on $\ell(u) = r$ that there exists a section $(p_v) \in \Gamma(\{> y w_J\}, \mathcal{B}_{ww_J})$ such that $p_{x s_1 \dots s_i} = \psi^{x s_1 \dots s_i}(\eta_x(m_x))$ for some $m_x \in (\mathcal{B}_w^J)^x$ for any $i = 0, \dots, r$ and such that $u_{y w_J}((p_v)) = u_{y w_J}((n_v))$.

For the base step we have $r = 0$ and there is nothing to prove.

If $z = (x s_1 s_2 \dots s_{r-1}) s_r$ then, by the inductive hypothesis, there exists a section $(q_v) \in \Gamma(\{> y w_J\}, \mathcal{B}_{ww_J})$ and an element $m_x \in (\mathcal{B}_w^J)^x$ such that $q_{x s_1 \dots s_i} = \psi^{x s_1 \dots s_i}(\eta_x(m_x))$ and $u_y((q_v)) = u_y((n_v))$ for $i = 0, \dots, r - 1$. Thus, by Lemma 6.4, the element $(p_v) \in \bigoplus_{v > y w_J} \mathcal{B}^y$ so that

$$p_{y s_1 \dots s_{r-1} s_r} = \varphi^{y s_1 \dots s_{r-1}}(p_{y s_1 \dots s_{r-1}})$$

and

$$p_{x s_1 \dots s_i} = q_{x s_1 \dots s_i} = \psi^{x s_1 \dots s_i}(\eta_x(m_x)) \quad \forall i < r$$

is a section on $\{> y w_J\}$ and verifies $u_{y w_J}((\tilde{n}_v)) = u_{y w_J}((n_v))$.

Finally, from the proof of Lemma 6.7 it follows that

$$\varphi^{y s_1 \dots s_{r-1}}(p_{y s_1 \dots s_{r-1}}) = \varphi^{y s_1 \dots s_{r-1}}(\psi^{y s_1 \dots s_{r-1}}(\eta_x(m_x))) = \psi^{x s_1 \dots s_r}(\eta_x(m_x)). \quad \square$$

The theorem above is the analogue of the following theorem, due to Deodhar:

Theorem 6.3. (See [7].) Let \mathcal{W} be a Weyl group with \mathcal{S} , set of simple reflections, and $J \subseteq \mathcal{S}$ such that \mathcal{W}_J is finite. Let w_J be the longest element of \mathcal{W}_J and $y, w \in \mathcal{W}^J$, then $P_{y, w}^{J, -1} = P_{y w_J, w w_J}$.

Acknowledgments

I would like to thank Peter Fiebig for drawing my attention to this problem and for helpful discussions, and Rocco Chirivì and Corrado De Concini for useful conversations. I would also like to thank the INdAM for funding part of my stay at the Friedrich-Alexander-Universität of Erlangen-Nürnberg. Finally, I am grateful to the Hausdorff Research Institute for Mathematics in Bonn for its hospitality during the program “On the Interaction of Representation Theory with Geometry and Combinatorics”, where the last version of this paper was written.

References

- [1] H.H. Andersen, J.C. Jantzen, W. Soergel, Representation of quantum groups at a p -th root of unity and of semisimple groups in characteristic p : Independence of p , *Astérisque* 220 (1994) 1–320.
- [2] A. Beilinson, J. Bernstein, Localisation de \mathfrak{g} -modules, *C. R. Acad. Sci. Paris* 292 (1981) 15–18.
- [3] T. Braden, R. MacPherson, From moment graphs to intersection cohomology, *Math. Ann.* 321 (2001) 533–551.
- [4] A. Björner, F. Brenti, *Combinatorics of Coxeter Groups*, Springer, New York, 2005.
- [5] F. Brenti, F. Caselli, M. Marietti, Special matchings and Kazhdan–Lusztig polynomials, *Adv. Math.* 202 (2006) 555–601.
- [6] J.-L. Brylinski, M. Kashiwara, Démonstration de la conjecture de Kazhdan et Lusztig sur les modules de Verma, *C. R. Acad. Sci. Paris* 27 (1980) 373–376.
- [7] V. Deodhar, On some geometric aspects of Bruhat orderings II. The parabolic analogue of Kazhdan–Lusztig polynomials, *J. Algebra* 111 (1987) 483–506.
- [8] M. Dyer, Hecke algebras and reflections in Coxeter groups, Ph.D. thesis, University of Sydney, 1987.
- [9] P. Fiebig, Sheaves on moment graphs and a localization of Verma flags, *Adv. Math.* 217 (2008) 683–712.
- [10] P. Fiebig, Moment graphs in representation theory and geometry, Skript der Vorlesung zu Bruhatgraphen im Wintersemester 2008/09, available at http://www.algeo.math.uni-erlangen.de/fileadmin/algeo/users/fiebig/Skripten/Skript_MomGra.pdf.
- [11] P. Fiebig, Lusztig’s conjecture as a moment graph problem, *Bull. Lond. Math. Soc.* 42 (6) (2010) 957–972.
- [12] P. Fiebig, The multiplicity one case of Lusztig’s conjecture, *Duke Math. J.* 153 (2010) 551–571.
- [13] P. Fiebig, Sheaves on affine Schubert varieties, modular representations and Lusztig’s conjecture, *J. Amer. Math. Soc.* 24 (2011) 133–181.
- [14] P. Fiebig, An upper bound on the exceptional characteristics for Lusztig’s character formula, *J. Reine Angew. Math. (Crelle’s J.)*, <http://dx.doi.org/10.1515/crelle.2011.170>.
- [15] P. Fiebig, G. Williamson, Parity sheaves, moment graphs and the p -smooth locus of Schubert varieties, arXiv:1008.0719, 2010.
- [16] M. Goresky, R. Kottwitz, R. MacPherson, Equivariant cohomology, Koszul duality and the localization theorem, *Invent. Math.* 131 (1) (1998) 23–83.
- [17] J.E. Humphreys, *Reflection Groups and Coxeter Groups*, Cambridge University Press, Cambridge, 1990.
- [18] D. Juteau, C. Mautner, G. Williamson, Parity sheaves, arXiv:0906.2994, 2009.
- [19] V. Kac, *Infinite Dimensional Lie Algebras*, Progr. Math., vol. 44, Birkhäuser Boston Inc., Boston, MA, 1983.
- [20] D. Kazhdan, G. Lusztig, Representation of Coxeter groups and Hecke algebras, *Invent. Math.* 53 (1979) 165–184.
- [21] D. Kazhdan, G. Lusztig, Schubert varieties and Poincaré duality, in: *Geometry of the Laplace Operator*, in: Proc. Sympos. Pure Math., vol. 36, Amer. Math. Soc., 1980, pp. 185–203.
- [22] D. Kazhdan, G. Lusztig, Tensor structures arising from affine Lie algebras, I, *J. Amer. Math. Soc.* 6 (1993) 905–947;
D. Kazhdan, G. Lusztig, Tensor structures arising from affine Lie algebras, II, *J. Amer. Math. Soc.* 6 (1993) 949–1011;
D. Kazhdan, G. Lusztig, Tensor structures arising from affine Lie algebras, III, *J. Amer. Math. Soc.* 7 (1994) 335–381;
D. Kazhdan, G. Lusztig, Tensor structures arising from affine Lie algebras, IV, *J. Amer. Math. Soc.* 7 (1994) 383–453.
- [23] M. Kashiwara, T. Tanisaki, Kazhdan–Lusztig conjecture for affine Lie algebras with negative level, *Duke Math. J.* 77 (1) (1995) 21–62.
- [24] S. Kumar, *Kac–Moody Groups, Their Flag Varieties and Representation Theory*, Progr. Math., vol. 204, Birkhäuser Boston Inc., Boston, MA, 2002.
- [25] G. Lusztig, Some problems in the representation theory of finite Chevalley groups, in: *The Santa Cruz Conference on Finite Groups*, Univ. California, Santa Cruz, CA, 1979, in: Adv. Proc. Sympos. Pure Math., vol. 37, Amer. Math. Soc., Providence, RI, 1980, pp. 313–317.
- [26] G. Williamson, Checking Lusztig’s conjecture around the Steinberg weight, in: *Representations of Finite Groups 2012*, Oberwolfach Report, 16/2012.