

DEGENERATE FLAG VARIETIES AND SCHUBERT VARIETIES: A CHARACTERISTIC FREE APPROACH

G. CERULLI IRELLI, M. LANINI, P. LITTELMANN

ABSTRACT. We consider the PBW filtrations over \mathbb{Z} of the irreducible highest weight modules in type A_n and C_n . We show that the associated graded modules can be realized as Demazure modules for group schemes of the same type and doubled rank. We deduce that the corresponding degenerate flag varieties are isomorphic to Schubert varieties in any characteristic.

INTRODUCTION

Introduced by Evgeny Feigin in 2010, degenerate flag varieties naturally arise from a representation theoretic context. In fact, given a finite dimensional, highest weight irreducible module $V(\lambda)$ for a simple finite dimensional, complex Lie algebra, the corresponding degenerate flag variety $\mathcal{F}\ell(\lambda)^a$ is the closure of a certain highest weight orbit in the projectivization of $V(\lambda)^a$, a degenerate version of $V(\lambda)$.

If the algebra one starts with is of type A_n or C_n , it was shown in [CL] that, surprisingly, degenerate flag varieties can be realized as Schubert varieties in a partial flag variety of same type and bigger rank. It is hence natural to ask whether also the modules $V(\lambda)^a$ are isomorphic to some already investigated objects. The aim of this paper is to address such a question and provide a positive answer to it. Feigin's degeneration procedure can be carried out over \mathbb{Z} (cf. [FFL3]) and it is in this generality that we decided to approach the problem.

Our main theorem is the realization of $V(\lambda)^a$ as a Demazure module for a group scheme of same type and doubled rank. This fact allows us to recover, as a corollary, the above-mentioned realization of $\mathcal{F}\ell(\lambda)^a$ as a Schubert variety. While the arguments in [CL] relied on a linear algebraic description of the degenerate flag variety due to Feigin (cf. [F1]), the proof we obtain here only uses the definition of $\mathcal{F}\ell(\lambda)^a$ as a closure of an highest weight orbit and it is hence more conceptual.

In what follows, we describe more in detail the main results of this article.

For simplicity let us start with the complex algebraic group $SL_n(\mathbb{C})$ and its Lie algebra $\mathfrak{g} = \mathfrak{sl}_n$. We fix a Cartan decomposition $\mathfrak{g} = \mathfrak{n}^- \oplus \mathfrak{h} \oplus \mathfrak{n}^+$, where \mathfrak{n} is the subalgebra of strictly upper triangular matrices, \mathfrak{h} is the Cartan subalgebra consisting of diagonal matrices and \mathfrak{n}^- is the subalgebra of strictly lower triangular matrices. Let $\mathfrak{b} = \mathfrak{h} \oplus \mathfrak{n}^+$ be the corresponding Borel subalgebra of \mathfrak{g} and let $B \subset G$ be the Borel subgroup with Lie algebra \mathfrak{b} .

We use the notation \tilde{B} , $\tilde{\mathfrak{b}}$, $\tilde{\mathfrak{n}}^+$, $\tilde{\mathfrak{h}}$ and $\tilde{\mathfrak{n}}^-$ for the corresponding subgroup of $\tilde{G} = SL_{2n}(\mathbb{C})$ and subalgebras of $\tilde{\mathfrak{g}} = \mathfrak{sl}_{2n}$. Let $\mathfrak{n}^{-,a} \subset \mathfrak{sl}_{2n}$ respectively $N^{-,a} \subseteq SL_{2n}(\mathbb{C})$ be the following commutative Lie subalgebra respectively commutative unipotent subgroup:

$$(1) \quad \mathfrak{n}^{-,a} := \left\{ \begin{pmatrix} 0 & N \\ 0 & 0 \end{pmatrix} \in \mathfrak{sl}_{2n} \mid N \in \mathfrak{n}^- \right\}, N^{-,a} := \left\{ \begin{pmatrix} \mathbb{I} & N \\ 0 & \mathbb{I} \end{pmatrix} \in SL_{2n} \mid N \in \mathfrak{n}^- \right\}.$$

We view $\mathfrak{n}^{-,a}$ as the abelianization of \mathfrak{n}^- , i.e. we have the canonical vector space isomorphism between the two vector spaces, but $\mathfrak{n}^{-,a}$ is endowed with the trivial Lie bracket. The enveloping algebra of $\mathfrak{n}^{-,a}$ is $S^\bullet(\mathfrak{n}^{-,a})$. The embedding $\mathfrak{n}^{-,a} \hookrightarrow \tilde{\mathfrak{b}}$ induces an embedding $S^\bullet(\mathfrak{n}^{-,a}) \hookrightarrow U(\tilde{\mathfrak{b}})$, so any $U(\tilde{\mathfrak{b}})$ -module inherits in a natural way the structure of a $S^\bullet(\mathfrak{n}^{-,a})$ -module.

A well investigated class of $U(\tilde{\mathfrak{b}})$ -modules are the Demazure modules: let μ be a dominant integral weight for $\tilde{\mathfrak{g}}$ and let $\tilde{V}(\mu)$ be the corresponding irreducible representation. For an element w of the Weyl group \tilde{W} of $\tilde{\mathfrak{g}}$, the weight space $\tilde{V}(\mu)_{w\mu}$ of weight $w\mu$ is one-dimensional, fix a generator $v_{w\mu}$. Recall that the Demazure submodule $\tilde{V}(\mu)_w$ is by definition the cyclic $U(\tilde{\mathfrak{b}})$ -module generated by $v_{w\mu}$, i.e. $\tilde{V}(\mu)_w = U(\tilde{\mathfrak{b}}) \cdot v_{w\mu}$, and the Schubert variety $X(w)$ is the closure of the orbit $\tilde{B} \cdot [v_{w\mu}] \subseteq \mathbb{P}(\tilde{V}(\mu))$.

A special class of $S^\bullet(\mathfrak{n}^{-,a})$ -modules has been investigated in [FFL1, FFL2]. Let λ be a dominant integral weight for \mathfrak{g} , let $V(\lambda)$ be the corresponding irreducible representation and fix a highest weight vector v_λ . The PBW filtration on $U(\mathfrak{n}^-)$ induces a filtration on the cyclic $U(\mathfrak{n}^-)$ -module $V(\lambda) = U(\mathfrak{n}^-) \cdot v_\lambda$, the associated graded space $V^a(\lambda) := \text{gr } V(\lambda)$ becomes a module for the associated graded algebra $S^\bullet(\mathfrak{n}^-) := \text{gr } U(\mathfrak{n}^-) \simeq S^\bullet(\mathfrak{n}^{-,a})$.

The action of $\mathfrak{n}^{-,a}$ on $V^a(\lambda)$ can be integrated to an action of $N^{-,a}$, in analogy with the classical case we call the closure of the orbit $\mathcal{F}_\lambda^a := \overline{N^{-,a} \cdot [v_\lambda]} \subseteq \mathbb{P}(V^a(\lambda))$ the degenerate flag variety.

The aim of this article is to connect these two constructions and extend the results in [CL] to an algebraically closed field k of arbitrary characteristic. In fact, the results hold even over \mathbb{Z} . For simplicity, we formulate them in the introduction for an algebraically closed field k . In the following we consider the case $G = SL_n(k)$ and $\tilde{G} = SL_{2n}(k)$, respectively $G = Sp_{2m}(k)$, and $\tilde{G} = Sp_{4m}(k)$, and we replace the irreducible module of highest weight λ by the Weyl module of highest weight λ , using the same notation $V(\lambda)$. For the precise description of the highest weight $\Psi(\lambda)$ see Definitions 2.1 and 5.3, for a description of the Weyl group element $\tau \in \tilde{W}$ see Definitions 2.2 and 5.1, and the construction of the Lie algebra $\mathfrak{n}^{-,a}$ in the symplectic case can be found in Section 5. For a dominant G -weight λ let λ^* be the dual dominant weight, so for the symplectic case we have $\lambda = \lambda^*$, and in the SL_n case we have $\lambda^* = \sum_{i=1}^{n-1} m_i \omega_{n-i}$ for $\lambda = \sum_{i=1}^{n-1} m_i \omega_i$ (notation as in [B]).

Theorem. *Let λ be a dominant G -weight.*

- i) The Demazure submodule $\tilde{V}_k(\Psi(\lambda^*))_\tau$ of the \tilde{G} -module $\tilde{V}_k(\Psi(\lambda^*))$ is, as an $\mathfrak{n}^{-,a}$ -module, isomorphic to the abelianized module $V^a(\lambda)$.*

ii) The Schubert variety $X(\tau) \subset \mathbb{P}(\tilde{V}(\Psi(\lambda^*))_\tau)$ is isomorphic to the degenerate flag variety $\mathcal{F}^a(\lambda)$, and this isomorphism induces an $S^\bullet(\mathfrak{n}^{-,a})$ -module isomorphism

$$H^0(X(\tau), \mathcal{L}_{\Psi(\lambda^*)}) \simeq (V^a(\lambda))^*.$$

Using the isomorphism above, we deduce the defining relations for $V^a(\lambda)$ from the defining relations of the Demazure module. Translated back into the language of the abelianized algebras we get the following: in the SL_n -case let $R^{++} = R^+$ be the set of positive roots, in the symplectic case set

$$R^{++} = \{\epsilon_i - \epsilon_j \mid 1 \leq i < j \leq m\} \cup \{2\epsilon_i \mid 1 \leq i \leq m\}.$$

Corollary. *The abelianized module $V^a(\lambda)$ is as a cyclic $S^\bullet(\mathfrak{n}^{-,a})$ -module isomorphic to $S^\bullet(\mathfrak{n}^{-,a})/I(\lambda)$, where $I(\lambda)$ is the ideal:*

$$I(\lambda) = S^\bullet(\mathfrak{n}^{-,a})(U(\mathfrak{n}^+) \circ \text{span}\{f_\alpha^{(\langle \lambda, \alpha^\vee \rangle + 1)}; \alpha \in R^{++}\}) \subseteq S^\bullet(\mathfrak{n}^{-,a}).$$

The identification of the degenerate flag variety as a Schubert variety implies immediately (see also [FF, FFIL]):

Corollary. *The degenerate flag variety $\mathcal{F}^a(\lambda)$ is projectively normal and has rational singularities.*

1. SOME SPECIAL COMMUTATIVE UNIPOTENT SUBGROUPS

Let k be a field. Given a subspace $\mathfrak{N} \subseteq M_n(k)$ and a vector space automorphism $\eta : \mathfrak{N} \rightarrow \mathfrak{N}$, denote by $\mathfrak{N}_\eta^a \subseteq M_{2n}(k)$ respectively $N_\eta^a \subseteq GL_{2n}(k)$ the following commutative nilpotent Lie subalgebra of $M_{2n}(k)$, respectively commutative unipotent subgroup of $GL_{2n}(k)$:

$$\mathfrak{N}_\eta^a := \left\{ \begin{pmatrix} 0 & \eta(A) \\ 0 & 0 \end{pmatrix} \mid A \in \mathfrak{N} \right\}, \quad N_\eta^a := \left\{ \begin{pmatrix} \mathbb{I} & \eta(A) \\ 0 & \mathbb{I} \end{pmatrix} \mid A \in \mathfrak{N} \right\}.$$

If $\mathfrak{N} \subseteq M_n(k)$ is a Lie subalgebra, then we think of \mathfrak{N}_η^a as an abelianized version of \mathfrak{N} . Similarly one may think of N_η^a as an abelianized version of a subgroup $N \subseteq GL_{2n}(k)$. We will be more precise in the following examples.

Example 1.1. Let k be an algebraically closed field of characteristic zero. We fix as a maximal torus $T \subset SL_n$ the subgroup of diagonal matrices, and let B be the Borel subgroup of upper triangular matrices. Let us denote by \mathfrak{sl}_n , \mathfrak{b} and \mathfrak{h} the corresponding Lie algebras and let $\mathfrak{g} = \mathfrak{n}^- \oplus \mathfrak{h} \oplus \mathfrak{n}^+$ be the Cartan decomposition. The choice of a maximal torus and a Borel subgroup as above determines the set of positive roots Φ^+ and hence, according to the adjoint action of \mathfrak{h} , the root space decomposition $\mathfrak{n}^- = \bigoplus_{\alpha \in \Phi^+} \mathfrak{n}_{-\alpha}^-$. In this example we set $\mathfrak{N} = \mathfrak{n}^-$, and $N = U^-$ is the unipotent radical of the opposite Borel subgroup B^- . The map η is the identity map, so we just omit it. We write henceforth $\mathfrak{n}^{-,a}$ for $\mathfrak{N}_\eta^a \subset \mathfrak{sl}_{2n}(k)$ and $N^{-,a}$ for $N_\eta^a \subset SL_{2n}$.

Note that $\mathfrak{n}^{-,a} \subset \mathfrak{sl}_{2n}$ is a Lie subalgebra of the Borel subalgebra $\tilde{\mathfrak{b}} \subset \mathfrak{sl}_{2n}$ and $N^{-,a}$ is an abelian subgroup of the Borel subgroup $\tilde{B} \subset SL_{2n}$ (of upper triangular matrices). We can think of $N^{-,a}$ as an abelianized version of U^- .

The subgroup $N^{-,a}$, as well as the Lie algebra $\mathfrak{n}^{-,a}$, is stable under conjugation with respect to the maximal torus $\tilde{T} \subset SL_{2n}$, where $\tilde{T} \subset SL_{2n}$ consists of the diagonal matrices. The group $N^{-,a}$ hence decomposes as a product of root subgroups of the group SL_{2n} , and $\mathfrak{n}^{-,a}$ decomposes into the direct sum of root subspaces for the Lie algebra \mathfrak{sl}_{2n} . We get an induced map $\phi : \Phi^+ \rightarrow \tilde{\Phi}^+$ between the set of positive roots of \mathfrak{sl}_n and the positive roots of \mathfrak{sl}_{2n} , such that $\mathfrak{n}^{-,a} = \bigoplus_{\alpha \in \Phi^+} \mathfrak{n}_{\phi(\alpha)}^{-,a}$.

Example 1.2. Let k be an algebraically closed field of characteristic 0. Let $\{e_1, \dots, e_{2n}\}$ be the canonical basis of k^{2n} , we fix a non-degenerate skew symmetric form by the conditions $\langle e_i, e_j \rangle = \delta_{j, 2n-i+1} = -\langle e_j, e_i \rangle$ for $1 \leq i \leq n$, $1 \leq j \leq 2n$. Let Sp_{2m} be the associated symplectic group. By the choice of the form we can fix as a Borel subgroup B the subgroup of upper triangular matrices in Sp_{2m} , and let T be its maximal torus consisting of diagonal matrices. Let us denote by \mathfrak{sp}_{2m} , \mathfrak{b} and \mathfrak{h} the corresponding Lie algebras and let $\mathfrak{g} = \mathfrak{n}^- \oplus \mathfrak{h} \oplus \mathfrak{n}^+$ be the Cartan decomposition.

The choice of the torus and the Borel subgroup as above determines a set of positive roots Φ^+ and hence, according to the adjoint action of \mathfrak{h} , the root space decomposition $\mathfrak{n}^- = \bigoplus_{\alpha \in \Phi^+} \mathfrak{n}_{-\alpha}^-$. In this example we set $\mathfrak{N} = \mathfrak{n}^-$, and $N = U^-$ is the unipotent radical of the opposite Borel subgroup B^- . Let $\eta : \mathfrak{n}^- \rightarrow \mathfrak{n}^-$ be the linear map sending a matrix $(m_{i,j})_{1 \leq i,j \leq 2n}$ to the matrix $(m'_{i,j})_{1 \leq i,j \leq 2n}$, where $m'_{i,j} = m_{i,j}$ if $i \leq n$ or $j \leq n$ and $m'_{i,j} = -m_{i,j}$ if both indices are strictly larger than n . We write henceforth $\mathfrak{n}_{\eta}^{-,a}$ for $\mathfrak{N}_{\eta}^a \subset \mathfrak{sp}_{4n}(k)$ and $N_{\eta}^{-,a}$ for $N_{\eta}^a \subset Sp_{4n}$.

Note that $\mathfrak{n}_{\eta}^{-,a} \subset \mathfrak{sp}_{4n}$ is a Lie subalgebra of the Borel subalgebra (of upper triangular matrices) $\mathfrak{b} \subset \mathfrak{sp}_{4n}$ and $N_{\eta}^{-,a}$ is an abelian subgroup of the Borel subgroup $\tilde{B} \subset Sp_{4n}$ (of upper triangular matrices). We can think of $N_{\eta}^{-,a}$ as an abelianized version of U^- .

The subgroup $N_{\eta}^{-,a}$ is stable under conjugation with respect to the maximal torus $\tilde{T} \subset Sp_{4n}$, where $\tilde{T} \subset Sp_{4n}$ consists of the diagonal matrices. The group $N_{\eta}^{-,a}$ hence decomposes as a product of root subgroups of the group Sp_{4n} and $\mathfrak{n}^{-,a}$ decomposes into the direct sum of root subspaces for the Lie algebra \mathfrak{sp}_{4n} . We get an induced map $\phi : \Phi^+ \rightarrow \tilde{\Phi}^+$ between the set of positive roots of \mathfrak{sp}_{2n} and the positive roots of \mathfrak{sp}_{4n} , such that $\mathfrak{n}_{\eta}^{-,a} = \bigoplus_{\alpha \in \Phi^+} \mathfrak{n}_{\phi(\alpha), \eta}^{-,a}$.

Example 1.3. To get a characteristic free approach for G as above, let $G_{\mathbb{Z}}$ be a split and connected simple algebraic \mathbb{Z} -group of type A_n or C_n . For any commutative ring A set $G_A = (G_{\mathbb{Z}})_A$, and for a field set $G = G_k$, for this and the following see also [J]. Then G_k is for any algebraically closed field a reduced k -group, and it is connected and reductive. Its Lie algebra $\text{Lie}(G_{\mathbb{Z}})$ is a free Lie algebra of finite rank and $\text{Lie } G_k = \text{Lie}(G_{\mathbb{Z}}) \otimes_{\mathbb{Z}} k$. Let $T_{\mathbb{Z}} \subset G_{\mathbb{Z}}$ be a split maximal torus and set $T_A = (T_{\mathbb{Z}})_A$ for any ring A and $T = T_k$. We have a root space decomposition $\text{Lie } G = \text{Lie } T \oplus \bigoplus_{\alpha \in \Phi} (\text{Lie } G)_{\alpha}$ where $(\text{Lie } G)_{\alpha} = (\text{Lie } G_{\mathbb{Z}})_{\alpha} \otimes_{\mathbb{Z}} k$, and corresponding root subgroups (defined over \mathbb{Z}) $x_{\alpha} : \mathbb{G}_a \rightarrow G$ such that the tangent map dx_{α} induces an isomorphism between the Lie algebra of the additive group \mathbb{G}_a and $(\text{Lie } G)_{\alpha}$. The functor which associates to any commutative ring A the group $x_{\alpha}(\mathbb{G}_a(A)) = x_{\alpha}(A)$ is a closed subgroup of G denoted by U_{α} , and we have $\text{Lie}(U_{\alpha}) = (\text{Lie } G)_{\alpha}$. Over \mathbb{Z} we denote the corresponding subgroup by $U_{\alpha, \mathbb{Z}}$, over a field k we have $U_{\alpha} = (U_{\alpha, \mathbb{Z}})_k$.

The construction described above in Examples 1.1 and 1.2 makes (in this language) sense over \mathbb{Z} or over any field. As before, let \tilde{G} be the group of the same type but twice the rank, we denote the corresponding Borel subgroup, maximal torus etc. by \tilde{B}, \tilde{T} etc. The construction in the examples above associates to every root $\alpha \in \Phi$ a root $\phi(\alpha)$ in the root system of \tilde{G} . For the \mathbb{Z} -group $G_{\mathbb{Z}}$ we have the subgroup $U_{\mathbb{Z}}^{-}$ and the Lie algebra $\mathfrak{n}_{\mathbb{Z}}^{-} = \bigoplus_{\alpha \in \Phi^+} \mathfrak{n}_{\mathbb{Z}, -\alpha}^{-}$, and we associate to this pair a new pair given by a commutative subgroup $N_{\eta}^{-,a}$ of the \mathbb{Z} -group $\tilde{G}_{\mathbb{Z}}$ and an abelian Lie algebra $\mathfrak{n}_{\mathbb{Z}, \eta}^{-,a}$. The first is the subgroup of the Borel subgroup $\tilde{B}_{\mathbb{Z}} \subset \tilde{G}_{\mathbb{Z}}$ generated by the commuting root subgroups $U_{\phi(\alpha), \mathbb{Z}}$, $\alpha \in \Phi^+$, and the second is the abelian Lie algebra $\mathfrak{n}_{\mathbb{Z}, \eta}^{-,a} = \bigoplus_{\alpha \in \Phi^+} \widetilde{(\text{Lie } G)_{\mathbb{Z}, \phi(\alpha)}}$ given as the sum of root subalgebras.

2. A SPECIAL SCHUBERT VARIETY – THE SL_n -CASE

We want to realize in the situation of Example 1.1 the abelianized representation $V(\lambda)^a$ for $N_{\eta}^{-,a}$ as a Demazure submodule of an irreducible representation for the group SL_{2n} .

2.1. A special Weyl group element. Let \tilde{W} be the Weyl group of $SL_{2n}(\mathbb{C})$, it is the symmetric group \mathcal{S}_{2n} generated by the transpositions s_i , $i = 1, \dots, 2n - 1$. Let $\mathfrak{h} \subset \mathfrak{g} = \mathfrak{sl}_n$ (respectively $\tilde{\mathfrak{h}} \subset \tilde{\mathfrak{g}} = \mathfrak{sl}_{2n}$) be the Cartan subalgebra of traceless complex diagonal matrices. For an element $\alpha \in \mathfrak{h}^*$ and an element $h \in \mathfrak{h}$ we denote by $\langle h, \alpha \rangle$ the evaluation of α in h . Let $\{\epsilon_1, \dots, \epsilon_n\}$ be the elements of the dual vector space \mathfrak{h}^* such that $\langle h, \epsilon_i \rangle$ is the i -th entry in the diagonal matrix $h \in \mathfrak{h}$. We use the same notation $\langle \tilde{h}, \tilde{\alpha} \rangle$ for elements $\tilde{h} \in \tilde{\mathfrak{h}}$ and $\tilde{\alpha} \in \tilde{\mathfrak{h}}^*$, and the linear forms $\{\tilde{\epsilon}_1, \dots, \tilde{\epsilon}_{2n}\}$ in $\tilde{\mathfrak{h}}^*$ are defined as above.

The roots of \mathfrak{g} (resp. $\tilde{\mathfrak{g}}$) are the elements $\alpha_{i,j} := \epsilon_i - \epsilon_j$ (resp. $\tilde{\alpha}_{i,j} = \tilde{\epsilon}_i - \tilde{\epsilon}_j$) for $i \neq j$. We choose as a Borel subalgebra of \mathfrak{g} the subalgebra \mathfrak{b} of upper triangular matrices. The corresponding simple roots are $\alpha_1, \dots, \alpha_{n-1}$ given by $\alpha_i := \alpha_{i, i+1}$. For every root α , we denote by α^\vee its coroot: this is the unique element of \mathfrak{h} such that the reflection $s_\alpha \in \mathfrak{h}^*$ along α acts as $s_\alpha(\lambda) = \lambda - \langle \alpha^\vee, \lambda \rangle \alpha$. Moreover we denote by E_α the corresponding root vector. We denote by $\omega_i = \epsilon_1 + \dots + \epsilon_i$ (resp. $\tilde{\omega}_i = \tilde{\epsilon}_1 + \dots + \tilde{\epsilon}_i$) the i -th fundamental weight of \mathfrak{g} (resp. $\tilde{\mathfrak{g}}$), where $i = 1, 2, \dots, n - 1$ (resp. $i = 1, \dots, 2n - 1$). They are characterized by the property $\langle \alpha_i^\vee, \omega_j \rangle = \delta_{i,j}$.

Definition 2.1. Let $\Psi : \mathfrak{h}^* \rightarrow \tilde{\mathfrak{h}}^*$ be the linear map defined on the weight lattice as follows

$$\Psi\left(\sum_{i=1}^{n-1} a_i \omega_i\right) := \sum_{i=1}^{n-1} a_i \tilde{\omega}_{2i}.$$

Note that Ψ sends dominant weights to dominant weights. For every fundamental weight $\tilde{\omega}_k$, we denote the corresponding parabolic subgroup by $P_{\tilde{\omega}_k}$ and by $\tilde{W}_{\tilde{\omega}_k}$ the corresponding subgroup of \tilde{W} , which is the Weyl group of the semi-simple part of $P_{\tilde{\omega}_k}$. Note that $\tilde{W}_{\tilde{\omega}_k}$ is generated by all the simple transpositions s_i 's but s_k . Let $\rho = \omega_1 + \dots + \omega_{n-1}$. Then $\Psi(\rho) = \tilde{\omega}_2 + \tilde{\omega}_4 + \dots + \tilde{\omega}_{2n-2}$. The parabolic subgroup $Q = P_{\tilde{\omega}_2 + \dots + \tilde{\omega}_{2n-2}}$ which is the stabilizer of $\Psi(\rho)$ will play an important role. The Weyl group of the semisimple part of Q is denoted by \tilde{W}^J .

Definition 2.2. We define in the Weyl group \tilde{W} the element τ as follows:

$$(2) \quad \tau = (s_n s_{n+1} \cdots s_{2n-3} s_{2n-2})(s_{n-1} s_n \cdots s_{2n-4}) \cdots (s_4 s_5 s_6)(s_3 s_4) s_2.$$

It is easy to see that the decomposition is reduced and τ is a minimal length representative in its class in \tilde{W}/\tilde{W}^J . Another description of τ can be given by viewing τ as a permutation of the set $\{1, \dots, 2n\}$:

$$(3) \quad \tau(t) = \begin{cases} n+k & \text{if } t = 2k \\ k & \text{if } t = 2k-1 \end{cases} \quad (k = 1, 2, \dots, n).$$

It follows now immediately from (3):

Lemma 2.3. In the irreducible $SL_{2n}(\mathbb{C})$ -representation $\tilde{V}(\tilde{\omega}_{2i}) = \Lambda^{2i}\mathbb{C}^{2n}$ let v_0 be the highest weight vector $v_0 = e_1 \wedge e_2 \wedge \dots \wedge e_{2i}$. Then (up to sign)

$$\tau(v_0) = v_\tau = e_1 \wedge e_2 \wedge \dots \wedge e_i \wedge e_{n+1} \wedge e_{n+2} \wedge \dots \wedge e_{n+i}.$$

Let $\lambda = b_1 \epsilon_1 + \dots + b_{n-1} \epsilon_{n-1}$, $b_1 \geq \dots \geq b_{n-1} \geq 0$, be a dominant weight for $SL_n(\mathbb{C})$.

Lemma 2.4. $\tau(\Psi(\lambda)) = b_1 \tilde{\epsilon}_1 + \dots + b_{n-1} \tilde{\epsilon}_{n-1} + b_1 \tilde{\epsilon}_{n+1} + \dots + b_{n-1} \tilde{\epsilon}_{2n-1}$.

Proof. Follows directly from Lemma 2.3 above. \square

In Example 1.1 we have introduced a map $\phi : \Phi^+ \rightarrow \tilde{\Phi}^+$ between the positive roots of \mathfrak{sl}_n and the positive roots of \mathfrak{sl}_{2n} . Note that the image of $\alpha = \epsilon_i - \epsilon_j$, $1 \leq i < j \leq n$ is the root $\phi(\alpha) = \tilde{\epsilon}_j - \tilde{\epsilon}_{n+i}$.

Lemma 2.5. *i)* Let λ be a dominant weight for $SL_n(\mathbb{C})$. For a positive SL_{2n} -root $\tilde{\alpha}$ we have $\langle \tilde{\alpha}^\vee, \tau(\Psi(\lambda)) \rangle < 0$ only if the root space of $\tilde{\alpha}$ lies in \mathfrak{n}^{-a} .

ii) Let λ be a dominant SL_n -weight and let $\alpha = \epsilon_i - \epsilon_j$ be a positive SL_n -root, then:

$$\langle \alpha^\vee, \lambda \rangle = -\langle \phi(\alpha)^\vee, \tau(\Psi(\lambda)) \rangle.$$

iii) Let λ be a dominant weight for $SL_n(\mathbb{C})$ and let $\tilde{\alpha} = \tilde{\epsilon}_p - \tilde{\epsilon}_q$ be a positive SL_{2n} -root. Then $E_{\tilde{\alpha}} v_\tau \neq 0$ in $\tilde{V}(\Psi(\lambda))$ only if $\tilde{\alpha}$ is of the form $\tilde{\alpha} = \tilde{\epsilon}_j - \tilde{\epsilon}_{n+i}$, $1 \leq i < j \leq n$ and $\langle (\epsilon_i - \epsilon_j)^\vee, \lambda \rangle > 0$.

Proof. Let $\tilde{\alpha} = \tilde{\epsilon}_i - \tilde{\epsilon}_j$ be a positive root. Lemma 2.4 implies that for $\lambda = b_1 \epsilon_1 + \dots + b_{n-1} \epsilon_{n-1}$ we get

$$\langle \tilde{\alpha}^\vee, \tau(\Psi(\lambda)) \rangle = \begin{cases} b_i - b_j \geq 0 & \text{if } 1 \leq i < j \leq n, \\ b_i - b_{j-n} \geq 0 & \text{if } 1 \leq i \leq n, n+i \leq j \leq 2n, \\ b_i - b_{j-n} \leq 0 & \text{if } 1 \leq i \leq n, n+1 \leq j < n+i, \\ b_{i-n} - b_{j-n} \geq 0 & \text{if } n+1 \leq i < j \leq 2n, \end{cases}$$

which proves the lemma. \square

The decomposition in (2) is reduced, but if we apply τ to a fundamental weight, then it is possible to omit some of the reflections. A simple calculation shows:

Lemma 2.6. Let $\tilde{\omega}_{2i}$ be the $2i$ -th fundamental weight for $SL_{2n}(\mathbb{C})$. Then

$$\tau(\tilde{\omega}_{2i}) = (s_n s_{n+1} \cdots s_{n+i-1}) \cdots (s_{i+2} \cdots s_{2i+1})(s_{i+1} \cdots s_{2i-1} s_{2i})(\tilde{\omega}_{2i}).$$

Let $L(i)$ be the semisimple part of the Levi subgroup of $SL_{2n}(\mathbb{C})$ associated with the simple roots $\tilde{\alpha}_{i+1}, \tilde{\alpha}_{i+2}, \dots, \tilde{\alpha}_{i+n-1}$, denote by $\mathfrak{l}(i)$ the Lie algebra of $L(i)$. Note that $L(i)$ is isomorphic to $SL_n(\mathbb{C})$. Let $\varpi_1, \dots, \varpi_{n-1}$ be the fundamental weights of $L(i)$, the enumeration is such that the simple root $\tilde{\alpha}_{i+j}$ of $L(i) \subseteq SL_{2n}(\mathbb{C})$ corresponds to ϖ_j .

The restriction of $\tilde{\omega}_{2i}$ to $L(i)$ is ϖ_i . Let $W^{L(i)}$ be the Weyl group of $L(i)$, we can identify it with the subgroup of the Weyl group of SL_{2n} generated by the reflections $s_{i+1}, s_{i+2}, \dots, s_{i+n-1}$. Using Lemma 2.6, it is easy to see:

Lemma 2.7. $(s_n s_{n+1} \cdots s_{n+i-1}) \cdots (s_{i+2} \cdots s_{2i+1})(s_{i+1} \cdots s_{2i-1} s_{2i})$ is a reduced decomposition of the longest word of $W^{L(i)}$ modulo the stabilizer $W_{\varpi_i}^{L(i)}$ of ϖ_i in $W^{L(i)}$.

3. THE FUNDAMENTAL REPRESENTATIONS - THE \mathfrak{sl}_n -CASE

We switch now to Lie algebras and hyperalgebras over \mathbb{Z} . Fix a Chevalley basis for the Lie algebra $\mathfrak{g}_{\mathbb{Z}} = \mathfrak{sl}_{n,\mathbb{Z}} \subset \mathfrak{sl}_{n,\mathbb{C}}$:

$$\{f_\alpha, e_\alpha : \alpha \in \Phi^+; h_1, \dots, h_{n-1}\},$$

where $f_\alpha \in \mathfrak{g}_{\mathbb{Z}, -\alpha}$, $e_\alpha \in \mathfrak{g}_{\mathbb{Z}, \alpha}$ and $h_i \in \mathfrak{h}_{\mathbb{Z}}$. For any $m \in \mathbb{Z}_{\geq 1}$, we define the following elements in $U(\mathfrak{g})$:

$$(4) \quad e_\alpha^{(m)} = \frac{e_\alpha^m}{m!} \quad f_\alpha^{(m)} = \frac{f_\alpha^m}{m!} \quad \binom{h_i}{m} = \frac{h_i(h_i-1)\cdots(h_i-m+1)}{m},$$

for $m = 0$ we set: $e_\alpha^{(0)} = f_\alpha^{(0)} = \binom{h_i}{0} = 1$. Recall that the hyperalgebra $U_{\mathbb{Z}}(\mathfrak{sl}_n)$ of $(SL_n)_{\mathbb{Z}}$ is the \mathbb{Z} -subalgebra of the complex enveloping algebra $U(\mathfrak{sl}_n)$ generated by the elements defined in (4). We will use capital letters to denote the Chevalley basis elements for $\mathfrak{sl}_{2n,\mathbb{Z}}$ (e.g. $E_{\tilde{\alpha}}, F_{\tilde{\alpha}}, H_i$) and the generators of the hyperalgebra $U_{\mathbb{Z}}(\mathfrak{sl}_{2n})$ (e.g. $E_{\tilde{\alpha}}^{(m)}, F_{\tilde{\alpha}}^{(m)}, \binom{H_i}{m}$). Similarly, let $U_{\mathbb{Z}}(\tilde{\mathfrak{b}})$ be the subalgebra generated by all $E_{\tilde{\alpha}}^{(m)}$ for $m \geq 0$ and $\tilde{\alpha} > 0$, and all $\binom{H_i}{m}$, $i = 1, \dots, 2n-1$, $m \geq 0$. Denote by $U_{\mathbb{Z}}(\mathfrak{l}(i))$ the hyperalgebra associated with $\mathfrak{l}(i)$, i.e. it is the subalgebra generated by all $F_{\tilde{\alpha}}^{(m)}, E_{\tilde{\alpha}}^{(m)}$, $m \geq 0$, $\tilde{\alpha} > 0$ and a root of the Levi subgroup $L(i)$, and by all $\binom{H_j}{m}$, $j = i+1, \dots, i+n-1$.

Let μ be a dominant integral weight for $SL_{2n}(\mathbb{C})$ and denote by $\tilde{V}(\mu)$ the irreducible $SL_{2n}(\mathbb{C})$ -representation of highest weight μ . Fix a highest weight vector v_μ , the corresponding \mathbb{Z} -form is $\tilde{V}_{\mathbb{Z}}(\mu) = U_{\mathbb{Z}}(\mathfrak{sl}_{2n})v_\mu$. To define the Demazure module $\tilde{V}_{\mathbb{Z}}(\mu)_w$, fix a representative \tilde{w} of w in the simply connected Chevalley group associated with $\mathfrak{sl}_{2n,\mathbb{Z}}$ and set $v_w := \tilde{w}(v_\mu)$. The Demazure module $\tilde{V}_{\mathbb{Z}}(\lambda)_w$ is the cyclic $U_{\mathbb{Z}}(\tilde{\mathfrak{b}})$ -subrepresentation $U_{\mathbb{Z}}(\tilde{\mathfrak{b}}).v_w \subseteq \tilde{V}_{\mathbb{Z}}(\mu)$.

Lemma 3.1. *The Demazure module $\tilde{V}_{\mathbb{Z}}(\ell\tilde{\omega}_{2i})_\tau$ contained in $\tilde{V}_{\mathbb{Z}}(\ell\tilde{\omega}_{2i})$ is the Weyl module $V_{\mathbb{Z}}(\ell\varpi_i)$ of highest weight $\ell\varpi_i$ for $U_{\mathbb{Z}}(\mathfrak{l}(i))$.*

Proof. Consider $\Psi(\ell\omega_i) = \ell\tilde{\omega}_{2i}$ and recall that the restriction of $\tilde{\omega}_{2i}$ to $\mathfrak{l}(i)$ is ϖ_i . So the $U_{\mathbb{Z}}(\mathfrak{l}(i))$ -submodule $U_{\mathbb{Z}}(\mathfrak{l}(i))v_{\ell\tilde{\omega}_{2i}} \subseteq \tilde{V}_{\mathbb{Z}}(\ell\tilde{\omega}_{2i})$ is the Weyl module $V_{\mathbb{Z}}(\ell\varpi_i)$ of highest

weight $\ell\varpi_i$ for $U_{\mathbb{Z}}(\mathfrak{l}(i))$. Let $U_{\mathbb{Z}}(\mathfrak{b}(i))$ be the subalgebra of $U_{\mathbb{Z}}(\mathfrak{l}(i))$ generated by the $E_{\tilde{\alpha}}^{(m)}$, $m \geq 0$, $\tilde{\alpha} > 0$ and a root of $\mathfrak{l}(i)$, and all $\binom{H_j}{m}$, $j = i+1, \dots, i+n-1$.

The Weyl module $V_{\mathbb{Z}}(\ell\varpi_i)$ is a cyclic $U_{\mathbb{Z}}(\mathfrak{b}(i))$ -module and is generated by a lowest weight vector of the form $\tilde{w}_{0,i}(v_{\ell\varpi_i})$, where $\tilde{w}_{0,i}$ is an appropriate representative (in the Chevalley group associated with $\mathfrak{l}(i)$) of the longest element $w_{0,i}$ of the Weyl group $W^{L(i)}$ of $\mathfrak{l}(i)$. Recall that $W^{L(i)}$ can be identified with the subgroup of \tilde{W} generated by $s_{i+1}, \dots, s_{n+i-1}$. Now

$$\tilde{V}_{\mathbb{Z}}(\ell\tilde{\omega}_{2i})_{\tau} = U_{\mathbb{Z}}(\tilde{\mathfrak{b}})v_{\tau} = U_{\mathbb{Z}}(\tilde{\mathfrak{b}})v_{w_{0,i}} = U_{\mathbb{Z}}(\mathfrak{b}(i))v_{w_{0,i}} = V_{\mathbb{Z}}(\ell\varpi_i) \subseteq \tilde{V}_{\mathbb{Z}}(\ell\tilde{\omega}_{2i})$$

by Lemma 2.5, Lemma 2.6 and Lemma 2.7. \square

The previous result implies in particular:

Corollary 3.2. *rank $\tilde{V}_{\mathbb{Z}}(\Psi(\ell\omega_i))_{\tau} = \text{rank } V_{\mathbb{Z}}(\ell\omega_i)$.*

Let $\iota : \mathfrak{sl}_n \rightarrow \mathfrak{sl}_n$ be the Chevalley involution defined by $\iota|_{\mathfrak{h}} = -1$ and ι exchanges e_{α} and $-f_{\alpha}$. It follows that $\iota(\mathfrak{n}_{\mathbb{Z}}^{-}) = \mathfrak{n}_{\mathbb{Z}}^{+}$, and this map extends to an isomorphism of the corresponding hyperalgebras $\iota : U_{\mathbb{Z}}(\mathfrak{n}^{-}) \rightarrow U_{\mathbb{Z}}(\mathfrak{n}^{+})$ and the associated graded versions obtained via the PBW filtration: $\iota : S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{-}) \rightarrow S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{+})$.

Let $\lambda = \sum a_j \omega_j$ be a dominant weight and set $\lambda^* := \sum a_j \omega_{n-j}$. Fix a highest weight vector $v_{\lambda} \in V_{\mathbb{Z}}(\lambda)$ and a lowest weight vector $v_{w_0} \in V_{\mathbb{Z}}(\lambda)$, where w_0 is the longest word in the Weyl group of \mathfrak{sl}_n . We get two possible $S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{-,a})$ -structures on $V_{\mathbb{Z}}(\lambda)$: one uses the PBW filtration on $U_{\mathbb{Z}}(\mathfrak{n}^{-})$ to induce, via the highest weight vector, a PBW filtration on $V_{\mathbb{Z}}(\lambda)$ and passes to the associated graded module. One gets the module $V_{\mathbb{Z}}^a(\lambda)$ discussed before. Now one can do the same also for $U_{\mathbb{Z}}(\mathfrak{n}^{+})$, once replaced the highest weight vector by the lowest weight vector. We denote the cyclic $S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{+})$ -module (generated by the lowest weight vector) by $V_{\mathbb{Z}}^{a,+}(\lambda)$. Now via ι this module also becomes naturally a $S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{-})$ -module.

Lemma 3.3. *The $S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{-})$ -module $V_{\mathbb{Z}}^{a,+}(\lambda)$ is, as $S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{-})$ -module, isomorphic to $V_{\mathbb{Z}}^a(\lambda^*)$.*

Proof. Note that twisting the representation map with the Chevalley involution makes the lowest weight vector (the cyclic generator for the $U(\mathfrak{n}^{+})$ -action) into a cyclic generator for the $U(\mathfrak{n}^{-})$ -action. Recall that the Chevalley involution is equal to -1 on \mathfrak{h} , so after the twist this is now a highest weight vector of weight $\lambda^* = -w_0(\lambda)$, where w_0 is the longest word in W . Since the construction is compatible with the PBW filtrations with respect to the two algebras, the result for the associated graded modules follows immediately. \square

Proposition 3.4. *As $S_{\mathbb{Z}}(\mathfrak{n}^{-,a})$ -modules, $\tilde{V}_{\mathbb{Z}}(\Psi(\ell\omega_i))_{\tau} \simeq V_{\mathbb{Z}}^a(\ell\omega_{n-i})$.*

Proof. Let $\mathfrak{n}_i^{-,a} \subseteq \mathfrak{n}^{-,a}$ be the sum of all root subspaces of roots of the form $\tilde{\epsilon}_k - \tilde{\epsilon}_{n+\ell}$, where $1 \leq \ell \leq i \leq k \leq n$ and $\ell \neq k$. This is a commutative Lie subalgebra, which by Lemma 2.5 has the following property:

$$V_{\mathbb{Z}}(\Psi(\ell\omega_i))_{\tau} = U(\tilde{\mathfrak{b}}).v_{\tau} = U(\mathfrak{n}_i^{-,a}).v_{\tau} = S^{\bullet}(\mathfrak{n}_i^{-,a}).v_{\tau}.$$

Since $\mathfrak{n}^{-,a}$ is commutative, all root vectors in $\mathfrak{n}^{-,a}$ which are not in $\mathfrak{n}_i^{-,a}$ act trivially on $V_{\mathbb{Z}}(\Psi(\ell\omega_i))_{\tau}$.

Another way to describe $\mathfrak{n}_i^{-,a}$ is as the intersection $\mathfrak{n}^{-,a} \cap \mathfrak{l}(i)$. More precisely, this intersection is the nilpotent radical of the maximal parabolic subalgebra of $\mathfrak{l}(i)$ associated with the fundamental weight ϖ_{n-i} . By Lemma 3.1 we know that $V_{\mathbb{Z}}(\Psi(\ell\omega_i))_{\tau} = U(\mathfrak{b}(i)).v_{\tau} \simeq V_{\mathbb{Z}}(\ell\varpi_i)$, since v_{τ} is a lowest weight vector we get in addition

$$V_{\mathbb{Z}}(\Psi(\ell\omega_i))_{\tau} = U(\mathfrak{n}_i^{-,a}).v_{\tau} \simeq V_{\mathbb{Z}}(\ell\varpi_i).$$

Set $\mathfrak{n}^+(i) = \mathfrak{b}(i) \cap \tilde{\mathfrak{n}}^+$, by the isomorphism between $\mathfrak{l}(i)$ and \mathfrak{sl}_n we can identify \mathfrak{m} with $\mathfrak{n}^+ \subset \mathfrak{sl}_n$. Consider the associated PBW filtration on $V_{\mathbb{Z}}(\ell\varpi_i)$ by applying the PBW filtration of $U(\mathfrak{n}^+(i))$ to the lowest weight vector. Recall that after passing to the associated graded algebra $S^{\bullet}(\mathfrak{n}^+(i))$ all root vectors not contained in $\mathfrak{n}_i^{-,a}$ act trivially on $V_{\mathbb{Z}}^{a,+}(\ell\varpi_i)$. Remember that we add a “+” to indicate that this is the associated graded space with respect to the filtration by the nilpotent radical of the fixed Borel subalgebra and not, as usual, of the opposite nilpotent algebra. Since ϖ_i, ϖ_{n-i} are cominusculer, $\mathfrak{n}_i^{-,a}$ is commutative and the PBW filtration on $V_{\mathbb{Z}}(\ell\varpi_i)$ is already a grading. It follows that the action of $\mathfrak{n}_i^{-,a}$ on $V_{\mathbb{Z}}(\ell\varpi_i)$ and $V_{\mathbb{Z}}^{a,+}(\ell\varpi_i)$ are the same and hence the $\mathfrak{n}_i^{-,a}$ action on $V_{\mathbb{Z}}^{a,+}(\ell\varpi_i)$ and $V_{\mathbb{Z}}(\Psi(\ell\omega_i))_{\tau}$ are isomorphic, and hence so are the $\mathfrak{n}^{-,a}$ actions by trivial extension. The proposition follows now by Lemma 3.3. \square

4. THE GENERAL CASE FOR \mathfrak{sl}_n

4.1. We extend Proposition 3.4 to any dominant weight for \mathfrak{sl}_n . Recall that for a dominant weight $\lambda = a_1\omega_1 + \dots + a_{n-1}\omega_{n-1}$ we denote by λ^* the dominant weight given by $\lambda^* = a_{n-1}\omega_1 + \dots + a_1\omega_{n-1}$.

Theorem 4.1. *Let λ be a dominant \mathfrak{sl}_n -weight. The Demazure submodule $\tilde{V}_{\mathbb{Z}}(\Psi(\lambda^*))_{\tau}$ of the $(\mathfrak{sl}_{2n})_{\mathbb{Z}}$ -module $\tilde{V}_{\mathbb{Z}}(\Psi(\lambda^*))$ is, as an $\mathfrak{n}_{\mathbb{Z}}^{-,a}$ -module, isomorphic to $V_{\mathbb{Z}}^a(\lambda)$.*

The proof of Theorem 4.1 will be given in Section 4.7 (the strategy of proof is explained in Section 4.3). We deduce a useful corollary.

Corollary 4.2. *In particular, $V_{\mathbb{Z}}^a(\lambda)$ is free as a \mathbb{Z} -module.*

Proof. (of the corollary) The Demazure module $\tilde{V}_{\mathbb{Z}}(\Psi(\lambda^*))_{\tau}$ is a direct summand of the free \mathbb{Z} -module $\tilde{V}_{\mathbb{Z}}(\Psi(\lambda^*))$ and hence free as \mathbb{Z} -module. \square

4.2. The abelianized module $V_{\mathbb{Z}}^a(\lambda)$ is a cyclic module over the algebra $S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{-,a})$ having as a generator the image of a highest weight vector $v_{\lambda} \in V(\lambda)$ in $V_{\mathbb{Z}}^a(\lambda)$. Hence the module is isomorphic to $S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{-,a})/I_{\mathbb{Z}}(\lambda)$ where $I_{\mathbb{Z}}(\lambda)$ is the annihilator of v_{λ} in $S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{-,a})$.

We have an additional Lie algebra acting on $S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{-,a})$ as well as on $V_{\mathbb{Z}}^a(\lambda)$. Let \mathfrak{b} be the Borel subalgebra of $\mathfrak{g} = (\mathfrak{sl}_n)_{\mathbb{Z}} \otimes \mathbb{C}$ as in Example 1.1, so $\mathfrak{g} = \mathfrak{n}^{-} \oplus \mathfrak{h} \oplus \mathfrak{n}^{+}$. As free \mathbb{Z} -modules, $U_{\mathbb{Z}}(\mathfrak{n}^{-}) \simeq U_{\mathbb{Z}}(\mathfrak{g})/U_{\mathbb{Z}}^+(\mathfrak{h} + \mathfrak{n}^{+})$, so that the adjoint action of $U_{\mathbb{Z}}(\mathfrak{b})$ on $U_{\mathbb{Z}}(\mathfrak{g})$ induces a structure of $U_{\mathbb{Z}}(\mathfrak{b})$ -module on $U_{\mathbb{Z}}(\mathfrak{n}^{-})$ and hence on $S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{-,a})$. This action is compatible with the induced $U_{\mathbb{Z}}(\mathfrak{b})$ -action on $V_{\mathbb{Z}}^a(\lambda)$ (cf.[FFL3, Prop. 2.3.]). Recall that for a positive root α we have denoted by f_{α} the corresponding fixed Chevalley basis element in $(\mathfrak{sl}_n)_{-\alpha, \mathbb{Z}}$. Using the presentation of Demazure modules in terms of generators and relations by Joseph,

Mathieu and Polo (cf. [M], Lemme 26), we get as a consequence of the proof of Theorem 4.1 the following description of the ideal $I_{\mathbb{Z}}(\lambda)$ (cf. [FFL1], [FFL3]):

Corollary 4.3. *The abelianized module $V_{\mathbb{Z}}^a(\lambda)$ is isomorphic to $S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{-,a})/I_{\mathbb{Z}}(\lambda)$ (as a cyclic $S_{\mathbb{Z}}(\mathfrak{n}^{-,a})$ -module), where*

$$I_{\mathbb{Z}}(\lambda) = S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{-,a})(U_{\mathbb{Z}}(\mathfrak{n}^+) \circ \text{span}\{f_{\alpha}^{(\langle \tilde{\alpha}^{\vee}, \lambda \rangle + m)}; m \geq 1, \alpha > 0\}) \subseteq S_{\mathbb{Z}}(\mathfrak{n}^{-,a}).$$

4.3. The proof of Theorem 4.1 will be given in section 4.7, it needs some preparation. The strategy of the proof is summarized by the following diagram of $S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{-,a})$ -modules. For a dominant weight $\lambda = a_1\omega_1 + \dots + a_{n-1}\omega_{n-1}$ (so $\lambda^* = a_{n-1}\omega_1 + \dots + a_1\omega_{n-1}$), we get the following natural maps, the details are described below:

$$\begin{array}{ccccc} S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{-,a})/I_{\mathbb{Z}}(\lambda^*) & \xrightarrow{\simeq} & V_{\mathbb{Z}}^a(\lambda^*) & \xrightarrow{b} & V_{\mathbb{Z}}^a(a_1\omega_1^*) \otimes \cdots \otimes V_{\mathbb{Z}}^a(a_{n-1}\omega_{n-1}^*) \\ \uparrow f & & \downarrow a & & \downarrow c \\ S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{-,a})/M_{\mathbb{Z}}(\lambda^*) & \xrightarrow{\simeq} & \tilde{V}_{\mathbb{Z}}(\Psi(\lambda))_{\tau} & \xrightarrow{d} & \tilde{V}_{\mathbb{Z}}(a_1\Psi(\omega_1))_{\tau} \otimes \cdots \otimes \tilde{V}_{\mathbb{Z}}(a_{n-1}\Psi(\omega_{n-1}))_{\tau}. \end{array}$$

Let us describe the diagram above and the strategy of the proof. We recall that given a tensor product of cyclic $S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{-,a})$ -modules, the Cartan component of the tensor product is, by definition, the cyclic $S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{-,a})$ -submodule generated by the tensor product of the cyclic generators. Further, recall that the isomorphism $V_{\mathbb{Z}}^a(\ell\omega_j^*) \simeq \tilde{V}_{\mathbb{Z}}(\ell\Psi(\omega_j))_{\tau}$ sends the highest weight vector $v_{\ell\omega_j^*}$ to the extremal weight vector $v_{\tau(\ell\Psi(\omega_j))}$ and uses the Chevalley involution. The maps above are defined as follows:

- b is induced by the compatibility of the PBW filtration with the tensor product, and it is surjective onto the Cartan component of this tensor product.
- $I_{\mathbb{Z}}(\lambda^*)$ is the annihilator in $S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{-,a})$ of the image of the highest weight vector v_{λ^*} in $V_{\mathbb{Z}}^a(\lambda^*)$ and h is the corresponding quotient map.
- c is the isomorphism given by Proposition 3.4.
- d is the isomorphism onto the Cartan component of the tensor product. The fact that this is an isomorphism follows by standard monomial theory [LMS] or Frobenius splitting [R].
- a equals $c \circ b$ after identifying $\tilde{V}_{\mathbb{Z}}(\Psi(\lambda))_{\tau}$ with its image under d .
- $M_{\mathbb{Z}}(\lambda^*)$ is the annihilator in $S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{-,a})$ of the extremal weight vector $v_{\tau(\Psi(\lambda))}$ in $\tilde{V}_{\mathbb{Z}}(\Psi(\lambda))_{\tau}$ and g is the corresponding quotient map.
- f is going to be constructed in the proof.

In order to finish the proof we will show that $M_{\mathbb{Z}}(\lambda^*) \subseteq I_{\mathbb{Z}}(\lambda^*)$, and the inclusion induces the surjective map f which in turn shows that the map a is an isomorphism.

4.4. We first determine $M_{\mathbb{Z}}(\lambda^*)$. By [M, Lemme 26], the Demazure module $\tilde{V}_{\mathbb{Z}}(\Psi(\lambda))_{\tau}$ is isomorphic to the algebra $U_{\mathbb{Z}}(\tilde{\mathfrak{n}}^+)$ modulo the left ideal $\tilde{I}_{\mathbb{Z}}(\tau\Psi(\lambda))$ generated by the $E_{\tilde{\alpha}}^{(m)}$ for all $m \geq 1$ if $\langle \tilde{\alpha}^{\vee}, \tau\Psi(\lambda) \rangle \geq 0$, and $E_{\tilde{\alpha}}^{(-\langle \tilde{\alpha}^{\vee}, \tau\Psi(\lambda) \rangle + m)}$ for all $m \geq 1$ otherwise.

Proof. Let \bar{m} be an element of $M_{\mathbb{Z}}(\lambda^*)$ and choose a representative m in $\tilde{I}_{\mathbb{Z}}(\tau\Psi(\lambda))$. Since we are free to choose a representative modulo $I_{\mathbb{Z}}(\infty)$, we may assume (see section 4.4) that m is a sum of monomials of the form $rE_{\tilde{\alpha}}^{(\ell)}$ where $\ell = -\langle \tilde{\alpha}^\vee, \tau\Psi(\lambda) \rangle + k$ for some $k \geq 1$ and $\tilde{\alpha}$ of first type, and r is a monomial in the $E_{\tilde{\beta}}^{(q)}$, $q \geq 0$ and $\tilde{\beta}$ of first, second or third type.

If $\tilde{\gamma}$ is a root of third type and $\tilde{\beta}$ is any other positive root, then $[E_{\tilde{\gamma}}, E_{\tilde{\beta}}] = cE_{\tilde{\gamma}'}$, where $c \in \mathbb{Z}$ and either $c = 0$ or $\tilde{\gamma}'$ of third type. So if r has a factor $E_{\tilde{\gamma}}^{(p)}$, $p > 0$ and $\tilde{\gamma}$ a root of third type, then we can rewrite the monomial $rE_{\tilde{\alpha}}^{(\ell)}$ as a sum of monomials of the form $r'E_{\tilde{\gamma}}^{(p')}$, $p' > 0$. Since this sum is an element in $I_{\mathbb{Z}}(\infty)$, without loss of generality we will assume in the following that r has only factors of the form $E_{\tilde{\beta}}^{(\ell)}$, $\tilde{\beta}$ of first or second type.

If $\tilde{\gamma}$ is of second type and $\tilde{\beta}$ is of first type, then $[E_{\tilde{\gamma}}, E_{\tilde{\beta}}] = cE_{\tilde{\gamma}'}$, where $c \in \mathbb{Z}$ and either $c = 0$ or $\tilde{\gamma}'$ is of first or third type. So after reordering the factors we can assume without loss of generality in the following that $rE_{\tilde{\alpha}}^{(\ell)}$ is of the form $r = r_1r_2$, where r_1 is a monomial in the $E_{\tilde{\beta}}^{(\ell)}$, $\tilde{\beta}$ of first type, and r_2 is a monomial in the $E_{\tilde{\gamma}}^{(\ell)}$, $\tilde{\gamma}$ of second type.

Recall that for $\tilde{\gamma}$ of second type we have

$$\begin{aligned} E_{\tilde{\gamma}}E_{\tilde{\beta}_1} \cdots E_{\tilde{\beta}_m} &\equiv \sum_{i=1}^m E_{\tilde{\beta}_1} \cdots E_{\tilde{\beta}_{i-1}} (E_{\tilde{\gamma}}E_{\tilde{\beta}_i} - E_{\tilde{\beta}_i}E_{\tilde{\gamma}})E_{\tilde{\beta}_{i+1}} \cdots E_{\tilde{\beta}_m} \text{ mod } I_{\mathbb{Z}}(\infty) \\ &\equiv \sum_{i=1}^m E_{\tilde{\beta}_1} \cdots E_{\tilde{\beta}_{i-1}} (ad(E_{\tilde{\gamma}})(E_{\tilde{\beta}_i}))E_{\tilde{\beta}_{i+1}} \cdots E_{\tilde{\beta}_m} \text{ mod } I_{\mathbb{Z}}(\infty) \\ &\equiv ad(E_{\tilde{\gamma}})(E_{\tilde{\beta}_1} \cdots E_{\tilde{\beta}_m}) \text{ mod } I_{\mathbb{Z}}(\infty). \end{aligned}$$

An appropriate reformulation of the equality above holds also for the divided powers of the root vectors. It follows that $r_2E_{\tilde{\alpha}}^{(\ell)} \in m_{\mathbb{Z}}(\lambda^*)$ and hence $rE_{\tilde{\alpha}}^{(\ell)} = r_1r_2E_{\tilde{\alpha}}^{(\ell)} \in S_{\mathbb{Z}}^\bullet(\mathfrak{n}^{-a})m_{\mathbb{Z}}(\lambda^*)$, which implies that $M_{\mathbb{Z}}(\lambda^*)$ is generated by $m_{\mathbb{Z}}(\lambda^*)$ as a left $S_{\mathbb{Z}}^\bullet(\mathfrak{n}^{-a})$ -module. \square

4.6. To compare $M_{\mathbb{Z}}(\lambda^*)$ with $I_{\mathbb{Z}}(\lambda^*)$, we need a variant of the description of $m_{\mathbb{Z}}(\lambda^*)$. Let $\Delta(\mathfrak{b}_{\mathbb{Z}}) \subset \mathfrak{b}_{\mathbb{Z}}^1 \oplus \mathfrak{b}_{\mathbb{Z}}^2$ be the Lie subalgebra obtained as a diagonally embedded copy of $\mathfrak{b}_{\mathbb{Z}}$. Let $U_{\mathbb{Z}}(\Delta(\mathfrak{b}_{\mathbb{Z}})) \subset U_{\mathbb{Z}}((\mathfrak{b}_{\mathbb{Z}}^1) \oplus (\mathfrak{b}_{\mathbb{Z}}^2))$ be its hyperalgebra.

Lemma 4.5. $m_{\mathbb{Z}}(\lambda^*) = \langle U_{\mathbb{Z}}(\Delta(\mathfrak{b}_{\mathbb{Z}})) \circ E_{\tilde{\alpha}}^{(-\langle \tilde{\alpha}^\vee, \tau\Psi(\lambda) \rangle + m)} \mid \tilde{\alpha} \text{ of first type, } m \geq 1 \rangle_{\mathbb{Z}}$.

Proof. We assume first that k is an algebraically closed field of arbitrary characteristic. Let B be the subgroup of upper triangular invertible matrices in $SL_n(k)$, so $\text{Lie } B = \mathfrak{b}$. Let $B^1 \times B^2 \subset SL_{2n}(k)$ be the subgroup with Lie algebra $\mathfrak{b}^1 \oplus \mathfrak{b}^2$ and denote by $\Delta(B) \subset B^1 \times B^2$ the diagonally embedded group isomorphic to B .

Let \mathfrak{q} be the sum of the SL_{2n} -root spaces corresponding to roots of second or third type. Then $\tilde{\mathfrak{n}}^+ = \mathfrak{n}^{-a} \oplus \mathfrak{q}$ and we identify \mathfrak{n}^{-a} with $\tilde{\mathfrak{n}}^+/\mathfrak{q}$. The adjoint action of $B^1 \times B^2$ on \mathfrak{sl}_{2n} admits $\tilde{\mathfrak{n}}^+$ as well as \mathfrak{q} as submodules, so we get an induced $B^1 \times B^2$ -action on $\mathfrak{n}^{-a} = \tilde{\mathfrak{n}}^+/\mathfrak{q}$. This action naturally extends to the commutative hyperalgebra $S_k^\bullet(\mathfrak{n}^{-a})$.

If we replace the group action of $B^1 \times B^2$ by the induced action of the hyperalgebra $U_k(\mathfrak{b}^1 \oplus \mathfrak{b}^2)$ of the group, then we get the action of $U_k(\mathfrak{b}^1 \oplus \mathfrak{b}^2)$ on $U_k(\tilde{\mathfrak{n}}^+)$ respectively on $S_k^\bullet(\mathfrak{n}^{-a})$ discussed above, similarly for the action of $\Delta(B)$ and its hyperalgebra $U_k(\Delta(\mathfrak{b}))$.

Recall that for a root $\tilde{\alpha}$ of type 1 we have

$$U_k(\mathfrak{b}^1 \oplus \mathfrak{b}^2) \circ E_{\tilde{\alpha}}^{(m)} = \langle \text{Ad}((b^1, b^2)) \circ (E_{\tilde{\alpha}})^{(m)} \mid (b_1, b_2) \in B^1 \times B^2 \rangle,$$

i.e. the smallest $U_k(\mathfrak{b}^1 \oplus \mathfrak{b}^2)$ stable subspace containing $(E_{\tilde{\alpha}})^{(m)}$ is the linear span of the $B^1 \times B^2$ -orbit. The same holds in the other case, so we have:

$$U_k(\Delta(\mathfrak{b})) \circ E_{\tilde{\alpha}}^{(m)} = \langle \text{Ad}((b, b)) \circ (E_{\tilde{\alpha}})^{(m)} \mid b \in B \rangle.$$

Let \mathfrak{d} be the sum of the SL_{2n} -root spaces corresponding to roots of first or third type and let \mathfrak{d}^3 be just the sum of the root spaces corresponding to roots of third type, so $\mathfrak{d} = \mathfrak{n}^{-,a} \oplus \mathfrak{d}^3$. We identify $\mathfrak{d} \subset \mathfrak{sl}_{2n}$ with $M_n(k)$, formally this can be done by the map

$$\chi : \mathfrak{d} \rightarrow M_n(k), \quad \tilde{A} = \begin{pmatrix} 0 & A \\ 0 & 0 \end{pmatrix} \mapsto A,$$

where A is a $n \times n$ matrix. In the following we simplify the notation and omit the map χ . We freely identify \mathfrak{d} with $M_n(k)$, so we denote by A the $n \times n$ matrix as well as the $2n \times 2n$ -matrix $\tilde{A} \in \mathfrak{d}$. Note that for $(b_1, b_2) \in B^1 \times B^2$ we get

$$\text{Ad}((b_1, b_2)) \circ \tilde{A} = \begin{pmatrix} b_1 & 0 \\ 0 & b_2 \end{pmatrix} \begin{pmatrix} 0 & A \\ 0 & 0 \end{pmatrix} \begin{pmatrix} b_1^{-1} & 0 \\ 0 & b_2^{-1} \end{pmatrix} = \begin{pmatrix} 0 & b_1 A b_2^{-1} \\ 0 & 0 \end{pmatrix},$$

we just write $\text{Ad}((b_1, b_2)) \circ (A) = b_1 A b_2^{-1}$ and $\text{Ad}((b, b)) \circ (A) = b A b^{-1}$. Recall that χ is just a vector space isomorphism. If we equip in addition $M_n(k)$ with the trivial Lie bracket, then this becomes also a Lie algebra homomorphism. In this sense we identify also the (commutative) Lie subalgebras $\mathfrak{n}^{-,a}$ and \mathfrak{d}^3 with subalgebras of M_n . An elementary calculation shows how the $B^1 \times B^2$ -orbit through $E_{\tilde{\alpha}} = E_{\epsilon_n - \epsilon_{n+1}}$ breaks up into $\Delta(B)$ -orbits, recall that we identify \mathfrak{d} with $M_n(k)$ and $E_{\epsilon_n - \epsilon_{n+1}} \in \mathfrak{d}$ corresponds to $E_{n,1}$:

$$\begin{aligned} \{\text{Ad}((b_1, b_2)) \circ (E_{\tilde{\alpha}}) \mid (b_1, b_2) \in B^1 \times B^2\} &= \{b_1 E_{n,1} b_2^{-1} \mid b_1, b_2 \in B\} \\ &= \bigcup_{\lambda \in k} \{b(E_{n,1} + \lambda E_{1,1}) b^{-1} \mid b \in B\} \\ &\subset M_n(k). \end{aligned}$$

We conclude for the linear span:

$$\begin{aligned} U_k(\mathfrak{b}^1 \oplus \mathfrak{b}^2) \circ E_{\tilde{\alpha}}^{(m)} &= \langle \text{Ad}((b_1, b_2)) \circ (E_{\tilde{\alpha}})^{(m)} \mid (b_1, b_2) \in B^1 \times B^2 \rangle \\ &= \langle (\text{Ad}((b_1, b_2)) \circ (E_{\tilde{\alpha}}))^{(m)} \mid (b_1, b_2) \in B^1 \times B^2 \rangle \\ &= \langle (b_1 E_{n,1} b_2^{-1})^{(m)} \mid b_1, b_2 \in B \rangle \\ &= \langle \bigcup_{\lambda \in k} \{b(E_{n,1} + \lambda E_{1,1})^{(m)} b^{-1} \mid b \in B\} \rangle. \end{aligned}$$

Let $I(\mathfrak{d}^3) \subset U_k(\mathfrak{d})$ be the left ideal in the hyperalgebra generated by \mathfrak{d}^3 . It follows:

$$\begin{aligned} U_k(\mathfrak{b}^1 \oplus \mathfrak{b}^2) \circ E_{\tilde{\alpha}}^{(m)} \bmod I(\mathfrak{d}^3) &= \langle \bigcup_{\lambda \in k} \{b(E_{n,1} + \lambda E_{1,1}) b^{-1}\}^{(m)} \mid b \in B \rangle \bmod I(\mathfrak{d}^3) \\ &= \langle (b E_{n,1} b^{-1})^{(m)} \mid b \in B \rangle \bmod I(\mathfrak{d}^3) \end{aligned}$$

because the $E_{1,1}^{(\ell)}$, $\ell \geq 1$, lie in the $\Delta(B)$ -stable ideal $I(\mathfrak{d}^3)$. It follows:

$$\begin{aligned} U_k(\mathfrak{b}^1 \oplus \mathfrak{b}^2) \circ E_{\tilde{\alpha}}^{(m)} \bmod I(\mathfrak{d}^3) &= \langle \text{Ad}((b, b))(E_{\tilde{\alpha}})^{(m)} \mid b \in B \rangle \bmod I(\mathfrak{d}^3) \\ &= U_k(\Delta(\mathfrak{b})) \circ E_{\tilde{\alpha}}^{(m)} \bmod I(\mathfrak{d}^3). \end{aligned}$$

Since $I(\mathfrak{d}^3) \subset \tilde{I}_k(\infty)$, the equation holds in $S^\bullet(\mathfrak{n}^{-,a}) = U_k(\tilde{\mathfrak{n}}^+)/\tilde{I}(\infty)$ too and hence

$$U_k(\mathfrak{b}^1 \oplus \mathfrak{b}^2) \circ E_{\tilde{\alpha}}^{(m)} = U_k(\Delta(\mathfrak{b})) \circ E_{\tilde{\alpha}}^{(m)}$$

in $S^\bullet(\mathfrak{n}^{-,a})$. It is now easy to see that the same arguments prove the equality for all $E_{\tilde{\alpha}}^{(m)}$, $\tilde{\alpha}$ of first type and $m \geq 1$. We have clearly

$$U_{\mathbb{Z}}(\mathfrak{b}^1 \oplus \mathfrak{b}^2) \circ E_{\tilde{\alpha}}^{(m)} \supseteq U_{\mathbb{Z}}(\Delta(\mathfrak{b})) \circ E_{\tilde{\alpha}}^{(m)}.$$

Since we have equality after base change for fields of arbitrary characteristics, the equality of the modules holds also over \mathbb{Z} . In particular, the following equality holds in $S_{\mathbb{Z}}^\bullet(\mathfrak{n}^{-,a})$:

$$\begin{aligned} m_{\mathbb{Z}}(\lambda^*) &= \langle U_{\mathbb{Z}}(\mathfrak{b}_{\mathbb{Z}}^1 \oplus \mathfrak{b}_{\mathbb{Z}}^2) \circ E_{\tilde{\alpha}}^{(-\langle \tilde{\alpha}^\vee, \tau\Psi(\lambda) \rangle + m)} \mid \tilde{\alpha} \text{ of first type and } m \geq 1 \rangle_{\mathbb{Z}} \\ &= \langle U_{\mathbb{Z}}(\Delta(\mathfrak{b}_{\mathbb{Z}})) \circ E_{\tilde{\alpha}}^{(-\langle \tilde{\alpha}^\vee, \tau\Psi(\lambda) \rangle + m)} \mid \tilde{\alpha} \text{ of first type and } m \geq 1 \rangle_{\mathbb{Z}}. \end{aligned}$$

□

4.7. Proof of Theorem 4.1. Recall the identification of the abelianized version of $\mathfrak{n}^- \subset \mathfrak{sl}_n$ with $\mathfrak{n}^{-,a} \subset \mathfrak{sl}_{2n}$, which sends the image of a Chevalley generator f_α to $E_{\phi(\alpha)}$. By Lemma 2.5 (see also Lemma 3.3 for the twist $\lambda \leftrightarrow \lambda^*$) the elements $E_{\tilde{\alpha}}^{(-\langle \tilde{\alpha}^\vee, \tau\Psi(\lambda) \rangle + m)}$, where $\tilde{\alpha}$ is of first type and $m \geq 1$ are elements of $I_{\mathbb{Z}}(\lambda^*)$. Now the $U_{\mathbb{Z}}(\mathfrak{b})$ -module structure on $S_{\mathbb{Z}}^\bullet(\mathfrak{n}^{-,a})$ described in section 4.2 is the same as the one described above, so it follows that $m_{\mathbb{Z}}(\lambda^*) \subset I_{\mathbb{Z}}(\lambda^*)$ and hence $M_{\mathbb{Z}}(\lambda^*) \subset I_{\mathbb{Z}}(\lambda^*)$, which, as explained in Section 4.3, finishes the proof of the theorem. □

4.8. Let ρ be the sum of all fundamental weights for SL_n and set $\tilde{\rho} = \Psi(\rho)$. Let $Q_{\mathbb{Z}} \subset (SL_{2n})_{\mathbb{Z}}$ be the corresponding parabolic \mathbb{Z} -subgroup. Recall that $(N^{-,a})_{\mathbb{Z}}$ is a commutative subgroup of the Borel subgroup $\tilde{B}_{\mathbb{Z}}$. For any SL_{2n} -root $\tilde{\alpha}$ let $U_{\mathbb{Z},\tilde{\alpha}}$ be the associated root subgroup.

Lemma 4.6. *The orbit $\tilde{B}_{\mathbb{Z}}.\tau \subset (SL_{2n})_{\mathbb{Z}}/Q_{\mathbb{Z}}$ is equal to $N^{-,a}.\tau$, and the map $N^{-,a} \rightarrow N^{-,a}.\tau$, $u \mapsto u\tau$, is a bijection.*

Proof. We have $\tilde{B}_{\mathbb{Z}}.\tau = \prod_{\tilde{\alpha} > 0} U_{\mathbb{Z},\tilde{\alpha}}.\tau$, and the map $\prod_{\tilde{\alpha} \in \Gamma} U_{\mathbb{Z},\tilde{\alpha}} \rightarrow \prod_{\tilde{\alpha} \in \Gamma} U_{\mathbb{Z},\tilde{\alpha}}.\tau$ is a bijection, where Γ is the set of all positive roots of SL_{2n} such that $\tau^{-1}(\tilde{\alpha}) < 0$ and $\tau^{-1}(\tilde{\alpha})$ is not an element of the root system of $Q_{\mathbb{Z}}$. Now this condition is fulfilled if and only if $\langle \tau^{-1}(\tilde{\alpha}^\vee), \tilde{\rho} \rangle < 0$, or, equivalently, $\langle \tilde{\alpha}^\vee, \tau(\tilde{\rho}) \rangle < 0$. By Lemma 2.4 this is only possible if $\tilde{\alpha} = \tilde{\epsilon}_i - \tilde{\epsilon}_j$, $i < j$, is such that $1 \leq i \leq n$, $n+1 \leq j \leq 2n$ and $i > j - n$. But this implies that the root subgroup $U_{\mathbb{Z},\tilde{\alpha}}$ is a subgroup of $N^{-,a}$, and all root subgroups of $(SL_{2n})_{\mathbb{Z}}$ contained in $N^{-,a}$ satisfy this condition. It follows that $N^{-,a}$ is the product of all root subgroups corresponding to positive roots of SL_{2n} in Γ . □

Recall that the degenerate flag scheme $\mathcal{F}\ell(\lambda)_{\mathbb{Z}}^a$ is the closure of the $N_{\mathbb{Z}}^{-,a}$ -orbit through the highest weight vector in $\mathbb{P}(V_{\mathbb{Z}}^a(\lambda))$.

Theorem 4.7. *Let λ be a dominant weight for SL_n . The Schubert scheme $X_{\mathbb{Z}}(\tau) \subset \mathbb{P}(\tilde{V}_{\mathbb{Z}}(\Psi(\lambda))_{\tau})$ is isomorphic to the degenerate partial flag scheme $\mathcal{F}\ell(\lambda^*)_{\mathbb{Z}}^a$ for $(SL_n)_{\mathbb{Z}}$, and this isomorphism induces a module isomorphism $H^0(X_{\mathbb{Z}}(\tau), \mathcal{L}_{\Psi(\lambda)}) \simeq (V_{\mathbb{Z}}^a(\lambda^*))^*$.*

Proof. We consider only the case where λ is regular, the arguments in the general case are similar. With respect to the isomorphism in Lemma 4.6, the orbit

$$\tilde{B}_{\mathbb{Z}}.\tau \subset (SL_{2n})_{\mathbb{Z}}/(\tilde{P}_{\lambda})_{\mathbb{Z}} \hookrightarrow \mathbb{P}(\tilde{V}(\Psi(\lambda))),$$

through the extremal weight vector, which is the same as the $N^{-,a}$ -orbit, is mapped onto the $N_{\mathbb{Z}}^{-,a}$ -orbit through the highest weight vector in $\mathbb{P}(V_{\mathbb{Z}}^a(\lambda^*))$. By definition, the Schubert scheme $X_{\mathbb{Z}}(\tau)$ is the closure of the orbit $\tilde{B}_{\mathbb{Z}}.\tau$ and the degenerate flag scheme $\mathcal{F}\ell(\lambda^*)_{\mathbb{Z}}^a$ is the closure of the $N_{\mathbb{Z}}^{-,a}$ -orbit. It follows that the module isomorphism induces an isomorphism between the Schubert scheme $X_{\mathbb{Z}}(\tau) \subseteq \mathbb{P}(\tilde{V}_{\mathbb{Z}}(\Psi(\lambda))_{\tau})$ and the degenerate flag scheme $\mathcal{F}\ell(\lambda^*)_{\mathbb{Z}}^a$ in $\mathbb{P}(V_{\mathbb{Z}}^a(\lambda))$. Hence we get induced isomorphisms

$$H^0(X_{\mathbb{Z}}(\tau), \mathcal{L}_{\Psi(\lambda)}) \simeq (\tilde{V}_{\mathbb{Z}}(\Psi(\lambda))_{\tau})^* \simeq (V_{\mathbb{Z}}^a(\lambda^*))^*$$

for the dual modules. □

Let k be an algebraically closed field of arbitrary characteristic and denote by $V_k(\lambda) = V_{\mathbb{Z}}(\lambda) \otimes_{\mathbb{Z}} k$, $U_k(\mathfrak{sl}_n) = U_{\mathbb{Z}}(\mathfrak{sl}_n) \otimes_{\mathbb{Z}} k$, $U_k(\mathfrak{n}^-) = U_{\mathbb{Z}}(\mathfrak{n}^-) \otimes_{\mathbb{Z}} k$ etc. the objects obtained by base change. The PBW filtration

$$V_k(\lambda)_{\ell} = \langle Y_1^{(m_1)} \cdots Y_N^{(m_N)} v_{\lambda} \mid m_1 + \cdots + m_N \leq \ell, Y_1, \dots, Y_N \in \mathfrak{n}_k^- \rangle$$

and the associated graded space $V_k^a(\lambda)$ is defined in the same way as before, and by Corollary 4.2 we have $V_k(\lambda)_{\ell} = V_{\mathbb{Z}}(\lambda)_{\ell} \otimes_{\mathbb{Z}} k$ and $V_k^a(\lambda) = V_{\mathbb{Z}}^a(\lambda) \otimes_{\mathbb{Z}} k$. The group $N_k^{-,a}$ acts on the abelianized representation $V_k^a(\lambda)$, and the degenerate flag variety $\mathcal{F}\ell(\lambda)_k^a$ is the closure of the $N_k^{-,a}$ -orbit through the highest weight vector in $\mathbb{P}(V_k^a(\lambda))$.

Now by the results of [M, MR, R, RR] one knows that for Demazure modules we have $\tilde{V}_k(\Psi(\lambda))_{\tau} = \tilde{V}_{\mathbb{Z}}(\Psi(\lambda))_{\tau} \otimes_{\mathbb{Z}} k$, $X_k(\tau) = X_{\mathbb{Z}}(\tau) \otimes_{\mathbb{Z}} k$, etc., and that the Schubert varieties are Frobenius split, projectively normal and have rational singularities. It follows that $V_k^a(\lambda^*) = \tilde{V}_k(\Psi(\lambda))_{\tau}$ and $\mathcal{F}\ell(\lambda^*)_k^a = X_k(\tau)$, so the degenerate flag variety has in this case the same nice geometric properties as the Schubert variety. For a dominant SL_n -weight $\lambda = \sum_{i=1}^{n-1} a_i \omega_i$ let the support $\text{supp } \lambda$ of λ be the set $\{i \mid 1 \leq i \leq n-1, a_i \neq 0\}$.

Corollary 4.8. *The degenerate partial flag variety $\mathcal{F}\ell(\lambda)_k^a$ depends only on $\text{supp } \lambda$, it is a projectively normal variety, Frobenius split, with rational singularities.*

Remark 4.9. In [FF] the authors construct resolutions of the degenerate flag varieties given by towers of \mathbb{P}^1 -fibrations. The steps of the successive fibrations are indexed by the set of positive roots, which had been totally reordered. In fact their varieties are Bott-Samelson varieties (cf [CL, Appendix]) and such an order (which actually should be thought of as an order on the set of negative roots) is now natural since it corresponds to the subsequent steps of the Bott-Samelson variety indexed by the reduced expression (2) of τ , under the identification of $-\alpha_{i,j}$ with $\tilde{\alpha}_{j,i+n}$.

5. A SPECIAL SCHUBERT VARIETY – THE Sp_{2m} -CASE

As for the SL_n -case, we want to realize for Example 1.2 the abelianized representation $V_{\mathbb{Z}}(\lambda)^a$ for $N_{\mathbb{Z}}^{-,a}$ as a Demazure submodule in an irreducible representation for the larger group $Sp_{2(2m)}$.

5.1. A special Weyl group element. Let us keep the same notation as in the previous sections and denote by $\mathfrak{h} \subset \mathfrak{sp}_{2m}$ (respectively $\tilde{\mathfrak{h}} \subset \mathfrak{sp}_{2(2m)}$) the Cartan subalgebra of traceless complex diagonal matrices and by $\mathfrak{b} \subset \mathfrak{sp}_{2m}$ (respectively $\tilde{\mathfrak{b}} \subset \mathfrak{sp}_{2(2m)}$) the Borel subalgebra of traceless complex upper triangular matrices. Let $\{\epsilon_1, \dots, \epsilon_{2m}\}$ (resp. $\{\tilde{\epsilon}_1, \dots, \tilde{\epsilon}_{2(2m)}\}$) be a basis of the dual vector space \mathfrak{h}^* (resp. $\tilde{\mathfrak{h}}^*$). The choice of Cartan and Borel subalgebras we made determines the following set of positive roots for Sp_{2m} :

$$\alpha_{i,j} := \begin{cases} \epsilon_i - \epsilon_j & 1 \leq i < j \leq m, \\ \epsilon_i + \epsilon_{j-m} & 1 \leq i \leq m < j, i + j \leq 2m, \end{cases}$$

where the simple roots are $\{\alpha_i := \alpha_{i,i+1} \mid 1 \leq i \leq m-1\} \cup \{\alpha_m := 2\epsilon_m\}$. We will write $\tilde{\alpha}_{i,j}$ for the $Sp_{2(2m)}$ -roots. The Weyl group of $Sp_{2(2m)}$ is denoted \tilde{W} . This is the group of linear transformations of $\tilde{\mathfrak{h}}^*$ generated by the elements $\{r_i \mid 1 \leq i \leq 2m\}$, where r_i denotes the reflection with respect to the simple root $\tilde{\alpha}_i$.

Definition 5.1. We define in \tilde{W} a very special element:

$$\bar{\tau} = (r_{2m} \cdots r_{m+1}) \cdots (r_{2m} r_{2m-1} r_{2m-2}) (r_{2m} r_{2m-1}) r_{2m} (r_m \cdots r_{2m-2}) \cdots (r_4 r_5 r_6) (r_3 r_4) r_2.$$

Any element of the Weyl group \tilde{W} of $Sp_{2(2m)}$ can be identified with an element of the symmetric group on $4m$ letters \mathcal{S}_{4m} , via $r_i = s_i s_{4m-i}$, for $1 \leq i \leq 2m-1$, and $r_n = s_{2m}$ (where, as usual, s_i denotes the transposition exchanging i and $i+1$) and it acts on the basis $\{e_i \mid i = 1, \dots, 4m\}$ of \mathbb{C}^{4m} by permuting the indices. It is an easy check to see that under this identification $\bar{\tau}$ equals the element τ of Definition 2.2 for $n = 2m$ and we hence have the following (cf. Lemma 2.3):

Lemma 5.2. In the irreducible $Sp_{2(2m)}$ -representation $\tilde{V}(\tilde{\omega}_{2i}) \subset \Lambda^{2i} \mathbb{C}^{4m}$, $1 \leq i \leq 2m$, let $v_{\omega_{2i}}$ be the highest weight vector $v_{\omega_{2i}} = e_1 \wedge e_2 \wedge \dots \wedge e_{2i}$. Then (up to sign)

$$\bar{\tau}(v_{\omega_{2i}}) = e_1 \wedge e_2 \wedge \dots \wedge e_i \wedge e_{2m+1} \wedge e_{2m+2} \wedge \dots \wedge e_{2m+i}.$$

We denote by $\{\omega_i \mid 1 \leq i \leq m\}$, respectively $\{\tilde{\omega}_i \mid 1 \leq i \leq 2m\}$, the fundamental weights of \mathfrak{sp}_{2m} , respectively $\mathfrak{sp}_{2(2m)}$. They are characterized by the property $\langle \alpha_i^\vee, \omega_j \rangle = \delta_{i,j}$.

Definition 5.3. Let $\Psi : \mathfrak{h}^* \rightarrow \tilde{\mathfrak{h}}^*$ be the linear map defined on the weight lattice as follows

$$\Psi\left(\sum_{i=1}^{n-1} a_i \omega_i\right) := \sum_{i=1}^{n-1} a_i \tilde{\omega}_{2i}.$$

Note that Ψ sends dominant weights to dominant weights. Let $\lambda = b_1 \epsilon_1 + \dots + b_m \epsilon_m$, $b_1 \geq \dots \geq b_m \geq 0$ be a dominant weight for Sp_{2m} .

Lemma 5.4. $\bar{\tau}(\Psi(\lambda)) = b_1 \tilde{\epsilon}_1 + \dots + b_m \tilde{\epsilon}_m - b_m \tilde{\epsilon}_{m+1} - \dots - b_1 \tilde{\epsilon}_{2m}$.

Proof. Follows directly from Lemma 5.2 above. \square

As in the special linear case, we define a map from the set of negative roots of Sp_{2m} to the set of positive $Sp_{2(2m)}$ -roots by sending $\alpha_{i,j}$ to $\tilde{\alpha}_{j,i+2m}$. The following is the symplectic analogue of Lemma 2.5:

Lemma 5.5. *i) Let λ be a dominant weight for $Sp_{2m}(\mathbb{C})$. For a positive $Sp_{2(2m)}$ -root $\tilde{\alpha}$ we have $\langle \tilde{\alpha}^\vee, \tau(\Psi(\lambda)) \rangle < 0$ only if the $\tilde{\alpha}$ -root space lies in $Lie(\mathbb{U}(\mathfrak{n}^-))$.
ii) Let λ be a dominant Sp_{2m} -weight, let $\alpha = \alpha_{i,j}$ be a positive Sp_{2m} -root and let $\tilde{\alpha} = \tilde{\alpha}_{j,i+2m}$ be the $Sp_{2(2m)}$ positive root associated with $-\alpha$. Then we have:*

$$\langle \alpha^\vee, \lambda \rangle = -\langle \tilde{\alpha}^\vee, \tau(\Psi(\lambda)) \rangle.$$

iii) Let λ be a dominant weight for $Sp_{2m}(\mathbb{C})$ and let $\tilde{\alpha}$ be a positive $Sp_{2(2m)}$ -root. Then $E_{\tilde{\alpha}} v_{\tilde{\tau}} \neq 0$ in $\tilde{V}(\Psi(\lambda))$ only if $\tilde{\alpha} = \tilde{\alpha}_{j,i+2m}$, where $\alpha_{i,j}$ is a positive Sp_{2m} -root such that $\langle \alpha_{i,j}^\vee, \lambda \rangle > 0$.

Proof. Lemma 5.4 implies that for $\lambda = b_1 \epsilon_1 + \dots + b_{m-1} \epsilon_{m-1}$ we get

$$\langle (\tilde{\epsilon}_i - \tilde{\epsilon}_j)^\vee, \tau(\Psi(\lambda)) \rangle = \begin{cases} b_i - b_j \geq 0 & \text{if } 1 \leq i < j \leq m, \\ b_i + b_{2m-j+1} \geq 0 & \text{if } 1 \leq i \leq m < j \leq 2m, \\ -b_{2m-i+1} + b_{2m-j+1} \geq 0 & \text{if } m < i < j \leq 2m, \end{cases}$$

and

$$\langle (\tilde{\epsilon}_i + \tilde{\epsilon}_{j-2m})^\vee, \tau(\Psi(\lambda)) \rangle = \begin{cases} b_i + b_{j-2m} \geq 0 & \text{if } 1 \leq i \leq m, 2m < j \leq 3m, \\ b_i - b_{4m-j+1} \geq 0 & \text{if } 1 \leq i \leq m, \text{ and} \\ & 3m < j \leq 4m - i + 1, \\ b_i - b_{4m-j+1} \leq 0 & \text{if } 1 \leq i \leq m, 3m < j, \\ & \text{and } 4m - i + 1 < j, \\ -b_{2m-i+1} - b_{2m-j+1} \leq 0 & \text{if } m < i \leq 2m, 3m < j \end{cases}$$

where always $i + j \leq 4m$. This proves the corollary. \square

6. THE FUNDAMENTAL REPRESENTATIONS - THE \mathfrak{sp}_{2m} -CASE

Let $\mathfrak{n}_{\mathbb{Z}}^{-,a,i}$ be the direct sum of all root spaces of $Lie(Sp_{2(2m)})_{\mathbb{Z}}$ corresponding to positive roots β such that $\langle \beta^\vee, \tau(\tilde{\omega}_{2i}) \rangle < 0$ for all $1 \leq i \leq m$. By Lemma 5.5, such a space lies in $Lie(\mathbb{U}(\mathfrak{n}_{\mathbb{Z}}^-))$.

By [M, Lemme 26], the Demazure module $\tilde{V}_{\mathbb{Z}}(\ell\tilde{\omega}_{2i})_{\tau}$ is isomorphic to the algebra $U_{\mathbb{Z}}(\tilde{\mathfrak{n}})$ modulo the left ideal $\tilde{I}_{\mathbb{Z}}(\tau\ell\tilde{\omega}_{2i})$ generated by the $E_{k,l}^{(m)}$ for all $m \geq 1$ if $\langle \tilde{\alpha}_{k,l}^\vee, \tau\tilde{\omega}_{2i} \rangle \geq 0$, and $E_{k,l}^{(-\langle \tilde{\alpha}_{k,l}^\vee, \tau\ell\tilde{\omega}_{2i} \rangle + m)}$ for all $m \geq 1$ otherwise. Therefore all the root vectors (and their divided powers) not lying in $\mathfrak{n}_{\mathbb{Z}}^{-,a,i}$ act trivially on $\tilde{V}_{\mathbb{Z}}(\ell\tilde{\omega}_{2i})_{\tau}$ and hence in order to describe its structure as an $\mathfrak{n}_{\mathbb{Z}}^{-,a}$ -module it suffices to consider only the $\mathfrak{n}_{\mathbb{Z}}^{-,a,i}$ -action. Recall that $v_{\tilde{\tau}} = \tilde{\tau}(v_0)$ denotes the generator of $\tilde{V}_{\mathbb{Z}}(\Psi(\ell\omega_i))_{\tilde{\tau}}$, then the above discussion can be summarized as

$$(5) \quad \tilde{V}_{\mathbb{Z}}(\ell\tilde{\omega}_{2i})_{\tilde{\tau}} = U_{\mathbb{Z}}(\tilde{\mathfrak{n}}).v_{\tilde{\tau}} = U_{\mathbb{Z}}(\mathfrak{n}^{-,a}).v_{\tilde{\tau}} = U_{\mathbb{Z}}(\mathfrak{n}^{-,a,i}).v_{\tilde{\tau}}.$$

Recall that we embedded in $(SL_{2(2m)})_{\mathbb{Z}}$ a copy $L(i)_{\mathbb{Z}}$ of $(SL_{2m})_{\mathbb{Z}}$, so that we could identify the $SL_{2(2m)}$ -Demazure module $\tilde{V}_{\mathbb{Z}}^{SL_{2(2m)}}(\ell\tilde{\omega}_{2i})_{\bar{\tau}}$ generated by $\tau(v_0) = \bar{\tau}(v_0) = v_{\bar{\tau}}$ with the Weyl module $V_{\mathbb{Z}}^{L(i)}(\ell\varpi_i)$ for $L(i)_{\mathbb{Z}}$.

For $1 \leq k, l \leq 4m$, denote by $X_{k,l}$ the $4m \times 4m$ -matrix having a 1 in position (k, l) and whose all other entries are zero (for $k \neq l$ this is the $SL_{2(2m)}$ -root operator corresponding to the $SL_{2(2m)}$ -root $\alpha_{k,l}$), so that

$$\mathfrak{n}^{-,a,i} = \text{span}\{X_{r,s} + X_{4m-s+1,4m-r+1} \mid i+1 \leq r \leq 2m, 2m < s \leq 2m+i\}.$$

It is then immediate:

Lemma 6.1. *Every element $y \in \mathfrak{n}_{\mathbb{Z}}^{-,a,i}$ can be written in a unique way as $y = y_1 + y_2$, with $y_1 \in \mathfrak{n}^{-,a,i} \cap \text{Lie } L(i)_{\mathbb{Z}}$, $y_2 \in \text{span}\{X_{k,l} \mid l > 2m+k\}$. Moreover, y_2 is uniquely determined by y_1 .*

By the the previous lemma, the projection $p : (SL_{2(2m)})_{\mathbb{Z}} \rightarrow L(i)_{\mathbb{Z}}$ induces an isomorphism of vector spaces $\mathfrak{n}_{\mathbb{Z}}^{-,a,i} \simeq p(\mathfrak{n}_{\mathbb{Z}}^{-,a,i})$. Let us write $\bar{\mathfrak{n}}_{\mathbb{Z}}^{-,a,i}$ for $p(\mathfrak{n}_{\mathbb{Z}}^{-,a,i})$. Since the Lie algebras are commutative, we see that $p : \mathfrak{n}_{\mathbb{Z}}^{-,a,i} \rightarrow \bar{\mathfrak{n}}_{\mathbb{Z}}^{-,a,i}$ is not only an isomorphism of vector spaces, but it is in fact a Lie algebra isomorphism.

Corollary 6.2. $\tilde{V}_{\mathbb{Z}}(\Psi(\ell\omega_i))_{\bar{\tau}} = U_{\mathbb{Z}}(\bar{\mathfrak{n}}^{-,a,i}) \cdot v_{\bar{\tau}}$.

Proof. Let $y \in \mathfrak{n}_{\mathbb{Z}}^{-,a,i}$. By Lemma 6.1 we can write $y = p(y) + y_2$ with y_2 in the span of the matrices $X_{i,j}$ with $j > 2m+i$. Therefore y_2 acts trivially on $\tilde{V}_{\mathbb{Z}}(\Psi(\ell\omega_i))_{\bar{\tau}}$ and we conclude

$$\tilde{V}_{\mathbb{Z}}(\Psi(\ell\omega_i))_{\bar{\tau}} = U_{\mathbb{Z}}(\mathfrak{n}^{-,a,i}) \cdot v_{\bar{\tau}} = U_{\mathbb{Z}}(\bar{\mathfrak{n}}^{-,a,i}) \cdot v_{\bar{\tau}}.$$

□

For $0 \leq i \leq m-1$, let $Sp_{2m}(i)_{\mathbb{Z}}$ be a copy of $(Sp_{2m})_{\mathbb{Z}}$ sitting inside $L(i)_{\mathbb{Z}}$, defined with respect to the form given by the following matrix:

$$\begin{pmatrix} 0 & J_{m-i} & 0 & 0 \\ -J_{m-i} & 0 & 0 & 0 \\ 0 & 0 & 0 & J_i \\ 0 & 0 & -J_i & 0 \end{pmatrix},$$

where J_r denotes the $r \times r$ anti-diagonal matrix with entries $(1, 1, \dots, 1)$. Moreover, denote $(Sp_{2m}^{\text{std}})_{\mathbb{Z}} := Sp_{2m}(0)_{\mathbb{Z}}$.

Let σ be the permutation in $W^{L(i)}$ such that $\text{Lie } Sp_{2m}(i)_{\mathbb{Z}} = \sigma \text{Lie}(Sp_{2m}^{\text{std}})_{\mathbb{Z}} \sigma^{-1}$. Then the permutation σ is: keep $2m+1, \dots, 2m+i$ unchanged and move $2m-i+1, \dots, 2m$ in front of $i+1 \dots 2m-i$, so that

$$(6) \quad \begin{aligned} \sigma(e_1 \wedge e_2 \wedge \dots \wedge e_i \wedge e_{2m+1} \wedge e_{2m+2} \wedge \dots \wedge e_{2m+i}) = \\ e_1 \wedge e_2 \wedge \dots \wedge e_i \wedge e_{2m+1} \wedge e_{2m+2} \wedge \dots \wedge e_{2m+i}. \end{aligned}$$

Let $\mathfrak{b}_{\mathbb{Z}} \subset \text{Lie}(Sp_{2m}^{\text{std}})_{\mathbb{Z}}$ be the Borel subalgebra of upper triangular matrices. Let $\mathfrak{p}_{\mathbb{Z}}^i \subset \text{Lie}(Sp_{2m}^{\text{std}})_{\mathbb{Z}}$ be the maximal parabolic subalgebra associated with ϖ_i , and let $\mathfrak{p}_{\mathbb{Z}}^{i,n}$ be its nilpotent radical. Write $\mathfrak{Sp}_{\mathbb{Z}}^{i,n}$ for $\sigma \mathfrak{p}_{\mathbb{Z}}^{i,n} \sigma^{-1} \subset \text{Lie } Sp_{2m}(i)_{\mathbb{Z}}$.

Lemma 6.3. $V_{\mathbb{Z}}(\ell\varpi_i) = U_{\mathbb{Z}}(\mathbf{Sp}^{i,n}).v_{\bar{\tau}}$.

Proof. By (6), $v_{\bar{\tau}}$ is a lowest weight vector for $Sp_{2m}(i)$, as well as for $L(i)$, and we have that the module generated by this vector is $U_{\mathbb{Z}}(\text{Lie } Sp_{2m}(i)).v_{\bar{\tau}} = V_{\mathbb{Z}}(\ell\varpi_i)$. Since it is generated by a lowest weight vector, it is enough to consider the action of the nilpotent radical

$$U_{\mathbb{Z}}(\text{Lie } Sp_{2m}(i)).v_{\bar{\tau}} = U_{\mathbb{Z}}(\sigma\mathfrak{b}\sigma^{-1}).v_{\bar{\tau}} = U_{\mathbb{Z}}(\mathbf{Sp}^{i,n}).v_{\bar{\tau}}.$$

□

Observe that since $\bar{\mathfrak{n}}_{\mathbb{Z}}^{-,a,i} \subseteq \mathbf{Sp}_{\mathbb{Z}}^{i,n}$, the Weyl module $V_{\mathbb{Z}}(\ell\varpi_i)$ is naturally equipped with a structure of $\bar{\mathfrak{n}}_{\mathbb{Z}}^{-,a,i}$ -module. It is easy to check that:

Lemma 6.4. *Every element $x \in \mathbf{Sp}_{\mathbb{Z}}^{i,n}$ can be written in a unique way as $x = x_1 + x_2$, with $x_1 \in \bar{\mathfrak{n}}_{\mathbb{Z}}^{-,a,i}$, $x_2 \in \text{span} \{X_{k,l} \mid 2m - i < k \leq 2m, i + 1 < l < 2m\}$. Moreover, x_2 is uniquely determined by x_1 .*

As in Section 3, we consider the Chevalley involution $\iota : \mathfrak{sp}_{2m} \rightarrow \mathfrak{sp}_{2m}$ with $\iota|_{\mathfrak{h}} = -1$ and such that ι exchanges e_{α} and $-f_{\alpha}$. It induces an isomorphism $\iota : S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{-}) \rightarrow S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{+})$.

For a dominant weight λ , fix a highest weight vector $v_{\lambda} \in V_{\mathbb{Z}}(\lambda)$ and a lowest weight vector $v_{w_0} \in V_{\mathbb{Z}}(\lambda)$, where w_0 is the longest word in the Weyl group of \mathfrak{sp}_{2m} . Recall that considering the PBW filtration on $U_{\mathbb{Z}}(\mathfrak{n}^{-})$ and on $U_{\mathbb{Z}}(\mathfrak{n}^{+})$ provides $V_{\mathbb{Z}}(\lambda)$ with two possible $S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{-,a})$ -structures: in the first case looking at the PBW filtration on $V_{\mathbb{Z}}(\lambda)$ induced by the action of $U_{\mathbb{Z}}(\mathfrak{n}^{-})$ on the highest weight vector and taking the associated graded space provides the abelianized module $V_{\mathbb{Z}}^a(\lambda)$, while in the second case looking at the PBW filtration on $V_{\mathbb{Z}}(\lambda)$ induced by the action of $U_{\mathbb{Z}}(\mathfrak{n}^{+})$ on the lowest weight vector and taking the associated graded space produces a module that we denote by $V_{\mathbb{Z}}^{a,+}(\lambda)$. Now via ι this module also becomes naturally a $S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{-})$ -module and Lemma 3.3 holds in the symplectic case too:

Lemma 6.5. *The $S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{-})$ -module $V_{\mathbb{Z}}^{a,+}(\lambda)$ is, as $S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{-})$ -module, isomorphic to $V_{\mathbb{Z}}^a(\lambda)$.*

Observe that in the symplectic case there is no need of replacing λ by λ^* , since they coincide.

Lemma 6.6. *The Demazure module $\tilde{V}_{\mathbb{Z}}(\Psi(\ell\omega_i))_{\bar{\tau}}$ contained in $\tilde{V}_{\mathbb{Z}}(\Psi(\ell\omega_i))$ and $V_{\mathbb{Z}}(\ell\omega_i)^a$ are isomorphic as $S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}^{-})$ -modules.*

Proof. By Lemma 6.4, the projection $q : \mathbf{Sp}_{\mathbb{Z}}^{i,n} \rightarrow \bar{\mathfrak{n}}_{\mathbb{Z}}^{-,a,i}$ is an isomorphism of vector spaces. Moreover, if we write $x \in \mathbf{Sp}_{\mathbb{Z}}^{i,n}$ as $x = q(x) + x_2$, then x_2 lies in the span of the matrices $X_{i,j}$ with $i + 1 < j < 2m$ and hence $x_2.v_{\bar{\tau}} = 0$. Now, by [FFiL, Proposition 3.1], the PBW filtrations on $V_{\mathbb{Z}}(\ell\omega_i)$ with respect to the action of $\mathbf{Sp}_{\mathbb{Z}}^{i,n}$ and $\bar{\mathfrak{n}}_{\mathbb{Z}}^{-,a,i}$ are compatible and we get

$$\text{gr}V_{\mathbb{Z}}(\ell\varpi_i) = \text{gr}U_{\mathbb{Z}}(\mathbf{Sp}_{\mathbb{Z}}^{i,n}).v_{\bar{\tau}} \simeq \text{gr}U_{\mathbb{Z}}(\bar{\mathfrak{n}}_{\mathbb{Z}}^{-,a,i}).v_{\bar{\tau}}.$$

On the other hand, when we consider in $\tilde{V}_{\mathbb{Z}}(\Psi(\ell\omega_i))_{\bar{\tau}}$ the PBW filtration with respect to the action of $\mathbf{Sp}_{\mathbb{Z}}^{i,n}$ and go to the associated graded module, then the action of $(\mathbf{Sp}_{\mathbb{Z}}^{i,n})^a$ is isomorphic to the action of $\bar{\mathfrak{n}}_{\mathbb{Z}}^{-,a,i}$ on $\tilde{V}_{\mathbb{Z}}(\Psi(\ell\omega_i))_{\bar{\tau}}$. □

The previous result implies in particular:

Corollary 6.7. $\text{rank } \tilde{V}_{\mathbb{Z}}(\Psi(\ell\omega_i))_{\bar{\tau}} = \text{rank } V_{\mathbb{Z}}(\ell\omega_i).$

7. THE GENERAL CASE FOR \mathfrak{sp}_{2m}

We come now to the general case (notations as in Example 1.2):

Theorem 7.1. *Let λ be a dominant \mathfrak{sp}_{2m} -weight. The Demazure submodule $\tilde{V}_{\mathbb{Z}}(\Psi(\lambda))_{\bar{\tau}}$ of the $(Sp_{2(2m)})_{\mathbb{Z}}$ -module $\tilde{V}_{\mathbb{Z}}(\Psi(\lambda))$ is, as an $N_{\mathbb{Z},\eta}^{-,a}$ -module, isomorphic to the abelianized module $V_{\mathbb{Z}}^a(\lambda)$.*

As in the type **A**-case, the proof of the above theorem will provide us with a description of $V_{\mathbb{Z}}^a(\lambda)$ as an $S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}_{\eta}^{-,a})$ -module in terms of generators and relations. The abelianized module $V_{\mathbb{Z}}^a(\lambda)$ is a cyclic module over the algebra $S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}_{\eta}^{-,a})$ with the image of a highest weight vector $v_{\lambda} \in V(\lambda)$ in $V_{\mathbb{Z}}^a(\lambda)$ as a generator (cf. [FFL3, Proposition 2.3]). Hence the module is isomorphic to $S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}_{\eta}^{-,a})/I_{\mathbb{Z}}(\lambda)$ where $I_{\mathbb{Z}}(\lambda)$ is the annihilator of v_{λ} in $S_{\mathbb{Z}}(\mathfrak{n}_{\eta}^{-,a})$. As a consequence of the proof of Theorem 7.1, we obtain the description of the ideal $I_{\mathbb{Z}}(\lambda)$ in terms of generators given in [FFL1] and [FFL3] from Mathieu's generator and relation presentation of Demazure modules.

Let \mathfrak{b} be the Borel subalgebra of $\mathfrak{sp}_{2m} = (\mathfrak{sp}_{2m})_{\mathbb{Z}} \otimes \mathbb{C}$ as in Example 1.1 b), so $\mathfrak{sp}_{2m} = \mathfrak{n}^{-} \oplus \mathfrak{h} \oplus \mathfrak{n}^{+}$. As free \mathbb{Z} -modules, $U_{\mathbb{Z}}(\mathfrak{n}^{-}) \simeq U_{\mathbb{Z}}(\mathfrak{g})/U_{\mathbb{Z}}^{+}(\mathfrak{h} + \mathfrak{n}^{+})$, so that the adjoint action of $U_{\mathbb{Z}}(\mathfrak{b})$ on $U_{\mathbb{Z}}(\mathfrak{g})$ induces a structure of $U_{\mathbb{Z}}^{+}(\mathfrak{b})$ - and $B_{\mathbb{Z}}$ -module on $U_{\mathbb{Z}}(\mathfrak{n}^{-})$ and hence on $S_{\mathbb{Z}}(\mathfrak{n}_{\eta}^{-,a})$. This action is compatible with the $B_{\mathbb{Z}}$ -action on $V_{\mathbb{Z}}^a(\lambda)$ (cf. [FFL3, Prop. 2.3.]). Recall that for a positive root α we have denoted by f_{α} the corresponding fixed Chevalley basis element in $(\mathfrak{sp}_{2m})_{-\alpha,\mathbb{Z}}$. Let us set

$$R^{++} = \{\epsilon_i - \epsilon_j \mid 1 \leq i < j \leq m\} \cup \{2\epsilon_i \mid 1 \leq i \leq m\}$$

As a consequence of the proof of Theorem 7.1 we get the following description of the ideal $I_{\mathbb{Z}}(\lambda)$:

Corollary 7.2. *The abelianized module $V_{\mathbb{Z}}^a(\lambda)$ is as a cyclic $S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}_{\eta}^{-,a})$ -module isomorphic to $S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}_{\eta}^{-,a})/I_{\mathbb{Z}}(\lambda)$, where*

$$I_{\mathbb{Z}}(\lambda) = S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}_{\eta}^{-,a})(U_{\mathbb{Z}}(\mathfrak{n}^{+}) \circ \text{span}\{f_{\alpha}^{((\lambda, \alpha^{\vee})+m)} \mid m \geq 1, \alpha \in R^{++}\}) \subseteq S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}_{\eta}^{-,a}).$$

7.1. The proof of the theorem will be only sketched, since the strategy is the same as for the type **A**-case. We reproduce here the diagram of $S^{\bullet}(\mathfrak{n}_{\eta}^{-,a})$ -modules summarizing the main idea: for a dominant weight $\lambda = a_1\omega_1 + \dots + a_m\omega_m$, we have the following natural maps:

$$\begin{array}{ccccc} S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}_{\eta}^{-,a})/I_{\mathbb{Z}}(\lambda) & \xrightarrow[\simeq]{h} & V_{\mathbb{Z}}^a(\lambda) & \xrightarrow{b} & V_{\mathbb{Z}}^a(a_1\omega_1) \otimes \dots \otimes V_{\mathbb{Z}}^a(a_m\omega_m) \\ & & \downarrow a & & \downarrow c \\ S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}_{\eta}^{-,a})/M_{\mathbb{Z}}(\lambda) & \xrightarrow[\simeq]{g} & \tilde{V}_{\mathbb{Z}}(\Psi(\lambda))_{\tau} & \xrightarrow{d} & \tilde{V}_{\mathbb{Z}}(a_1\Psi(\omega_1))_{\tau} \otimes \dots \otimes \tilde{V}_{\mathbb{Z}}(a_m\Psi(\omega_m))_{\tau} \end{array}$$

where in the top row the action on the modules is twisted by the Chevalley involution, so the cyclic generators are lowest weight vectors, the maps c , d , a and g arise as in the proof of Theorem 4.1, so that again the main difficulty of the proof consists in producing the map f .

7.2. The first step consists in determining $M_{\mathbb{Z}}(\lambda)$. By [M, Lemme 26], the Demazure module $\tilde{V}_{\mathbb{Z}}(\Psi(\lambda))_{\tau}$ is isomorphic to the algebra $U_{\mathbb{Z}}(\tilde{\mathfrak{n}})$ modulo the left ideal $\tilde{I}_{\mathbb{Z}}(\tau\Psi(\lambda))$ generated by the elements $E_{k,l}^{(m)}$ for all $m \geq 1$ if $\langle \tilde{\alpha}_{k,l}^{\vee}, \tau\Psi(\lambda) \rangle \geq 0$, and $E_{k,l}^{(-\langle \tilde{\alpha}_{k,l}^{\vee}, \tau\Psi(\lambda) \rangle + m)}$ for all $m \geq 1$ otherwise.

7.3. The annihilator $M_{\mathbb{Z}}(\lambda)$ is the intersection of $U_{\mathbb{Z}}(\mathfrak{n}_{\eta}^{-,a}) \subset U_{\mathbb{Z}}(\tilde{\mathfrak{n}}^+)$ with the ideal $\tilde{I}_{\mathbb{Z}}(\tau\Psi(\lambda))$. To determine such an intersection, we fix a PBW basis and divide the positive roots in three families, exactly as in the proof of Theorem 4.1.

7.4. By Lemma 5.5(i), $\langle \tilde{\alpha}_{k,l}^{\vee}, \tau\Psi(\lambda) \rangle \geq 0$ if $\tilde{\alpha}_{k,l}$ is of third type. As in type **A**, we may hence proceed with the calculation modulo the left ideal generated by the divided powers of the corresponding $E_{k,l}$. Modulo such an ideal, by Lemma 5.4 and Lemma 5.5, $\tilde{I}_{\mathbb{Z}}(\tau\Psi(\lambda))$ is generated by the $E_{k,l}^{(m)}$, $m \geq 1$, for $\tilde{\alpha}_{k,l}$ of second type, and the $E_{k,l}^{(-\langle \tilde{\alpha}_{k,l}^{\vee}, \Psi(\lambda) \rangle + m)}$, $m \geq 1$, for $\tilde{\alpha}_{k,l}$ of first type.

7.5. Thus, we consider the subalgebra \mathfrak{a} generated by the $E_{k,l}^{(m)}$, for $\tilde{\alpha}_{k,l}$ of second type. Let $\mathfrak{b}_{\mathbb{Z}, \mathfrak{sl}_{2m}}$ be the Borel subalgebra of \mathfrak{sl}_{2m} consisting of traceless upper triangular matrices and let $\mathfrak{b}_{\mathbb{Z}}$ be the corresponding symplectic Borel subalgebra (of $(Sp_{2m}^{\text{std}})_{\mathbb{Z}}$). Let us embed $\mathfrak{b}_{\mathbb{Z}}$ in $\mathfrak{b}_{\mathbb{Z}, \mathfrak{sl}_{2m}} \oplus \mathfrak{b}_{\mathbb{Z}, \mathfrak{sl}_{2m}}$ via $A \mapsto (A, -\bar{A})$, where \bar{A} denotes the matrix which is skew-transposed to A , and let $\Delta^{-}(\mathfrak{b}_{\mathbb{Z}})$ be its image. Also \mathfrak{a} is embedded in $\mathfrak{b}_{\mathbb{Z}, \mathfrak{sl}_{2m}} \oplus \mathfrak{b}_{\mathbb{Z}, \mathfrak{sl}_{2m}}$, once we identify the latter with the Lie algebra generated by the divided powers of the SL_{4m} -root vectors of second type. The image of such an embedding contains $\Delta^{-}(\mathfrak{b}_{\mathbb{Z}})$. By taking fixed points with respect to the outer automorphism of \mathfrak{sl}_{2m} and \mathfrak{sl}_{4m} induced by the symmetry of the Dynkin diagram, it follows from Lemma 4.5,

$$(7) \quad U_{\mathbb{Z}}(\Delta^{-}(\mathfrak{b}_{\mathbb{Z}})) \langle \{E_{i,j}^{(m)} \mid \alpha_{i,j} \text{ of first type, } m \geq 1\} \rangle = U_{\mathbb{Z}}(\mathfrak{a}) \langle \{E_{i,j}^{(m)} \mid \alpha_{i,j} \text{ of first type, } m \geq 1\} \rangle.$$

Therefore,

$$\tilde{I}_{\mathbb{Z}}(\tau\Psi(\lambda)) \cap S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}_{\eta}^{-,a}) \simeq S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}_{\eta}^{-,a}) \circ U_{\mathbb{Z}}(\Delta^{-}(\mathfrak{b}_{\mathbb{Z}})) \text{span}\{f_{ij}^{\langle \alpha_{i,j}^{\vee}, \lambda \rangle + \ell} \mid \alpha \in R^{++}, \ell \geq 1\} =: M_{\mathbb{Z}}(\lambda).$$

7.6. *Proof of Theorem 7.1.* Since the roots of first type are precisely the ones coming from the elements f_{ij} with $\alpha_{i,j} \in R^{++}$ and since $\{f_{ij}^{\langle \alpha_{i,j}^{\vee}, \lambda \rangle + m} \mid \alpha_{i,j} \in R^{++}, m \geq 1\} \subseteq I_{\mathbb{Z}}(\lambda)$, we get a surjective morphism

$$(8) \quad \tilde{V}_{\mathbb{Z}}(\Psi\lambda)_{\bar{\tau}} \xrightarrow{g} S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}_{\eta}^{-,a})/M_{\mathbb{Z}}(\lambda) \xrightarrow{f} S_{\mathbb{Z}}^{\bullet}(\mathfrak{n}_{\eta}^{-,a})/I_{\mathbb{Z}}(\lambda) \simeq V_{\mathbb{Z}}(\lambda)^a.$$

This concludes the proof of the theorem. \square

7.7. Let ρ be the sum of the fundamental weights for Sp_{2m} and let $\tilde{\rho} = \Psi(\rho)$ be the corresponding dominant weight for $Sp_{2(2m)}$. Let $Q_{\mathbb{Z}} \subset (Sp_{2(2m)})_{\mathbb{Z}}$ be the corresponding parabolic subgroup. Recall that $N_{\mathbb{Z},\eta}^{-,a}$ is a commutative subgroup of the Borel subgroup $\tilde{B}_{\mathbb{Z}}$. For any $Sp_{2(2m)}$ -root $\tilde{\alpha}$ let $U_{\mathbb{Z},\tilde{\alpha}}$ be the associated root subgroup.

Lemma 7.3. *The orbit $\tilde{B}_{\mathbb{Z}}.\bar{\tau} \subset (Sp_{2(2m)})_{\mathbb{Z}}/Q_{\mathbb{Z}}$ is nothing but $N_{\mathbb{Z},\eta}^{-,a}.\bar{\tau}$, and the map $N_{\mathbb{Z},\eta}^{-,a} \rightarrow N_{\mathbb{Z},\eta}^{-,a}.\bar{\tau}$, $u \mapsto u\bar{\tau}$, is a bijection.*

Proof. We have $\tilde{B}_{\mathbb{Z}}.\bar{\tau} = \prod_{\tilde{\alpha}>0} U_{\mathbb{Z},\tilde{\alpha}}.\bar{\tau}$, and the map $\prod_{\tilde{\alpha}\in\Gamma} U_{\mathbb{Z},\tilde{\alpha}} \rightarrow \prod_{\tilde{\alpha}\in\Gamma} U_{\mathbb{Z},\tilde{\alpha}}.\bar{\tau}$ is a bijection, where Γ is the set of all positive roots of $Sp_{2(2m)}$ such that $\bar{\tau}^{-1}(\tilde{\alpha}) < 0$ and $\bar{\tau}^{-1}(\tilde{\alpha})$ is not an element of the root system of $Q_{\mathbb{Z}}$. Now this condition is fulfilled if and only if $\langle \bar{\tau}^{-1}(\tilde{\alpha}^{\vee}), \tilde{\rho} \rangle < 0$, or, equivalently, $\langle \tilde{\alpha}^{\vee}, \bar{\tau}(\tilde{\rho}) \rangle < 0$. By Lemma 5.4 this is not possible if $\tilde{\alpha}$ is of the form $\tilde{\alpha} = \tilde{\epsilon}_i - \tilde{\epsilon}_j$, $1 \leq i < j \leq 2m$. For the long roots this is only possible if $\tilde{\alpha} = 2\tilde{\epsilon}_j$, $j = m+1, \dots, 2m$, and for the roots $\alpha = \tilde{\epsilon}_i + \tilde{\epsilon}_j$, $1 \leq i < j \leq 2m$, this is only possible if either $i, j \geq m+1$ or $1 \leq i \leq m$ and $j = 2m+1-k$ is such that $1 \leq k < i$.

But this implies that the root subgroup $U_{\mathbb{Z},\tilde{\alpha}}$ is a subgroup of $N_{\mathbb{Z},\eta}^{-,a}$, and all root subgroups of $(Sp_{2(2m)})_{\mathbb{Z}}$ contained in $N_{\mathbb{Z},\eta}^{-,a}$ satisfy this condition. It follows that $N_{\mathbb{Z},\eta}^{-,a}.\bar{\tau}$ is the product of all root subgroups corresponding to positive roots of $Sp_{2(2m)}$ such that $\bar{\tau}^{-1}(\tilde{\alpha}) < 0$ and $\bar{\tau}^{-1}(\alpha)$ is not an element of the root system of $Q_{\mathbb{Z}}$ and hence $N_{\mathbb{Z},\eta}^{-,a}.\bar{\tau} = \tilde{B}_{\mathbb{Z}}.\bar{\tau} \subset (Sp_{2(2m)})_{\mathbb{Z}}/Q_{\mathbb{Z}}$. \square

Corollary 7.4. *The degenerate flag variety $\mathcal{F}\ell(\lambda)_k^a$ depends only on $\text{supp } \lambda$, it is a projectively normal variety, Frobenius split, with rational singularities.*

ACKNOWLEDGEMENTS

The work of M.L and P.L was partially supported by the DFG-Schwerpunkt Grant SP1388. The work of G.C.I. was financed by the national FIRB grant RBFR12RA9W ‘‘Perspectives in Lie Theory’’. We thank Corrado De Concini for many helpful conversations about this project.

REFERENCES

- [B] N. Bourbaki, *Éléments de mathématique. Fasc. XXXIV. Groupes et algèbres de Lie. Chapitres IV, V, VI.* Actualités Scientifiques et Industrielles, **1337** (1968), Hermann, Paris.
- [CL] G. Cerulli Irelli, M. Lanini, *Degenerate flag varieties of type A and C are Schubert varieties.* International Mathematics Research Notices, **15** (2015), 6353–6374.
- [FFL1] E. Feigin, G. Fourier, P. Littelmann, *PBW filtration and bases for irreducible modules in type A_n .* Transformation Groups, **16** (2011), no. 1, 71–89.
- [FFL2] E. Feigin, G. Fourier, P. Littelmann, *PBW filtration and bases for symplectic Lie algebras.* International Mathematics Research Notices, **24** (2011), 5760–5784.
- [FFL3] E. Feigin, G. Fourier, P. Littelmann, *PBW filtration over Z and compatible bases for $V(\lambda)$ in type A_n and C_n .* Symmetries, Integrable Systems and Representations, Springer Proceedings in Mathematics and Statistics, **40** (2012).
- [FF] E. Feigin, M. Finkelberg, *Degenerate flag varieties of type A: Frobenius splitting and BW theorem.* Mathematische Zeitschrift, **275** (2013), no. 1–2, 55–77.

- [FFiL] E. Feigin, M. Finkelberg, P. Littelmann, *Symplectic degenerate flag varieties*. Canadian Journal of Mathematics, **66** (2014), no. 6, 1250–1286.
- [F1] E. Feigin, \mathbb{G}_a^M *degeneration of flag varieties*. Selecta Mathematica, **18** (2012), no. 3, 513–537.
- [FH] W. Fulton, J. Harris *Representation Theory- A first course*. Graduate Texts in Mathematics, **129** (1991). Springer.
- [J] J. C. Jantzen *Representations of Algebraic Groups*. Pure and Applied Mathematics **131** (1987). Academic Press, Boston, MA.
- [H] J. E. Humphreys, *Introduction to Lie algebras and Representation Theory*. Graduate Texts in Mathematics, **9** (1972). Springer.
- [LMS] V. Lakshmibai, C. Musili, C. S. Seshadri, *Geometry of G/P* , Bulletin of the AMS, **1** (1979), no. 2, 432–435.
- [M] O. Mathieu, *Construction d'un groupe de Kac-Moody and applications*, Compositio Mathematica, **69** (1989), no. 1, 37–60.
- [MR] V. B. Mehta, A. Ramanathan *Schubert varieties in $G/B \times G/B$* , Compositio Mathematica, **67** (1988), 355–358.
- [P] C. Procesi *Lie Groups- An approach through Invariants and Representations*, Universitext, Springer (2007).
- [R] A. Ramanathan, *Equations defining Schubert varieties and Frobenius splitting of diagonals*, Inst. Hautes Étud. Sci. Publ. Math. **65** (1987), 61–90.
- [RR] S. Ramanan, A. Ramanathan *Projective normality of flag varieties and Schubert varieties*, Inventiones Mathematicae, **79** (1985), 217–224.

GIOVANNI CERULLI IRELLI: DIPARTIMENTO DI MATEMATICA. SAPIENZA-UNIVERSITÀ DI ROMA. PIAZZALE ALDO MORO 5, I-00185, ROME (ITALY)

E-mail address: `cerulli@mat.uniroma1.it`, `cerulli.math@googlemail.com`

MARTINA LANINI: SCHOOL OF MATHEMATICS, UNIVERSITY OF EDINBURGH, JAMES CLERK MAXWELL BUILDING, EDINBURGH, EH9 3FD (UK)

E-mail address: `m.lanini@ed.ac.uk`

PETER LITTELMANN: MATHEMATISCHES INSTITUT UNIVERSITÄT ZU KÖLN WEYERTAL 86-90 D-50931 KÖLN (GERMANY)

E-mail address: `peter.littelmann@math.uni-koeln.de`