

# ADDENDUM TO "EXISTENTIALLY COMPLETE CLOSURE ALGEBRAS"

Paolo Lipparini  
(14 maggio 1998)

*SOMMARIO. Si mostra, usando i metodi di [2], che la teoria universale delle algebre di chiusura è decidibile.*

A *Closure algebra* is a Boolean algebra together with an operator  $K$  satisfying the formal properties of closure in a topological space. McKinsey and Tarsky [1] announced that the (first-order) theory of closure algebras is undecidable. In contrast, the proof of [2, Theorem 1] shows:

**Theorem.** *The set of universal consequences of the theory of closure algebras is decidable.*

We now show how the arguments in [2] can be used to prove the theorem.

A universal sentence  $\psi$  is a consequence of the theory of closure algebras iff there is no closure algebra which is a model of the existential sentence  $\neg\psi$ , that is, iff no extension of the two-element closure algebra is a model of  $\neg\psi$ .

In [2] we dealt with a more general problem, that is, we considered an arbitrary closure algebra  $\hat{A}$  and an existential formula  $\varphi$  with constants from  $A$ , and we found necessary and sufficient conditions for the existence of an extension of  $\hat{A}$  satisfying  $\varphi$  (we supposed disjunctions were factored out, but this is no loss of generality). First, we showed how to reduce the existential formula  $\varphi$  to an equivalent form which involved only a polynomial  $P$  with certain coefficients  $a_{st}$  (defined in terms of the constants from  $A$ ); then we constructed increasing sequences  $w_{st}(m, \bar{a})$ , such that  $w_{st}(0, \bar{a}) = a_{st}$ , and such that an extension as above exists iff for all  $m$  the  $w_{st}(m, \bar{a})$ 's satisfy a certain property.

By our first remark here, we need to consider the particular case when  $\hat{A}$  is the two-element closure algebra; in this case each  $a_{st}$  and each  $w_{st}(m, \bar{a})$  is either 0 or 1; since each  $w_{st}(m, \bar{a})$  is increasing, as a function of  $m$ , it is eventually constant; moreover, the indexes  $s$  and  $t$  vary on a finite set, so that it is enough to consider a finite number of  $m$ 's (actually, just one). The procedure described in [2] is effective, and, by the preceding period, in this particular case, terminates in a finite number of steps, so that we have an effective way to decide whether a universal sentence is a consequence of the theory of closure algebras.

REMARKS. (i) A different proof can be found in [1]. We suspect that the algorithm furnished by the present proof is more efficient, but we have not checked this.

(ii) The constructions in [2] show that if a universal sentence fails in some closure algebra then it fails in a closure algebra whose underlying Boolean algebra is atomic and complete; that is, the sentence fails in a topological space (an atomic and complete Boolean algebra is isomorphic to the set of subsets of a set; we get a topological space by taking, as closed subsets, those  $x$ 's satisfying  $Kx=x$ ).

(iii) The same arguments show that set of universal consequences of the theory  $V_\alpha$  is decidable, too (cf. [2, Remark 2]).

We discovered the above arguments shortly after [2] had been printed, but only now we have written down the details.

References.

[1] McKinsey, A. Tarski, The algebra of topology, Ann.Math. (2) 45, 141-191 (1944).

[2] P. Lipparini, Existentially complete closure algebras, Bollettino UMI, Serie VI, Vol.I-D, N.1 (1982).

Paolo Lipparini  
Dipartimento di matematica  
Universita' di Tor Vergata  
Viale della Ricerca Scientifica  
I-00133 ROME  
ITALY