

# LOCAL UNIQUENESS AND NON-DEGENERACY OF BLOW UP SOLUTIONS OF MEAN FIELD EQUATIONS WITH SINGULAR DATA

DANIELE BARTOLUCCI, ALEKS JEVIKAR, YOUNGAE LEE, AND WEN YANG

**ABSTRACT.** We are concerned with the mean field equation with singular data on bounded domains. Under suitable non-degeneracy conditions we prove local uniqueness and non-degeneracy of bubbling solutions blowing up at singular points. The proof is based on sharp estimates for bubbling solutions of singular mean field equations and suitably defined Pohozaev-type identities.

**Keywords:** Mean field equations, uniqueness, non-degeneracy, blow up solutions, singular data.

## 1. INTRODUCTION

We are concerned with a sequence of solutions of the following mean field equation with singular data

$$\begin{cases} -\Delta u_n = \rho_n \frac{he^{u_n}}{\int_{\Omega} he^{u_n}} & \text{in } \Omega, \\ u_n = 0 & \text{on } \partial\Omega, \end{cases} \quad (\mathbf{P}_{\rho_n})$$

where  $\Omega \subset \mathbb{R}^2$  is a smooth bounded domain,  $h = h_* \exp(-4\pi \sum_{i=1}^N \alpha_i G(x, p_i))$ ,  $p_i$  are distinct points in  $\Omega$ ,  $\alpha_i \in (0, \infty) \setminus \mathbb{N}$ ,  $h_* \in C^\infty(\overline{\Omega})$ , and  $G$  is the Green function satisfying

$$\begin{cases} -\Delta G(x, p) = \delta_p & \text{in } \Omega, \\ G(x, p) = 0 & \text{on } \partial\Omega. \end{cases}$$

The mean field equation  $(\mathbf{P}_{\rho_n})$  (and its counterpart on compact surfaces) have been widely discussed in the last decades because of their several applications in Mathematics and Physics, such as Electroweak and Chern-Simons self-dual vortices [47, 49, 53], conformal metrics on surfaces with [50] or without conical singularities [35], statistical mechanics of two-dimensional turbulence [20] and of self-gravitating systems [52] and cosmic strings [45], and the theory of hyperelliptic curves [22] and of the Painlevé equations [24]. There are by now many results concerning existence [1, 3, 4, 5, 15, 21, 26, 28, 30, 31, 32, 36, 42], multiplicity [5, 29], uniqueness [6, 7, 10, 11, 12, 13, 14, 23, 33, 34, 40, 41, 48] and blow up analysis [2, 9, 16, 18, 17, 19, 25, 27, 37, 38, 39, 51, 54].

2010 *Mathematics Subject classification*: 35B32, 35J25, 35J61, 35J99, 82D15.

D. Bartolucci is partially supported by FIRB project “*Analysis and Beyond*”, by PRIN project 2012, ERC PE1.11, “*Variational and perturbative aspects in nonlinear differential problems*”, and by the Consolidate the Foundations project 2015 (sponsored by Univ. of Rome “Tor Vergata”), ERC PE1.11, “*Nonlinear Differential Problems and their Applications*”. Y. Lee is partially supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (No. NRF-2018R1C1B6003403). W. Yang is partially supported by NSFC No.11801550.

Our goal is to show that bubbling solutions of  $(\mathbf{P}_{\rho_n})$  blowing up at singular points  $p_i$  are unique and non-degenerate for  $n$  large enough.

**Definition 1.1.** Let  $u_n$  be a sequence of solutions of  $(\mathbf{P}_{\rho_n})$ . We say that  $u_n$  is a regular  $m$ -bubbling solution blowing up at the points  $q_j \notin \{p_1, \dots, p_N\}$ ,  $j = 1, \dots, m$ , if,

$$\frac{he^{u_n}}{\int_{\Omega} he^{u_n} dx} \rightharpoonup 8\pi \sum_{j=1}^m \delta_{q_j},$$

weakly in the sense of measures in  $\Omega$ .

We say that  $u_n$  is a singular  $m$ -bubbling solution blowing up at the points  $p_j \in \{p_1, \dots, p_N\}$ ,  $j = 1, \dots, m$ ,  $m \leq N$  if,

$$\frac{he^{u_n}}{\int_{\Omega} he^{u_n} dx} \rightharpoonup 8\pi \sum_{j=1}^m (1 + \alpha_j) \delta_{p_j},$$

weakly in the sense of measures in  $\Omega$ .

To state the main result and to compare it with the existing literature we introduce some notation. Let  $R(x, y) = \frac{1}{2\pi} \log |x - y| + G(x, y)$  be the regular part of  $G(x, y)$ . For what concerns regular bubbling solutions, for  $\mathbf{q} = (q_1, \dots, q_m) \in \overline{\Omega} \times \dots \times \overline{\Omega}$ , we let  $G_j^*(x) = 8\pi R(x, q_j) + 8\pi \sum_{l \neq j}^{1, \dots, m} G(x, q_l)$  and

$$\ell_{\text{reg}}(\mathbf{q}) = \sum_{j=1}^m [\Delta \log h(q_j)] h(q_j) e^{G_j^*(q_j)}.$$

For  $(x_1, \dots, x_m) \in \overline{\Omega} \times \dots \times \overline{\Omega}$ , we also define the  $m$ -vortex Hamiltonian,

$$\mathcal{H}_m(x_1, x_2, \dots, x_m) = \sum_{j=1}^m [\log(h(x_j)) + 4\pi R(x_j, x_j)] + 4\pi \sum_{l \neq j}^{1, \dots, m} G(x_l, x_j). \quad (1.1)$$

Then, by assuming suitable non-degeneracy conditions the authors in [8, 9] proved that regular  $m$ -bubbling solutions are unique and non-degenerate (see also [10] for an analogous result for the Gelfand equation).

**Theorem A** ([8, 9]). Let  $u_n^{(1)}$  and  $u_n^{(2)}$  be two regular  $m$ -bubbling solutions of  $(\mathbf{P}_{\rho_n})$ , with  $\rho_n^{(1)} = \rho_n = \rho_n^{(2)}$ , blowing up at the points  $q_j \notin \{p_1, \dots, p_N\}$ ,  $j = 1, \dots, m$ , where  $\mathbf{q} = (q_1, \dots, q_m)$  is a critical point of  $\mathcal{H}_m$ . Assume that,

- (1)  $\det(D^2 \mathcal{H}_m(\mathbf{q})) \neq 0$ ,
- (2)  $\ell_{\text{reg}}(\mathbf{q}) \neq 0$ .

Then there exists  $n_0 \geq 1$  such that  $u_n^{(1)} = u_n^{(2)}$  for all  $n \geq n_0$ . Moreover, the linearized problem at a  $m$ -bubbling solution  $u_n$

$$\begin{cases} \Delta \phi + \rho_n \frac{he^{u_n}}{\int_{\Omega} he^{u_n} dx} \left( \phi - \frac{\int_{\Omega} he^{u_n} \phi dx}{\int_{\Omega} he^{u_n} dx} \right) = 0 & \text{in } \Omega, \\ \phi = 0 & \text{on } \partial\Omega, \end{cases} \quad (1.2)$$

admits only the trivial solution  $\phi \equiv 0$  for any  $n \geq n_0$ .

The above condition (2) can be relaxed by assuming  $\ell_{\text{reg}}(\mathbf{q}) = 0$  and  $D(\mathbf{q}) \neq 0$ , where  $D(\mathbf{q})$  is a geometric quantity. Our aim is to extend the latter result to singular bubbling solutions. Even though the argument works out for more general situations we focus here on singular 1-bubbling solution blowing up at  $p_i$  for some  $i \in \{1, \dots, N\}$ , see also Remark 1.3. More precisely, we assume without loss of generality that  $\alpha_i \neq \alpha_j$  for  $i \neq j$  and we study the case  $\rho_n \rightarrow 8\pi(1 + \alpha_i)$  for some fixed  $i \in \{1, \dots, N\}$  and

$$\|u_n\|_{L^\infty(\Omega)} \rightarrow +\infty \quad \text{as } n \rightarrow +\infty.$$

We define

$$\ell(p_i) = \frac{2\pi^2}{(1 + \alpha_i) \sin\left(\frac{\pi}{1 + \alpha_i}\right)} \left( \frac{(1 + \alpha_i)}{\pi \bar{h}_1(p_i)} \right)^{\frac{1}{1 + \alpha_i}} \Delta \log h_*(p_i), \quad (1.3)$$

where  $h(x) = \bar{h}_1(x)|x - p_i|^{2\alpha_i}$ . Moreover, we define the 'desingularized' 1-vortex Hamiltonian to be

$$\mathcal{H}_{p_i}(x) = 8\pi(1 + \alpha_i)(R(x, p_i) - R(p_i, p_i)) + (\log \bar{h}_1(x) - \log \bar{h}_1(p_i)). \quad (1.4)$$

Our main results are the following.

**Theorem 1.1.** *Let  $u_n^{(1)}$  and  $u_n^{(2)}$  be two singular 1-bubbling solutions of  $(\mathbf{P}_{\rho_n})$ , with  $\rho_n^{(1)} = \rho_n = \rho_n^{(2)}$ , blowing up at the point  $p_i$  for some  $i \in \{1, \dots, N\}$ ,  $\alpha_i \in (0, \infty) \setminus \mathbb{N}$ . Assume that,*

- (1)  $p_i$  is a critical point of  $\mathcal{H}_{p_i}$ ,
- (2)  $\ell(p_i) \neq 0$ .

*Then there exists  $n_0 \geq 1$  such that  $u_n^{(1)} = u_n^{(2)}$  for all  $n \geq n_0$ .*

**Theorem 1.2.** *Let  $u_n$  be a singular 1-bubbling solution of  $(\mathbf{P}_{\rho_n})$ , blowing up at the point  $p_i$  for some  $i \in \{1, \dots, N\}$ ,  $\alpha_i \in (0, \infty) \setminus \mathbb{N}$ . Assume that the conditions (1)-(2) of Theorem 1.1 hold true. Then there exists  $n_0 \geq 1$  such that, for any  $n \geq n_0$ , (1.2) admits only the trivial solution  $\phi \equiv 0$ .*

Observe that we do not need the non-degeneracy of the Hamiltonian as in condition (1) of Theorem A. This is essentially due to the difference of the linearized problem, see (1.8) and the discussion later on. On the other hand, we do need to assume  $p_i$  to be a critical point of  $\mathcal{H}_{p_i}$ . For the regular blow up this is always the case since it is well-known [44] that for a regular  $m$ -bubbling solution blowing up at the points  $q_j \notin \{p_1, \dots, p_N\}$ , then  $\mathbf{q} = (q_1, \dots, q_m)$  has to be a critical point of  $\mathcal{H}_m$ .

**Remark 1.3.** *The argument yielding Theorems 1.1 and 1.2 works out for more general situations and can be carried out to prove local uniqueness of singular  $m$ -bubbling and even for mixed scenarios of singular  $m$ -bubbling and regular  $m'$ -bubbling solutions. The decision to focus on singular 1-bubbling is twofold: on one side the latter case is very subtle since in general the singular blow up point does not need be a critical point of the Hamiltonian  $\mathcal{H}_{p_i}$  and furthermore we are not assuming any non-degeneracy of  $\mathcal{H}_{p_i}$ , and on the other side we want to highlight the differences with respect to the regular case. We postpone the general situation to a future paper. The case  $\alpha \in (-1, 0)$  will be treated in a separate paper since we first need to derive suitable sharp estimates for bubbling solutions,*

which are still missing in this case. Finally, the case  $\alpha \in \mathbb{N}$  is by now out of reach due to the presence of non-simple (and non-radial) blow up [18, 37].

To prove Theorem 1.1 we argue by contradiction and we analyze the asymptotic behavior of the (normalized) difference of two distinct solutions for  $(\mathbf{P}_{\rho_n})$ ,

$$\xi_n = \frac{u_n^{(1)} - u_n^{(2)}}{\|u_n^{(1)} - u_n^{(2)}\|_{L^\infty(\Omega)}}. \quad (1.5)$$

Near the blow up point  $p_i$ , and after a suitable scaling,  $\xi_n$  converges to an entire solution of the linearized problem of the Liouville equation

$$\Delta v + |x|^{2\alpha_i} e^v = 0 \quad \text{in } \mathbb{R}^2. \quad (1.6)$$

Solutions of (1.6) with finite mass are completely classified [46] and for  $\alpha_i \in (0, \infty) \setminus \mathbb{N}$  take the form,

$$v(z) = v_\mu(z) = \log \frac{8(1 + \alpha_i)^2 e^\mu}{(1 + e^\mu |z|^{2(1+\alpha_i)})^2}, \quad \mu \in \mathbb{R}. \quad (1.7)$$

The freedom in the choice of  $\mu$  is due to the invariance of equation (1.6) under dilations. The linearized operator  $L$  relative to  $v_0$  is defined by,

$$L\phi := \Delta\phi + \frac{8(1 + \alpha_i)^2 |z|^{2\alpha_i}}{(1 + |z|^{2(1+\alpha_i)})^2} \phi \quad \text{in } \mathbb{R}^2. \quad (1.8)$$

It follows from [27, Corollary 2.2] that the  $L^\infty$ -bounded kernel of  $L$  has one eigenfunction  $Y_0$ , where,

$$Y_0(z) = \frac{1 - |z|^{2(1+\alpha_i)}}{1 + |z|^{2(1+\alpha_i)}} = \frac{\partial v_\mu}{\partial \mu} \Big|_{\mu=0}.$$

The main part of the proof of Theorem 1.1 is to show that, after scaling and for large  $n$ ,  $\xi_n$  is orthogonal to  $Y_0$ . This is done by a delicate analysis of a suitably defined Pohozaev-type identity first introduced in [43] and then exploited in [8, 10].

The proof of Theorem 1.2 follows the same strategy by analyzing the asymptotic behavior of

$$\Xi_n = \frac{\phi_n - \frac{\int_\Omega h e^{u_n} \phi_n dx}{\int_\Omega h e^{u_n} dx}}{\left\| \phi_n - \frac{\int_\Omega h e^{u_n} \phi_n dx}{\int_\Omega h e^{u_n} dx} \right\|_{L^\infty(\Omega)}},$$

for a non-trivial solution  $\phi_n$  of (1.2), which plays the role of (1.5).

The paper is organized as follows. In section 2 we introduce some preliminary results, in section 3 we estimate the  $L^\infty$ -norm of the difference of two solutions to  $(\mathbf{P}_{\rho_n})$  and in section 4 we then deduce the first estimates of  $\xi_n$ , the normalized difference of two solutions, away from the blow up point. In section 5 we introduce a Pohozaev-type identity to get refined estimates on  $\xi_n$  and prove Theorem 1.1. Finally, in section 6 we give the sketch of the proof of Theorem 1.2.

## 2. PRELIMINARY ESTIMATES ABOUT THE BLOW UP PHENOMENON AT THE SINGULAR POINT

In this section we collect some preliminary results which will be used in the sequel. Let us assume that  $i = 1$  and set  $p = p_1, 0 \neq \alpha = \alpha_1 \in (0, +\infty) \setminus \mathbb{N}$ . We define

$$\tilde{u}_n = u_n - \log \left( \int_{\Omega} h e^{u_n} dx \right), \quad \lambda_n = \max_{\Omega} \tilde{u}_n, \quad \sigma_n^{2(1+\alpha)} = e^{-\lambda_n}, \quad (2.1)$$

and

$$U_n(x) = \lambda_n - 2 \log(1 + \gamma_n e^{\lambda_n} |x - p|^{2+2\alpha}), \quad \gamma_n = \frac{\rho_n \bar{h}_1(p)}{8(1+\alpha)^2},$$

where

$$\bar{h}_1(x) = h_* \exp(-4\pi \bar{G}_1(x)) \quad \bar{G}_1(x) = \sum_{i=2}^N \alpha_i G(x, p_i) + R(x, p),$$

and  $R(x, y) = G(x, y) + \frac{1}{2\pi} \log |x - y|$  is the regular part of the Green function. Therefore, we have

$$h(x) = \bar{h}_1(x) |x - p|^{2\alpha},$$

and in any small enough ball centered at  $p$  it holds that  $\bar{h}_1 > 0$ . It has been shown in [2] (for  $\alpha \in (0, +\infty) \setminus \mathbb{N}$ ) and [17] (for  $\alpha \in (-1, 0)$ ) that

$$|\tilde{u}_n(x) - U_n(x)| \leq C, \quad \forall x \in B_r(p). \quad (2.2)$$

Actually the proofs in [18, 2] show that this estimate holds locally near  $p$ , but then the global estimate follows by looking at the Definition 1 and the Green representation formula.

More recently, it has been proved in [27] that if  $\alpha \in (0, +\infty) \setminus \mathbb{N}$ , then

$$\rho_n - 8\pi(1+\alpha) = \ell(p) e^{-\frac{\lambda_n}{1+\alpha}} + O(e^{-\lambda_n \frac{1+\epsilon_0}{1+\alpha}}) \text{ as } n \rightarrow +\infty, \quad (2.3)$$

and

$$\rho_{n,1} - 8\pi(1+\alpha) = \ell(p) e^{-\frac{\lambda_n}{1+\alpha}} + O(e^{-\lambda_n \frac{1+\epsilon_0}{1+\alpha}}) \text{ as } n \rightarrow +\infty, \quad (2.4)$$

where

$$\rho_{n,1} = \int_{B(p, r_0)} h e^{\tilde{u}_n}, \quad \ell(p) = \frac{2\pi^2}{(1+\alpha) \sin(\frac{\pi}{1+\alpha})} \left( \frac{(1+\alpha)}{\pi \bar{h}_1(p)} \right)^{\frac{1}{1+\alpha}} \Delta \log h_*(p),$$

and

$$\epsilon_0 = 2 - 2(1-\alpha)^+ = \begin{cases} 2, & \text{if } \alpha \geq 1, \\ 2\alpha, & \text{if } \alpha \in (0, 1). \end{cases}$$

Next, we set

$$R_{n,1} = \rho_{n,1} R(x, p),$$

let  $r_0 > 0$  be a small positive number and set

$$v_n = \tilde{u}_n - (R_{n,1}(x) - R_{n,1}(p)), \quad x \in B(p, r_0), \quad (2.5)$$

and as in [54], we denote  $\psi_n$  as the solution of

$$\begin{cases} \Delta\psi_n = 0, & \text{in } B(p, 4r_0), \\ \psi_n = v_n - \frac{1}{8\pi r_0} \int_{|x-p|=4r_0} v_n ds & \text{on } \partial B(p, 4r_0). \end{cases}$$

By the Mean value Theorem, we have  $\psi(0) = 0$ . It has been proved in [54] that

$$v_n - U_n - \psi_n(x) = \sigma_n \psi_{n,1} \left( \frac{x-p}{\sigma_n} \right) + \sigma_n^2 \psi_{n,2} \left( \frac{x-p}{\sigma_n} \right) + O(\sigma_n^2) \text{ in } B(p, 4r_0), \quad (2.6)$$

where

$$\psi_{n,1}(y) = -\frac{2(1+\alpha)a_{n,1}}{\alpha} \frac{y_1}{1 + \gamma_n |y|^{2(1+\alpha)}}, \quad y = (y_1, y_2) \in \mathbb{R}^2, \quad (2.7)$$

and

$$\psi_{n,2}(y) = -a_n \log(2 + |y|) + a_{n,0} + O(|y|^{-\epsilon_0}), \quad y = (y_1, y_2) \in \mathbb{R}^2 \setminus B(0, R_0), \quad (2.8)$$

for suitable  $R_0 \geq 1$ . Here  $a_{n,0}$  is a uniformly bounded sequence,

$$a_n = \frac{\pi}{(1+\alpha) \sin(\frac{\pi}{1+\alpha})} \left( \frac{8(1+\alpha)^2}{\rho_n \bar{h}_1(p)} \right)^{\frac{1}{1+\alpha}} \Delta \log h_*(p),$$

and, composing with suitable rotations, we can assume that

$$(a_{n,1}, 0) = \nabla \log \left( \bar{h}_1(x) e^{R_{n,1}(x) + \psi_n(x)} \right) \Big|_{x=p}. \quad (2.9)$$

Moreover, it has been shown in [27, Lemma 3.2] that

$$\psi_n(x) = O(\sigma_n^2), \quad x \in B(p, 4r_0). \quad (2.10)$$

Since  $\psi_n$  is harmonic, then we also have

$$|\nabla \psi_n(x)| = O(\sigma_n^2), \quad x \in B(p, 3r_0). \quad (2.11)$$

We also have, see [27, Lemma 3.1],

$$u_n(x) - \rho_n G(x, p) = O(\sigma_n), \quad x \in \bar{\Omega} \setminus B(p, r_0). \quad (2.12)$$

Also, we will need the following improved estimate obtained by matching (2.6) and (2.12).

**Lemma 2.1.** *It holds,*

$$\lambda_n - \log \left( \int_{\Omega} h e^{u_n} \right) + 2 \log \gamma_n + 8\pi(1+\alpha)R(p, p) = O(\sigma_n). \quad (2.13)$$

*Proof.* Putting  $c_n = \log \left( \int_{\Omega} h e^{u_n} \right)$  and picking any  $|x-p| = 2r_0$  in (2.6) and (2.12), we conclude that

$$\rho_n G(x, p) - c_n - (R_{n,1}(x) - R_{n,1}(p)) - U_n(x) = O(\sigma_n).$$

Clearly we have

$$U_n(x) = -\lambda_n - 4(1+\alpha) \log |x-p| - 2 \log(\gamma_n) + O(\sigma_n^{2(1+\alpha)}),$$

and we find that

$$-\frac{\rho_n}{2\pi} \log |x - p| + \rho_n R(x, p) - c_n - \rho_{n,1} R(x, p) + \rho_{n,1} R(p, p) + \lambda_n + 4(1 + \alpha) \log |x - p| + 2 \log(\gamma_n) = O(\sigma_n),$$

and then the desired conclusion easily follows from (2.3) and (2.4).  $\square$

Finally, similar arguments used in the estimate (2.12), yield

$$\nabla(\tilde{u}_n - \rho_n G(x, p)) = O(\sigma_n), \quad x \in \overline{\Omega} \setminus B(p, r_0). \quad (2.14)$$

### 3. ESTIMATE OF THE $L^\infty$ -NORM

The proof of Theorem 1.1 is obtained by contradiction and we assume that two distinct solutions  $u_n^{(i)}$ ,  $i = 1, 2$ , exist for  $(\mathbf{P}_{\rho_n})$ , whence in particular with the same  $\rho_n$ , which satisfy

$$\rho_n \rightarrow 8\pi(1 + \alpha) \quad \text{as } n \rightarrow +\infty,$$

where  $\alpha = \alpha_1$ . We also assume without loss of generality that

$$p_1 = 0 \in \Omega.$$

Then we define

$$\tilde{u}_n^{(i)} = u_n^{(i)}(x) - \log \left( \int_{\Omega} h e^{u_n^{(i)}} \right), \quad \lambda_n^{(i)} = \max_{\overline{\Omega}} \tilde{u}_n^{(i)},$$

and in particular  $v_n^{(i)}$  defined as in (2.5). Also we set

$$U_n^{(i)}(x) = \lambda_n^{(i)} - 2 \log(1 + \gamma_n e^{\lambda_n^{(i)}} |x - p|^{2(1+\alpha)}), \quad i = 1, 2, \quad \gamma_n = \frac{\rho_n \bar{h}_1(p)}{8(1 + \alpha)^2}.$$

There is no loss of generality in assuming that

$$\lambda_n^{(1)} \leq \lambda_n^{(2)}.$$

To simplify the notation, we set

$$\sigma_n^{2(1+\alpha)} = e^{-\lambda_n^{(1)}}.$$

Then we have

**Lemma 3.1.** (i)  $|\lambda_n^{(1)} - \lambda_n^{(2)}| = O(\sum_{i=1}^2 e^{-\frac{\epsilon_0}{1+\alpha} \lambda_n^{(i)}}) = O(e^{-\frac{\epsilon_0}{1+\alpha} \lambda_n^{(1)}}) = O(\sigma_n^{2\epsilon_0}).$

$$(ii) \quad \|\tilde{u}_n^{(1)} - \tilde{u}_n^{(2)}\|_{L^\infty(B(0, r_0))} \leq |\lambda_n^{(2)} - \lambda_n^{(1)}| + O(\lambda_n^{(1)} \sigma_n^2).$$

$$(iii) \quad \|\tilde{u}_n^{(1)} - \tilde{u}_n^{(2)}\|_{L^\infty(\Omega \setminus B(0, r_0))} \leq O(\sigma_n).$$

*Proof.* (i) In view of (2.3), we find that

$$\ell(p) e^{-\frac{\lambda_n^{(1)}}{1+\alpha}} + O\left(e^{-\frac{1+\epsilon_0}{1+\alpha} \lambda_n^{(1)}}\right) = \ell(p) e^{-\frac{\lambda_n^{(2)}}{1+\alpha}} + O\left(e^{-\frac{1+\epsilon_0}{1+\alpha} \lambda_n^{(2)}}\right),$$

which immediately implies, since  $\ell(p) \neq 0$ ,

$$\lambda_n^{(1)} - \lambda_n^{(2)} = O(e^{-\frac{\epsilon_0}{1+\alpha} \lambda_n^{(1)}}) + e^{-\frac{\epsilon_0}{1+\alpha} \lambda_n^{(2)}},$$

as claimed.

(ii) By using  $\lambda_n^{(1)} \leq \lambda_n^{(2)}$ , it is not difficult to see that

$$\begin{aligned} U_n^{(2)} - U_n^{(1)} &= (\lambda_n^{(2)} - \lambda_n^{(1)}) \left( \frac{1 - e^{\lambda_n^{(1)}} \gamma_n |x - p|^{2(1+\alpha)}}{1 + e^{\lambda_n^{(1)}} \gamma_n |x - p|^{2(1+\alpha)}} \right) + O((\lambda_n^{(2)} - \lambda_n^{(1)})^2) \\ &\leq |\lambda_n^{(2)} - \lambda_n^{(1)}| + O((\lambda_n^{(2)} - \lambda_n^{(1)})^2), \end{aligned}$$

uniformly in  $B(0, r)$  for any  $r > 0$ . Also, in view of (2.10), and since the  $\psi_n^{(i)}$ 's are harmonic, we find that

$$\psi_n^{(2)}(x) - \psi_n^{(1)}(x) = O(\sigma_n^2), \quad |\nabla(\psi_n^{(2)}(x) - \psi_n^{(1)}(x))| = O(\sigma_n^2),$$

uniformly in  $B(0, 3r_0)$ , we use this gradient estimate to evaluate the difference,

$$|a_{n,2} - a_{n,1}| = |\nabla(\psi_n^{(2)}(0) - \psi_n^{(1)}(0))| = O(\sigma_n^2),$$

which implies that

$$\begin{aligned} |\psi_{n,2}^{(2)}(x) - \psi_{n,1}^{(1)}(x)| &\leq \frac{2(1+\alpha)}{\alpha} |a_{n,1}^{(2)} - a_{n,1}^{(1)}| |x_1| + O(\lambda_n^{(2)} - \lambda_n^{(1)}) \\ &= O(\sigma_n^2) + O(\lambda_n^{(2)} - \lambda_n^{(1)}), \end{aligned}$$

uniformly in  $B(0, r_0)$ . Also it is easy to see that

$$|\psi_{n,2}^{(2)} - \psi_{n,2}^{(1)}| = O(\lambda_n^{(1)} \sigma_n^2).$$

Therefore, in view of (2.6) and Lemma 3.1, we finally conclude that,

$$\|\tilde{u}_n^{(1)}(x) - \tilde{u}_n^{(2)}(x)\|_{L^\infty(B(0, r_0))} \leq |\lambda_n^{(2)} - \lambda_n^{(1)}| + O(\lambda_n^{(1)} \sigma_n^2),$$

which is (ii).

(iii) Next we obtain the estimate in  $\Omega \setminus B(0, r_0)$ , by using the Green's representation formula,

$$\begin{aligned} \tilde{u}_n^{(1)}(x) - \tilde{u}_n^{(2)}(x) &= \rho_n \int_{\Omega} G(y, x) h(y) (e^{\tilde{u}_n^{(1)}(y)} - e^{\tilde{u}_n^{(2)}(y)}) dy \\ &= \rho_n \int_{B(0, r_0)} (G(y, x) - G(0, x)) h(y) (e^{\tilde{u}_n^{(1)}(y)} - e^{\tilde{u}_n^{(2)}(y)}) dy \\ &\quad + G(0, x) \int_{B(0, r_0)} \rho_n h(y) (e^{\tilde{u}_n^{(1)}(y)} - e^{\tilde{u}_n^{(2)}(y)}) dy \\ &\quad + \rho_n \int_{\Omega \setminus B(0, r_0)} G(y, x) h(y) (e^{\tilde{u}_n^{(1)}(y)} - e^{\tilde{u}_n^{(2)}(y)}) dy. \end{aligned}$$

In view of (2.2) and since  $\rho_n$  is the same for the two solutions, then we have

$$\begin{aligned} \rho_{n,1}^{(1)} - \rho_{n,1}^{(2)} &= \rho_n \int_{B(0, r_0)} h(y) e^{\tilde{u}_n^{(1)}(y)} - \rho_n \int_{B(0, r_0)} h(y) e^{\tilde{u}_n^{(2)}(y)} \\ &= \rho_n \int_{\Omega \setminus B(0, r_0)} h(y) \left( e^{\tilde{u}_n^{(2)}(y)} - e^{\tilde{u}_n^{(1)}(y)} \right) dy = O(e^{-\lambda_n}). \end{aligned}$$



Then, by using (2.2) once more, for  $x \in \Omega \setminus B(0, r_0)$  we have,

$$\begin{aligned}
& \tilde{u}_n^{(1)}(x) - \tilde{u}_n^{(2)}(x) \\
&= \rho_n \int_{B(0, r_0)} (G(y, x) - G(0, x)) h(y) \left( e^{\tilde{u}_n^{(2)}(y)} - e^{\tilde{u}_n^{(1)}(y)} \right) dy \\
&\quad + G(0, x) (\rho_{n,1}^{(1)} - \rho_{n,2}^{(2)}) + O(e^{-\lambda_n}) \\
&= \rho_n \int_{B(0, r_0)} (G(y, x) - G(0, x)) h(y) \left( e^{\tilde{u}_n^{(2)}(y)} - e^{\tilde{u}_n^{(1)}(y)} \right) dy + O(e^{-\lambda_n}) \\
&= \int_{B(0, r_0)} O(1) \left( \sum_{i=1,2} \frac{|y|^{2\alpha+1} e^{\lambda_n^{(i)}}}{(1 + \gamma_n e^{\lambda_n^{(i)}} |y|^{2+2\alpha})^2} \right) dy + O(e^{-\lambda_n}) \\
&= O(\sigma_n),
\end{aligned}$$

uniformly in  $x \in \Omega \setminus B(0, r_0)$ . Therefore we conclude that

$$\|\tilde{u}_n^{(1)}(x) - \tilde{u}_n^{(2)}(x)\|_{L^\infty(\Omega \setminus B(0, r_0))} \leq O(\sigma_n),$$

as claimed.  $\square$

#### 4. ESTIMATE OF THE DIFFERENCE AWAY FROM THE BLOW UP POINT

Let

$$\xi_n = \frac{\tilde{u}_n^{(1)} - \tilde{u}_n^{(2)}}{\|\tilde{u}_n^{(1)} - \tilde{u}_n^{(2)}\|_{L^\infty(\Omega)}}. \quad (4.1)$$

Clearly  $\xi_n$  satisfies

$$\begin{cases} \Delta \xi_n + \rho_n h(x) c_n(x) \xi_n(x) = 0 & \text{in } \Omega, \\ \xi_n = -d_n & \text{on } \partial\Omega, \end{cases} \quad (4.2)$$

for some constant  $d_n$  satisfying  $|d_n| \leq 1$  and

$$c_n(x) = \frac{e^{\tilde{u}_n^{(1)}} - e^{\tilde{u}_n^{(2)}}}{\tilde{u}_n^{(1)} - \tilde{u}_n^{(2)}}.$$

To simplify the notations, we set

$$\lambda_n = \lambda_n^{(1)} \quad \text{and} \quad \sigma_n^{2+2\alpha} = e^{-\lambda_n}.$$

Then by defining

$$\hat{\xi}_n(z) = \xi_n(\sigma_n z), \quad |z| < 4\sigma_n^{-1} r_0,$$

we prove the following

**Lemma 4.1.** *There exists a constant  $b_0 \in \mathbb{R}$ , such that  $\hat{\xi}_n(z) \rightarrow b_0 \hat{\xi}_0(z)$  in  $C_{\text{loc}}^0(\mathbb{R}^2)$ , where*

$$\hat{\xi}_0(z) = \frac{1 - \gamma |z|^{2+2\alpha}}{1 + \gamma |z|^{2+2\alpha}}, \quad z \in \mathbb{R}^2,$$

where  $\gamma = \frac{\pi \bar{h}_1(0)}{1+\alpha}$ .

*Proof.* By Lemma 3.1, we see that

$$\begin{aligned} c_n(x) &= e^{\tilde{u}_n^{(1)}(x)} \left( 1 + O(\|\tilde{u}_n^{(1)} - \tilde{u}_n^{(2)}\|_{L^\infty(\Omega)}) \right) \\ &= e^{\tilde{u}_n^{(1)}(x)} (1 + O(|\lambda_n^{(2)} - \lambda_n^{(1)}| + \sigma_n)), \end{aligned}$$

and then by (2.6), (2.10) and (2.11)

$$e^{-\lambda_n} c_n(\sigma_n z) = \frac{e^{C\sigma_n(1+O(|\lambda_n^{(2)} - \lambda_n^{(1)}| + \sigma_n))}}{(1 + \gamma_n |z|^{2+2\alpha})^2} \rightarrow \frac{1}{(1 + \gamma |z|^{2+2\alpha})^2} \quad \text{in } C_{\text{loc}}^2(\mathbb{R}^2),$$

where  $\gamma = \frac{\pi \bar{h}_1(0)}{1+\alpha}$ .

We define

$$\Omega_{\sigma_n} = \{z \in \mathbb{R}^2 \mid \sigma_n z \in \Omega\}.$$

By using (4.2), we have

$$\begin{cases} \Delta \hat{\xi}_n + \rho_n \bar{h}_1(\sigma_n z) |z|^{2\alpha} \sigma_n^{2+2\alpha} & \text{in } \Omega_{\sigma_n}, \\ \hat{\xi}_n(z) = -d_n & \text{on } \partial\Omega_{\sigma_n}, \end{cases}$$

and since  $|\hat{\xi}_n| \leq 1$ , then we conclude that  $\hat{\xi}_n \rightarrow \hat{\xi}$  in  $C_{\text{loc}}^0(\mathbb{R}^2)$ , where  $\hat{\xi}$  is a solution of

$$\Delta \hat{\xi} + \frac{8\gamma(1+\alpha)^2 |z|^{2\alpha}}{(1 + \gamma |z|^{2(1+\alpha)})^2} \hat{\xi} = 0 \text{ in } \mathbb{R}^2 \quad \text{and} \quad |\hat{\xi}(z)| \leq 1 \text{ in } \mathbb{R}^2.$$

It follows from [27, Corollary 2.2] that  $\hat{\xi}(z) = b_0 \xi_0(z)$ , for some constant  $b_0$ , as claimed.  $\square$

Next, we have

**Lemma 4.2.** *For any  $r_0$  small enough we have*

$$\xi_n(x) = -b_0 + o(1), \quad x \in \Omega \setminus B(0, r_0),$$

where  $b_0$  is defined by Lemma 4.1.

*Proof.* It follows from (2.2) that

$$c_n(x) \rightarrow 0 \quad \text{in } C_{\text{loc}}^0(\overline{\Omega} \setminus \{0\}).$$

Since  $\|\xi_n\|_{L^\infty(\Omega)} \leq 1$ , then (4.2) implies that

$$\xi_n \rightarrow \xi_0 \quad \text{in } C_{\text{loc}}^0(\overline{\Omega} \setminus \{0\}),$$

where

$$\Delta \xi_0 = 0 \text{ in } \Omega \setminus \{0\} \quad \text{and} \quad \|\xi_0\|_{L^\infty(\Omega)} \leq 1.$$

As a consequence,  $\xi_0$  is smooth in  $\Omega \setminus \{0\}$  and in particular

$$\Delta \xi_0 = 0 \quad \text{in } \Omega.$$

Therefore  $\xi_0 = -b$  in  $\Omega$  for some constant  $b$  and

$$\xi_n \rightarrow -b \quad \text{in } C_{\text{loc}}^0(\overline{\Omega} \setminus \{0\}). \tag{4.3}$$

In particular  $-\xi_n(x) = d_n \rightarrow b$  for  $x \in \partial\Omega$ . Let  $\phi_n = \frac{1-\gamma_n e^{\lambda_n}|x|^{2+2\alpha}}{1+\gamma_n e^{\lambda_n}|x|^{2+2\alpha}}$  and let us fix  $d \in (0, r_0)$ . Then, by using (2.6), (2.10) and (2.11), we find that

$$\begin{aligned} & \int_{\partial B(0,d)} \left( \phi_n \frac{\partial \xi_n}{\partial \nu} - \xi_n \frac{\partial \phi_n}{\partial \nu} \right) d\sigma = \int_{B(0,d)} (\phi_n \Delta \xi_n - \xi_n \Delta \phi_n) dx \\ &= \int_{B(0,d)} \left\{ -\rho_n \xi_n \phi_n \bar{h}_1(x) |x|^{2\alpha} \left( \frac{e^{\tilde{u}_n^{(1)}} - e^{\tilde{u}_n^{(2)}}}{\tilde{u}_n^{(1)} - \tilde{u}_n^{(2)}} \right) + 8(1+\alpha)^2 \gamma_n \xi_n \phi_n |x|^{2\alpha} e^{U_n} \right\} dx \\ &= \int_{B(0,d)} \rho_n \xi_n \phi_n \left\{ -\bar{h}_1(x) |x|^{2\alpha} e^{\tilde{u}_n^{(1)}} (1 + O(|\tilde{u}_n^{(1)} - \tilde{u}_n^{(2)}|)) + \bar{h}_1(0) |x|^{2\alpha} e^{U_n} \right\} dx \\ &= \int_{B(0,d)} \rho_n \xi_n \phi_n |x|^{2\alpha} e^{U_n} \left\{ -\bar{h}_1(x) e^{O(\sigma_n)} (1 + O(|\tilde{u}_n^{(1)} - \tilde{u}_n^{(2)}|)) + \bar{h}_1(0) \right\} dx. \end{aligned}$$

Therefore, by the scaling  $x = \sigma_n z$ , we see that,

$$\begin{aligned} & \int_{\partial B(0,d)} \left( \phi_n \frac{\partial \xi_n}{\partial \nu} - \xi_n \frac{\partial \phi_n}{\partial \nu} \right) d\sigma \\ &= \int_{B(0,d/\sigma_n)} \rho_n \hat{\xi}_n(z) \hat{\phi}_n(z) |z|^{2\alpha} \frac{O(1)(\sigma_n |z| + |\hat{u}_n^{(1)} - \hat{u}_n^{(2)}| + \sigma_n)}{(1 + \gamma_n |z|^{2+2\alpha})^2} dz. \end{aligned}$$

In view of Lemma 3.1 we obtain

$$\int_{\partial B(0,d)} \left( \phi_n \frac{\partial \xi_n}{\partial \nu} - \xi_n \frac{\partial \phi_n}{\partial \nu} \right) d\sigma = O(\sigma_n + \sigma_n^{2\epsilon_0}). \quad (4.4)$$

Let  $\zeta_n = \int_0^{2\pi} \xi_n(r, \theta) d\theta$ , where  $r = |x|$ . Then, for any fixed  $R > 0$ , (4.4) yields

$$(\zeta_n)'(r) \phi_n(r) - \zeta_n(r) \phi_n'(r) = \frac{O(\sigma_n + \sigma_n^{2\epsilon_0})}{r}, \quad \forall r \in (R\sigma_n, r_0].$$

Also for any  $R > 0$  large enough, and for any  $r \in (R\sigma_n, r_0]$ , we also obtain that

$$\phi_n(r) = -1 + O\left(\frac{\sigma_n^{2+2\alpha}}{r^{2+2\alpha}}\right), \quad \phi_n'(r) = O\left(\frac{\sigma_n^{2+2\alpha}}{r^{3+2\alpha}}\right),$$

and so we conclude that

$$\zeta_n'(r) = \frac{O(\sigma_n + \sigma_n^{2\epsilon_0})}{r} + O\left(\frac{\sigma_n^{2+2\alpha}}{r^{3+2\alpha}}\right), \quad \forall r \in (R\sigma_n, r_0]. \quad (4.5)$$

Integrating (4.5) we obtain that

$$\zeta_n(r) = \zeta_n(R\sigma_n) + o(1) + O(R^{-(2+2\alpha)}), \quad \forall r \in (R\sigma_n, r_0]. \quad (4.6)$$

In view of Lemma 4.1, we also have

$$\zeta_n(R\sigma_n) = -2\pi b_0 + o_R(1) + o_n(1),$$

where  $\lim_{R \rightarrow +\infty} o_R(1) = 0$  and  $\lim_{n \rightarrow +\infty} o_n(1) = 0$ . Then by (4.6) we have

$$\zeta_n(r) = -2\pi b_0 + o_R(1) + o_n(1)(1 + O(R)), \quad \forall r \in (R\sigma_n, r_0]. \quad (4.7)$$

In view of (4.3), we see that

$$\zeta_n = -2\pi b + o_n(1) \text{ in } C_{\text{loc}}(\Omega \setminus \{0\}),$$

which implies that  $b = b_0$ . Hence, we finish the proof.  $\square$

Next, we need a refined estimate about  $\xi_n$  which will be needed in next section.

**Lemma 4.3.**

$$\xi_n(x) = -d_n + A_n G(0, x) + o(\sigma_n) \quad \text{in } C^1(\Omega \setminus B(0, 2r_0)), \quad (4.8)$$

where

$$A_n = \int_{\Omega} f_n^*(x) \quad \text{and} \quad f_n^*(x) = \rho_n c_n(x) h(x) \xi_n(x).$$

Moreover, there is a constant  $C > 0$ , which does not depend on  $R > 0$ , such that

$$|\xi_n(x) + d_n - A_n G(0, x)| \leq C \sigma_n \left( \frac{1_{B(0, 2r_0)}(x)}{|x|} + 1_{\Omega \setminus B(0, 2r_0)}(x) \right), \quad x \in \Omega \setminus B(0, R\sigma_n). \quad (4.9)$$

*Proof.* By the Green representation formula we find that,

$$\begin{aligned} \xi_n(x) &= -d_n + \int_{\Omega} G(y, x) f_n^*(y) dy \\ &= -d_n + A_n G(0, x) + \int_{\Omega} (G(y, x) - G(0, x)) f_n^*(y) dy, \end{aligned} \quad (4.10)$$

while, by Lemma 3.1, we also find that

$$c_n(x) \xi_n(x) = \frac{e^{\tilde{u}_n^{(1)}} - e^{\tilde{u}_n^{(2)}}}{\|\tilde{u}_n^{(1)} - \tilde{u}_n^{(2)}\|_{L^\infty(\Omega)}} = e^{\tilde{u}_n^{(1)}} \xi_n(x) (1 + O(\lambda_n^{(2)} - \lambda_n^{(1)} + \sigma_n)). \quad (4.11)$$

Thus, for  $x \in \Omega \setminus B(0, 2r_0)$ , we see from (2.2), (2.6) that

$$\begin{aligned} \int_{\Omega} (G(y, x) - G(0, x)) f_n^*(y) dy &= \int_{B(0, r_0)} (G(y, x) - G(0, x)) f_n^*(y) dy + O(e^{-\lambda_n}) \\ &= \int_{B(0, r_0)} \langle \partial_y G(y, x) |_{y=0, y} \rangle f_n^*(y) dy + O(1) \left( \frac{|y|^{2+2\alpha} e^{\lambda_n}}{(1 + e^{\lambda_n} |y|^{2+2\alpha})^2} dy \right) \\ &= \int_{B(0, r_0)} \langle \partial_y G(y, x) |_{y=0, y} \rangle f_n^*(y) dy + O(\sigma_n^2). \end{aligned} \quad (4.12)$$

By using (2.2), (2.6) and Lemma 3.1, after scaling we see that for  $x \in \Omega \setminus B(0, 2r_0)$ , it holds

$$\begin{aligned} &\int_{B(0, r_0)} \langle \partial_y G(y, x) |_{y=0, y} \rangle f_n^*(y) dy \\ &= \sigma_n^{3+2\alpha} \int_{B(0, r_0/\sigma_n)} \langle \partial_y G(y, x) |_{y=0, z} \rangle \rho_n \bar{h}_1(\sigma_n z) |z|^{2\alpha} e^{\hat{U}_n + \hat{R}_{n,1}(z) - \hat{R}_{n,1}(0)} \hat{\xi}_n(z) dz \\ &\quad + O(\sigma_n^{1+2\epsilon_0} + \sigma_n^2) \\ &= \sigma_n \int_{B(0, r_0/\sigma_n)} \frac{\langle \partial_y G(y, x) |_{y=0, z} \rangle \rho_n \bar{h}_1(0) |z|^{2\alpha} \hat{\xi}_n(z)}{(1 + \gamma_n |z|^{2+2\alpha})^2} dz + O(\sigma_n^{1+2\epsilon_0} + \sigma_n^2). \end{aligned}$$

Therefore, in view of Lemma 4.1, for  $x \in \Omega \setminus B(0, 2r_0)$  we find that,

$$\begin{aligned} &\int_{B(0, r_0)} \langle \partial_y G(y, x) |_{y=0, y} \rangle f_n^*(y) dy \\ &= \sigma_n \sum_{h=1}^2 \partial_{y_h} G(y, x) |_{y=0} \rho_n \bar{h}_1(0) b_0 \int_{B(0, r_0/\sigma_n)} \frac{z_h |z|^{2\alpha} \hat{\xi}_0(z)}{(1 + \gamma_n |z|^{2+2\alpha})^2} dz + o(\sigma_n) \quad (4.13) \\ &= \sigma_n \sum_{h=1}^2 \partial_{y_h} G(y, x) |_{y=0} \rho_n \bar{h}_1(0) b_0 \int_{\mathbb{R}^2} \frac{z_h |z|^{2\alpha} \hat{\xi}_0(z)}{(1 + \gamma_n |z|^{2+2\alpha})^2} dz + o(\sigma_n). \end{aligned}$$

From (4.10)-(4.13), we see that the estimate (4.8) holds in  $C^0(\Omega \setminus B(0, r_0))$ . The proof of the fact that (4.8) holds in  $C^1(\Omega \setminus B(0, r_0))$  is similar and we skip it here to avoid repetitions.

From (4.11), (2.6) and suitable scaling, we see that there exists  $C > 0$ , which is independent of  $R > 0$  such that for  $x \in B(0, 2r_0) \setminus B(0, \sigma_n R)$ , it holds that

$$\begin{aligned}
 |\xi_n(x) + d_n - A_n G(0, x)| &\leq \left| \int_{B(0, 3r_0)} (G(y, x) - G(0, x)) f_n^*(y) dy \right| + O(e^{-\lambda_n}) \\
 &\leq \left| \frac{1}{2\pi} \int_{B(0, 3r_0)} \log \frac{|x|}{|x-y|} f_n^*(y) dy \right| + O\left( \int_{B(0, 3r_0)} \frac{e^{\lambda_n} |y|^{1+2\alpha}}{(1 + e^{\lambda_n} |y|^{2+2\alpha})^2} dy \right) + O(e^{-\lambda_n}) \\
 &\leq O(1) \left( \int_{B(0, 3r_0/\sigma_n)} \frac{|\log |x| - \log |x - \sigma_n z|| |z|^{2\alpha}}{(1 + |z|^{2+2\alpha})^2} dz \right) + O(\sigma_n) \\
 &\leq O(1) \left( \int_{\frac{|x|/\sigma_n}{2} \leq |z| \leq 2|x|/\sigma_n} \frac{|\log |x| - \log |x - \sigma_n z|| |z|^{2\alpha}}{(1 + |z|^{2+2\alpha})^2} dz \right) \\
 &\quad + O(1) \left( \int_{B(0, 3r_0/\sigma_n)} \frac{\sigma_n |z|^{2\alpha+1}}{|x| (1 + |z|^{2+2\alpha})^2} dz \right) + O(\sigma_n) \\
 &\leq O(1) \left( \frac{\sigma_n}{|x|} \right) + O(1) (\log |z| |z|^{-2} |_{|z|=|x|/\sigma_n}) + O(\sigma_n) \leq C \left( \frac{\sigma_n}{|x|} \right).
 \end{aligned} \tag{4.14}$$

By (4.10), (4.11) and (2.6), we also see that for  $x \in \Omega \setminus B(0, 2r_0)$ , it holds that

$$|\xi_n(x) + d_n - A_n G(x, 0)| = O\left( \int_{B(0, r_0)} \frac{e^{\lambda_n} |y|^{1+2\alpha}}{(1 + e^{\lambda_n} |y|^{2+2\alpha})^2} dy \right) + O(e^{-\lambda_n}) = O(\sigma_n). \tag{4.15}$$

By (4.14) and (4.15) we obtain (4.9), which concludes the proof of Lemma 4.3.  $\square$

## 5. ESTIMATES VIA POHOZAEV IDENTITIES

From now on, for a given function  $f(y, x)$ , we shall use  $\partial$  and  $D$  to denote the partial derivatives with respect to  $y$  and  $x$  respectively. With a small abuse of notation, for a function  $f(x)$  we will use both  $\nabla$  and  $D$  to denote its gradient.

We define

$$\varphi_n(y) = \rho_n(R(y, 0) - R(0, 0)), \tag{5.1}$$

and

$$v_n^{(i)} = \tilde{u}_n^{(i)} - \varphi_n(y), \quad i = 1, 2. \tag{5.2}$$

Recall the definition of  $\xi_n$  which satisfies (4.2). Our aim is to show that the projection of  $\xi_n$  on the radial part kernel is zero, i.e.,  $b_0 = 0$ . We shall accomplish it by exploiting the following Pohozaev identity to derive a more accurate estimate on  $\xi_n$ .

**Lemma 5.1.** ([43]) *For any fixed  $r \in (0, r_0)$ , it holds*

$$\begin{aligned} & \frac{1}{2} \int_{\partial B(0,r)} r \langle Dv_n^{(1)} + Dv_n^{(2)}, D\xi_n \rangle d\sigma - \int_{\partial B(0,r)} r \langle v, D(v_n^{(1)} + v_n^{(2)}) \rangle \langle v, D\xi_n \rangle d\sigma \\ &= \int_{\partial B(0,r)} \frac{r\rho_n h(x)}{\|v_n^{(1)} - v_n^{(2)}\|_{L^\infty(\Omega)}} (e^{v_n^{(1)} + \varphi_n} - e^{v_n^{(2)} + \varphi_n}) d\sigma \\ & \quad - \int_{B(0,r)} \frac{\rho_n h(x)(e^{v_n^{(1)} + \varphi_n} - e^{v_n^{(2)} + \varphi_n})}{\|v_n^{(1)} - v_n^{(2)}\|_{L^\infty(\Omega)}} \left(2 + 2\alpha + \langle D(\log \bar{h}_1(x) + \varphi_n(x)), x \rangle\right) dx. \end{aligned} \quad (5.3)$$

*Proof.* See [8] for a proof of this identity.  $\square$

Let

$$\begin{aligned} \Phi(y, 0) &= -8\pi(1 + \alpha) \log |y| + 8\pi(1 + \alpha)(R(y, 0) - R(0, 0)) \\ & \quad + \log(\bar{h}_1(y)) - \log(\bar{h}_1(0)). \end{aligned} \quad (5.4)$$

Recall the definition of  $A_n$  given in Lemma 4.3. Then we have

**Lemma 5.2.**

$$\begin{aligned} \text{L.H.S. of (5.3)} &= -4(1 + \alpha)A_n - \frac{(8(1 + \alpha)^2)^3 b_0 e^{-\lambda_n}}{2\rho_n \bar{h}_1(0)} \int_{\Omega \setminus B(0,r)} |y|^{2\alpha} e^{\Phi(y,0)} \\ & \quad + o(\sigma_n^2) + O(\sigma_n |A_n|) + O(r^{-3} \sigma_n^3). \end{aligned}$$

*Proof.* Let

$$G_n(x) = \rho_n G(x, 0), \quad (5.5)$$

so that

$$\nabla(G_n(x) - \varphi_n)(x) = -\frac{\rho_n}{2\pi} \frac{x}{|x|^2}. \quad (5.6)$$

In view of (2.14), we have

$$\begin{aligned} \nabla v_n^{(i)}(x) &= \nabla(\bar{u}_n^{(i)} - G_n(x)) + \nabla(G_n(x) - \varphi_n(x)) \\ &= \nabla(G_n(x) - \varphi_n(x)) + O(\sigma_n), \quad x \in \bar{\Omega} \setminus B(0, r_0), \end{aligned}$$

for any fixed small  $r_0 > 0$ . As a consequence, for fixed  $r > r_0$ , we find that

$$\begin{aligned} \text{L.H.S. of (5.3)} &= \int_{\partial B(0,r)} r \langle D(G_n - \varphi_n), D\xi_n \rangle d\sigma - 2 \int_{\partial B(0,r)} r \langle v, D(G_n - \varphi_n) \rangle \langle v, D\xi_n \rangle d\sigma \\ & \quad + O(\sigma_n \|D\xi_n\|_{L^\infty(\partial B(0,r))}) \\ &= \int_{\partial B(0,r)} \frac{\rho_n}{2\pi} \langle D\xi_n, v \rangle d\sigma + O(\sigma_n \|D\xi_n\|_{L^\infty(\partial B(0,r))}) \\ &= \int_{\partial B(0,r)} 4(1 + \alpha) \langle D\xi_n, v \rangle d\sigma + O(\sigma_n \|D\xi_n\|_{L^\infty(\partial B(0,r))}), \end{aligned} \quad (5.7)$$

where we used (2.3). Therefore, as a consequence of Lemma 4.3, we conclude that

$$\text{L.H.S. of (5.3)} = 4(1 + \alpha) \int_{\partial B(0,r)} \langle D\xi_n, v \rangle d\sigma + O(\sigma_n |A_n|) + o(\sigma_n^2). \quad (5.8)$$

In this particular case, we have  $A_n = 0$ .

To estimate the right hand side of (5.8), we need a refined estimate about  $\xi_n$  on  $\partial B(0, r)$ . So, by the Green representation formula with  $x \in \partial B(0, r)$ , we find that

$$\begin{aligned}\xi_n(x) &= -d_n + \int_{\Omega} G(y, x) f_n^*(y) dy \\ &= -d_n + A_n G(0, x) + \sum_{h=1}^2 B_{n,h} \partial_{y_h} G(y, x) \big|_{y=0} + \frac{1}{2} \sum_{h,k=1}^2 C_{n,h,k} \partial_{y_h y_k}^2 G(y, x) \big|_{y=0} \\ &\quad + \int_{\Omega} \Psi_n(y, x) f_n^*(y),\end{aligned}\tag{5.9}$$

where

$$A_n = \int_{\Omega} f_n^*(y) dy, \quad B_{n,h} = \int_{B(0,r)} y_h f_n^*(y) dy, \quad C_{n,h,k} = \int_{B(0,r)} y_h y_k f_n^*(y) dy,$$

and

$$\begin{aligned}\Psi_n(y, x) &= G(y, x) - G(0, x) - \langle \partial_y G(y, x) \big|_{y=0}, y \rangle 1_{\partial B(0,r)}(y) \\ &\quad - \frac{1}{2} \langle \partial_y^2 G(y, x) \big|_{y=0}, y, y \rangle 1_{B(0,r)}(y).\end{aligned}$$

At this point, let us fix  $\bar{\theta} \in (0, \frac{r}{2})$ . By Lemma 3.1 and Lemma 4.2, we find that,

$$\begin{aligned}f_n^*(y) &= \rho_n \bar{h}_1 |y|^{2\alpha} e^{\tilde{u}_n^{(1)}} (\xi_n(y) + O(\|\tilde{u}_n^{(1)} - \tilde{u}_n^{(2)}\|_{L^\infty(\Omega)})) \\ &= \rho_n \bar{h}_1 |y|^{2\alpha} e^{\tilde{u}_n^{(1)}} (-b_0 + o(1)),\end{aligned}\tag{5.10}$$

for any  $y \in \partial\Omega \setminus B(0, \bar{\theta})$ . By (2.4), (2.12), (2.13) and (5.10), we conclude that

$$\begin{aligned}f_n^*(y) &= \rho_n \bar{h}_1 |y|^{2\alpha} e^{\rho_{n,1}^{(1)} G(y,0) - \lambda_n - 2 \log(\gamma_n) - 8\pi(1+\alpha)R(0,0)} (-b_0 + o(1)) \\ &= (8(1+\alpha)^2)^2 \frac{e^{-\lambda_n}}{\rho_n \bar{h}_1(0)} |y|^{2\alpha} e^{\Phi(y,0)} (-b_0 + o(1)) \quad \text{for } y \in \Omega \setminus B(0, \bar{\theta}),\end{aligned}\tag{5.11}$$

where

$$\Phi(y, 0) = -4(1+\alpha) \log |y| + 8\pi(1+\alpha)(R(y, 0) - R(0, 0)) + \log(\bar{h}_1(0)) - \log(\bar{h}_1(0)).$$

On the other hand, by (2.6), we have for  $y \in B(0, \bar{\theta})$ ,

$$f_n^*(y) = \rho_n h e^{\tilde{u}_n^{(1)}} (\xi_n + O(\|\tilde{u}_n^{(1)} - \tilde{u}_n^{(2)}\|_{L^\infty(\Omega)})) = O\left(\frac{|y|^{2\alpha} e^{\lambda_n}}{(1 + e^{\lambda_n} |y|^{2+2\alpha})^2}\right). \tag{5.12}$$

Next, by (5.10), for  $y \in B(0, \bar{\theta})$  and  $x \in \partial B(0, r)$ , we get

$$\Psi_n(y, x) = O\left(\frac{|y|^3}{|x|^3}\right), \quad \text{and} \quad \nabla_x \Psi_n(y, x) = O\left(\frac{|y|^3}{|x|^4}\right). \tag{5.13}$$

Let us define

$$\bar{G}_n(x) = A_n G(0, x) + \sum_{h=1}^2 B_{n,h} \partial_{y_h} G(y, x) \big|_{y=0} + \frac{1}{2} \sum_{h,k=1}^2 C_{n,h,k} \partial_{y_h y_k}^2 G(y, x) \big|_{y=0}, \tag{5.14}$$

so that, by (5.11)-(5.13), we conclude that for  $x \in \partial B(0, r)$ , it holds

$$\begin{aligned}
\zeta_n(x) + d_n - \overline{G}_n(x) &= \int_{\Omega \setminus B(0, \overline{\theta})} \Psi_n(y, x) f_n^*(y) dy + \int_{B(0, \overline{\theta})} \Psi_n(y, x) f_n^*(y) dy \\
&= -b_0 \int_{\Omega \setminus B(0, \overline{\theta})} \frac{(8(1+\alpha)^2)^2 e^{-\lambda_n}}{\rho_n \overline{h}_1(0)} \Psi_n(y, x) |y|^{2\alpha} e^{\Phi(y, 0)} dy \\
&\quad + O\left(\int_{B(0, \overline{\theta})} \frac{|y|^3}{|x|^3} \frac{|y|^{2\alpha} e^{\lambda_n}}{(1 + e^{\lambda_n} |y|^{2+2\alpha})^2} dy\right) + o(e^{-\lambda_n}) \\
&= -b_0 \int_{\Omega \setminus B(0, \overline{\theta})} \frac{(8(1+\alpha)^2)^2 e^{-\lambda_n}}{\rho_n \overline{h}_1(0)} \Psi_n(y, x) |y|^{2\alpha} e^{\Phi(y, 0)} dy \\
&\quad + O\left(\frac{m_{n, \alpha}}{|x|^3}\right) + o(e^{-\lambda_n}) \text{ in } C^1(\partial B(0, r)),
\end{aligned} \tag{5.15}$$

where

$$m_{n, \alpha} = \begin{cases} \sigma_n^3, & \text{if } 2\alpha > 1, \\ \sigma_n^3 \log(\sigma_n^{-1}), & \text{if } 2\alpha = 1, \\ \sigma_n^{2+2\alpha} \overline{\theta}^{1-2\alpha}, & \text{if } 2\alpha < 1. \end{cases}$$

Let us set

$$\zeta_n^*(x) = -b_0 \int_{\Omega \setminus B(0, \overline{\theta})} \frac{(8(1+\alpha)^2)^2 e^{-\lambda_n}}{\rho_n \overline{h}_1(0)} \Psi_n(y, x) |y|^{2\alpha} e^{\Phi(y, 0)} dy \tag{5.16}$$

and then substitute (5.15) into (5.8), to derive that

$$\text{L.H.S. of (5.3)} = \int_{\partial B(0, r)} 4(1+\alpha) \langle \nu, D(\overline{G}_n + \zeta_n^*)(x) \rangle d\sigma + O(\sigma |A_n|) + O\left(\frac{m_{n, \alpha}}{r^3}\right) + o(\sigma_n^2). \tag{5.17}$$

To estimate the right hand side of (5.17), we notice that for any pair of (smooth enough) functions  $u$  and  $v$ , it holds

$$\begin{aligned}
&\Delta u (\nabla v \cdot x) + \Delta v (\nabla u \cdot x) \\
&= \operatorname{div} (\nabla u (\nabla v \cdot x) + \nabla v (\nabla u \cdot x) - \nabla u \cdot \nabla v(x)).
\end{aligned} \tag{5.18}$$

In view of (5.14), we also see that, for any  $\underline{\theta} \in (0, r)$ ,

$$\Delta \overline{G}_n(x) = A_n = \int_{\Omega} f_n^* dy = \int_{\Omega} \frac{\rho_n h(e^{\tilde{u}_n^{(1)}} - e^{\tilde{u}_n^{(2)}})}{\|\tilde{u}_n^{(1)} - \tilde{u}_n^{(2)}\|_{L^\infty(\Omega)}} = 0 \text{ for } x \in B(0, r) \setminus B(0, \underline{\theta}), \tag{5.19}$$

and moreover, by using (5.5) and (5.1), we have

$$\Delta(G_n - \varphi_n)(x) = 0 \quad \text{for } x \in B(0, r) \setminus B(0, \underline{\theta}). \tag{5.20}$$



By using (5.18)-(5.20) and (5.6), we conclude that

$$\begin{aligned}
0 &= \int_{B(0,r) \setminus B(0,\underline{\theta})} [\Delta \bar{G}_n (\nabla (G_n - \varphi_n) \cdot x) + \Delta (G_n - \varphi_n) (\nabla \bar{G}_n \cdot x)] dx \\
&= \int_{\partial(B(0,r) \setminus B(0,\underline{\theta}))} \left( \frac{\partial \bar{G}_n}{\partial \nu} (\nabla (G_n - \varphi_n) \cdot x) + \frac{\partial (G_n - \varphi_n)}{\partial \nu} (\nabla \bar{G}_n \cdot x) - \nabla \bar{G}_n \cdot \nabla (G_n - \varphi_n) \langle x, \nu \rangle \right) d\sigma \\
&= -\frac{\rho_n}{2\pi} \int_{\partial(B(0,r) \setminus B(0,\underline{\theta}))} \frac{\partial \bar{G}_n}{\partial \nu} d\sigma,
\end{aligned}$$

and thus,

$$\int_{\partial B(0,r)} \frac{\partial \bar{G}_n}{\partial \nu}(x) d\sigma = \int_{\partial B(0,\underline{\theta})} \frac{\partial \bar{G}_n}{\partial \nu}(x) d\sigma. \quad (5.21)$$

At this point, let us denote by  $o_{\underline{\theta}}(1)$  any quantity which converges to 0 as  $\underline{\theta} \rightarrow 0^+$ , and then observe that,

$$4(1+\alpha) \int_{\partial B(0,\underline{\theta})} \langle \nu, A_n D_x G(0, x) \rangle d\sigma = -4(1+\alpha) A_n + o_{\underline{\theta}}(1). \quad (5.22)$$

Since,  $D_i D_h \log |x| = \frac{\delta_{ih} |x|^2 - 2x_i x_h}{|x|^4}$ , then we find that,

$$\int_{\partial B(0,\underline{\theta})} \langle \nu, D_x \partial_{y_h} (\log |y-x|) |_{y=0} \rangle d\sigma = - \int_{\partial B(0,\underline{\theta})} \sum_{i=1}^2 \frac{x_i}{|x|} \left( \frac{\delta_{ih} |x|^2 - 2x_i x_h}{|x|^4} \right) d\sigma = 0. \quad (5.23)$$

We observe that, if  $h = k$  then  $D_i \log |x| = \frac{x_i}{|x|^2}$ ,

$$D_i D_{hh}^2 \log |x| = -\frac{2x_i}{|x|^4} - \frac{4x_h \delta_{ih}}{|x|^4} + \frac{8x_h^2 x_i}{|x|^6},$$

and thus,

$$\int_{\partial B(0,\underline{\theta})} \left\langle \nu, D_x \frac{\partial^2}{\partial y_h^2} \log \frac{1}{|y-x|} \Big|_{y=0} \right\rangle d\sigma = \int_{\partial B(0,\underline{\theta})} \left( \frac{2}{|x|^3} - \frac{4x_h^2}{|x|^5} \right) d\sigma = 0. \quad (5.24)$$

If  $h \neq k$ , then

$$D_i D_{hk}^2 \log |x| = -\frac{2(x_h \delta_{ki} + x_k \delta_{hi})}{|x|^4} + \frac{8x_k x_i x_h}{|x|^6},$$

which implies that

$$\int_{\partial B(0,\underline{\theta})} \left\langle \nu, D_x \frac{\partial^2}{\partial y_h \partial y_k} \log \frac{1}{|y-x|} \Big|_{y=0} \right\rangle d\sigma = \int_{\partial B(0,\underline{\theta})} \left( \frac{4x_h x_k}{|x|^5} - \frac{8x_h x_k}{|x|^5} \right) d\sigma = 0. \quad (5.25)$$

By (5.21)-(5.25), we conclude that

$$4(1+\alpha) \int_{\partial B(0,r)} \langle \nu, D_x \bar{G}_n(x) \rangle d\sigma = -4(1+\alpha) A_n + o_{\underline{\theta}}(1). \quad (5.26)$$

Next we estimate the other terms in (5.17), that is  $4(1+\alpha) \int_{\partial B(0,r)} \langle \nu, D_x \zeta_n^*(x) \rangle d\sigma$ , where  $\zeta_n^*$  is defined in (5.16). Clearly we have

$$\begin{aligned}
D_x \Psi_n(y, x) &= D_x \left( G(y, x) - G(0, x) - \langle \partial_y G(y, x) |_{y=0}, y \rangle 1_{B(0,r)}(y) \right) \\
&\quad - \frac{1}{2} D_x \left( \frac{1}{2} \langle \partial_y^2 G(y, x) |_{y=0}(y), y \rangle 1_{B(0,r)}(y) \right).
\end{aligned}$$

If  $y \in \Omega \setminus B(0, \bar{\theta})$  and  $x \in \partial B(0, \theta)$  with  $\theta \ll (\bar{\theta})^2$ , then we find that

$$|D_x G(x, y)| \leq \frac{C}{\sqrt{\theta}} \text{ for some constant } C > 0, \quad (5.27)$$

which implies

$$\int_{\partial B(0, \theta)} \langle v, D_x G(y, x) \rangle dx = o_\theta(1).$$

Thus (5.23)-(5.25) and (5.27) imply that

$$\begin{aligned} & 4(1 + \alpha) \int_{\partial B(0, \theta)} \langle v, D_x \Psi_n(y, x) \rangle dx \\ &= -4(1 + \alpha) \int_{\partial B(0, \theta)} \langle v, D_x G(0, x) \rangle dx \\ &\quad - 4(1 + \alpha) \int_{\partial B(0, \theta)} \left\langle v, D_x \left\langle \partial_y G(y, x) \mid_{y=0}, y \right\rangle 1_{B(0, r_0)}(y) \right\rangle dx \\ &\quad - 2(1 + \alpha) \int_{\partial B(0, \theta)} \left\langle v, D_x \left\langle \partial_y^2 G(y, x) \mid_{y=0}, y, y \right\rangle 1_{B(0, r_0)}(y) \right\rangle dx + o_\theta(1) \\ &= 4(1 + \alpha) + o_\theta(1) \quad \text{for } y \in \Omega \setminus B(0, \bar{\theta}), \quad \text{and } x \in \partial B(0, \theta). \end{aligned} \quad (5.28)$$

We observe that

$$-\Delta_x \Psi_n(y, x) = \delta_y \quad \text{for } x \in B(0, r) \setminus B(0, \theta)$$

and let us choose  $u(x) = \Psi_n(y, x)$  and  $v(x) = G_n(x) - \varphi_n(x)$  in (5.18). Then we consider the following two cases:

(i) If  $y \in \partial B(0, r) \setminus B(0, \bar{\theta})$ , then from (5.18) and (5.28), we obtain that

$$\begin{aligned} & 4(1 + \alpha) \int_{\partial B(0, r)} \langle v, D_x \Psi_n(y, x) \rangle dx \\ &= 4(1 + \alpha) \int_{\partial B(0, \theta)} \langle v, D_x \Psi_n(y, x) \rangle dx - 4(1 + \alpha) = o_\theta(1). \end{aligned} \quad (5.29)$$

(ii) If  $y \in \Omega \setminus B(0, r)$ , then we see from (5.18) and (5.28) that

$$\begin{aligned} & 4(1 + \alpha) \int_{\partial B(0, r)} \langle v, D_x \Psi_n(y, x) \rangle dx = 4(1 + \alpha) \int_{\partial B(0, \theta)} \langle v, D_x \Psi_n(y, x) \rangle dx \\ &= 4(1 + \alpha) + o_\theta(1). \end{aligned} \quad (5.30)$$

and by (5.16), and (5.29)-(5.30), we finally conclude that

$$\begin{aligned} & 4(1 + \alpha) \int_{\partial B(0, r)} \langle v, D_x \tilde{\zeta}_n^*(x) \rangle dx \\ &= -\frac{(8(1 + \alpha)^2)^3 b_0 e^{-\lambda_n}}{2\rho_n \bar{h}_1(0)} \int_{\Omega \setminus B(0, \bar{\theta})} \left( \int_{\partial B(0, r)} \langle v, D_x \Psi_n \rangle \right) |y|^{2\alpha} e^{\Phi(y, 0)} dx dy \\ &= -\frac{(8(1 + \alpha)^2)^3 b_0 e^{-\lambda_n}}{2\rho_n \bar{h}_1(0)} \int_{\Omega \setminus B(0, r)} |y|^{2\alpha} e^{\Phi(y, 0)} dy + o(e^{-\lambda_n}). \end{aligned} \quad (5.31)$$

Obviously from (5.17), (5.26) and (5.31) we get the conclusion of Lemma 5.2  $\square$

To estimate the right hand side of (5.3) of Lemma 5.1, we recall, see for example (5.10), that

$$f_n^*(x) = \rho_n h(x) e^{\tilde{u}_n^{(1)}} (\tilde{\zeta}_n + o(1)).$$

Recall also the definitions of  $\Phi(x, 0)$  and  $\mathcal{H}_p = \mathcal{H}_0$  in (5.4) and (1.4), respectively and the definition of  $\ell(p)$  after (2.4). A crucial point in our proof is the following estimate.

**Lemma 5.3.** (i)

$$\int_{\partial B(0,r)} r f_n^* d\sigma = -\frac{128(1+\alpha)^4 b_0 \pi e^{-\lambda_n}}{\rho_n \bar{h}_1(0) r^{2+2\alpha}} - \frac{32(1+\alpha)^4 b_0 \pi e^{-\lambda_n}}{\rho_n \bar{h}_1(0) r^{2\alpha}} \Delta \log h_*(0) + O(r^{1-2\alpha} e^{-\lambda_n}) + \frac{o(e^{-\lambda_n})}{r^{2+2\alpha}},$$

(ii)

$$\int_{B(0,r)} f_n^*(x) dx = \frac{64(1+\alpha)^4 b_0 e^{-\lambda_n}}{\rho_n \bar{h}_1(0)} \int_{\Omega \setminus B(0,r)} |x|^{2\alpha} e^{\Phi(x,0)} dx + \frac{o(e^{-\lambda_n})}{r^{2+2\alpha}},$$

(iii)

$$\begin{aligned} & \int_{B(0,r)} f_n^* \langle D(\log \bar{h}_1 + \varphi_n), x \rangle dx \\ &= -2b_0 \ell(p) \sigma_n^2 + o(\sigma_n) |\nabla \mathcal{H}_0(0)| + O(m_{n,1}(\alpha)) \\ & \quad + O(\sigma_n^{2\epsilon_0} + \lambda_n \sigma_n^2) (|\nabla \mathcal{H}_0(0)| \sigma_n + \sigma_n^2) + O(\sigma_n^{2+\epsilon_0}) \\ & \quad + \left( O(R^{-2\alpha}) + O(\lambda_n) |A_n| + O\left(\frac{1}{R}\right) \right) (|\nabla \mathcal{H}_0(0)| \sigma_n + \sigma_n^2). \end{aligned}$$

where  $O(m_{n,1}(\alpha))$  is defined after (5.36) and  $O(1)$  is used to denote any quantity uniformly bounded with respect to  $r$ ,  $R$  and  $n$ .

*Proof.* (i) We first observe that (5.11) implies that

$$\int_{\partial B(0,r)} r f_n^*(x) d\sigma = \int_{\partial B(0,r)} \frac{(8(1+\alpha)^2)^2 e^{-\lambda_n} (-b_0 + o(1)) |x|^{2\alpha} e^{\mathcal{H}_0(x)}}{\rho_n \bar{h}_1(0) |x|^{3+4\alpha}} d\sigma. \quad (5.32)$$

Clearly we have

$$\mathcal{H}_0(x) = \langle D\mathcal{H}_0(0), x \rangle + \frac{1}{2} \langle D_x^2 \mathcal{H}_0|_{x=0} x, x \rangle + O(|x|^3). \quad (5.33)$$

By (5.32) and (5.33), we obtain,

$$\begin{aligned} & \int_{\partial B(0,r)} r f_n^*(x) d\sigma \\ &= \int_{\partial B(0,r)} \frac{(8(1+\alpha)^2)^2 e^{-\lambda_n}}{\rho_n \bar{h}_1(0) |x|^{3+3\alpha}} \left( b_0 (1 + \langle D\mathcal{H}_0, x \rangle) + \frac{1}{2} \langle D_x^2 \mathcal{H}_0|_{x=0} x, x \rangle + O(|x|^3) + o(1) \right) d\sigma \\ &= - \int_{\partial B(0,r)} \frac{(8(1+\alpha)^2)^2 e^{-\lambda_n} b_0 (1 + \frac{\Delta \mathcal{H}_0}{4} |x|^2)}{\rho_n \bar{h}_1(0) |x|^{3+2\alpha}} d\sigma + O(r^{1-2\alpha} e^{-\lambda_n}) + \frac{o(e^{-\lambda_n})}{r^{2+2\alpha}} \\ &= -\frac{128(1+\alpha)^4 b_0 \pi e^{-\lambda_n}}{\rho_n \bar{h}_1(0) r^{2+2\alpha}} - \frac{32(1+\alpha)^4 b_0 \pi e^{-\lambda_n}}{\rho_n \bar{h}_1(0) r^{2\alpha}} \Delta \log(h_*(0)) + O(r^{1-2\alpha} e^{-\lambda_n}) + \frac{o(e^{-\lambda_n})}{r^{2+2\alpha}}, \end{aligned}$$

which proves (i).

(ii) We notice that  $A_n = \int_{\Omega} f_n^* = 0$ , and thus

$$\int_{B(0,r)} f_n^*(x) dx = - \int_{\Omega \setminus B(0,r)} f_n^*(x) dx. \quad (5.34)$$

By (5.11) we see that

$$\begin{aligned} - \int_{\Omega \setminus B(0,r)} f_n^* dx &= \int_{\Omega \setminus B(0,r)} \frac{64(1+\alpha)^4 b_0 e^{-\lambda_n}}{\rho_n \bar{h}_1(0)} |x|^{2\alpha} e^{\Phi(x,0)} dx + \frac{o(e^{-\lambda_n})}{r^{2+2\alpha}} \\ &= \frac{64(1+\alpha)^4 b_0 e^{-\lambda_n}}{\rho_n \bar{h}_1(0)} \int_{\Omega \setminus B(0,r)} |x|^{2\alpha} e^{\Phi(x,0)} dx + \frac{o(e^{-\lambda_n})}{r^{2+2\alpha}}, \end{aligned} \quad (5.35)$$

which proves (ii).

(iii) By (2.3) and (2.6), we see that

$$\tilde{u}_n(x) = U_n(x) + 8\pi(1+\alpha)(R(x,0) - R(0,0)) + \eta_n(x), \quad x \in B(0,r),$$

where

$$\eta_n(x) = \sigma_n \psi_{n,1}(\sigma_n^{-1}x) + \sigma_n^2 \psi_{n,2}(\sigma_n^{-1}x) + O(\sigma_n^2),$$

see (2.7), (2.8) and (2.10). Thus, we set

$$\omega_n(r) = \|\tilde{u}_n^{(1)} - \tilde{u}_n^{(2)}\|_{L^\infty(B(0,r))},$$

and use Lemma 3.1 and (2.4), we deduce that

$$\begin{aligned} &\int_{B(0,r)} f_n^* \langle D(\log \bar{h}_1(x) + \varphi_n), x \rangle dx \\ &= \int_{B(0,r)} \frac{\rho_n \bar{h}_1(0) |x|^{2\alpha} e^{\lambda_n + \mathcal{H}_0(x) + \eta_n(x)}}{(1 + \gamma_n e^{\lambda_n} |x|^{2+2\alpha})^2} (\xi_n - \frac{\omega_n(r)}{2} \xi_n^2 + O(\omega_n^2(r))) \langle D\mathcal{H}_0(x), x \rangle dx \\ &= \int_{B(0,r)} \frac{\rho_n \bar{h}_1(0) |x|^{2\alpha} e^{\lambda_n + \mathcal{H}_0(x) + \eta_n(x)}}{(1 + \gamma_n e^{\lambda_n} |x|^{2+2\alpha})^2} (\xi_n - \frac{\omega_n(r)}{2} \xi_n^2 + O(\omega_n^2(r))) \\ &\quad \times \langle D\mathcal{H}_0(0) + D^2\mathcal{H}_0(0)x + O(|x|^2), x \rangle dx \\ &= \int_{B(0,\sigma_n^{-1}r)} \frac{\rho_n \bar{h}_1(0) |z|^{2\alpha}}{(1 + \gamma_n |z|^{2+2\alpha})^2} \left( \hat{\xi}_n - \frac{\omega_n}{2} \hat{\xi}_n^2 + \langle D\mathcal{H}_0(0), \sigma_n z \rangle + \eta_n + O(\sigma_n^2 |z|^2) + O(\omega_n^2) \right) \\ &\quad \times \langle D\mathcal{H}_0(0) + D^2\mathcal{H}_0(0) \cdot \sigma_n z + O(\sigma_n^2 |z|^2), \sigma_n z \rangle dz =: K_{n,r}. \end{aligned} \quad (5.36)$$

Set

$$\begin{aligned} m_{n,1}(\alpha) &= \begin{cases} \sigma_n^3 & \text{if } \alpha > \frac{1}{2} \\ \log(r\sigma_n^{-1})\sigma_n^3 & \text{if } \alpha = \frac{1}{2} \\ r^{1-2\alpha}\sigma_n^{2(1+\alpha)} & \text{if } \alpha \in (0, \frac{1}{2}) \end{cases}, \quad m_{n,2}(\alpha) = \begin{cases} \sigma_n^4 & \text{if } \alpha > 1 \\ \log(r\sigma_n^{-1})\sigma_n^4 & \text{if } \alpha = 1 \\ r^{2-2\alpha}\sigma_n^{2(1+\alpha)} & \text{if } \alpha \in (0, 1) \end{cases}, \\ m_{n,3}(\alpha) &= \begin{cases} \sigma_n^5 & \text{if } \alpha > \frac{3}{2} \\ \log(r\sigma_n^{-1})\sigma_n^5 & \text{if } \alpha = \frac{3}{2} \\ r^{3-2\alpha}\sigma_n^{2(1+\alpha)} & \text{if } \alpha \in (0, \frac{3}{2}) \end{cases}. \end{aligned}$$

Using (5.36) together with (2.5), (2.7) and Lemma 3.1, we conclude that

$$\begin{aligned} K_{n,r} &= \int_{B(0,\sigma_n^{-1}r)} \frac{\rho_n \bar{h}_1(0) |z|^{2\alpha}}{(1 + \gamma_n |z|^{2(1+\alpha)})^2} \\ &\quad \times \left( \hat{\xi}_n - \frac{\omega_n(r)}{2} (\hat{\xi}_n)^2 + \sigma_n (\psi_{n,1}(z) + \langle D\mathcal{H}_0(0), z \rangle) + O(\sigma_n^2 |z|^2) + O((\sigma_n^{2\epsilon_0} + \lambda_n \sigma_n^2)^2) + \sigma_n^2 \tilde{\psi}_{n,2}(z) \right) \\ &\quad \times \langle D\mathcal{H}_0(0) + D^2\mathcal{H}_0(0) \cdot \sigma_n z + O(\sigma_n^2 |z|^2), \sigma_n z \rangle dz \end{aligned}$$

$$\begin{aligned}
&= \int_{B(0, r\sigma_n^{-1})} \frac{\rho_n \bar{h}_1(0) |z|^{2\alpha} \left( \hat{\xi}_n - \frac{\omega_n(r)}{2} (\hat{\xi}_n)^2 + \sigma_n (\psi_{n,1}(z) + \langle D\mathcal{H}_0(0), z \rangle) \right)}{(1 + \gamma_n |z|^{2(1+\alpha)})^2} \\
&\times \langle D\mathcal{H}_0(0) + \sigma_n D^2\mathcal{H}_0(0) \cdot z, z \rangle \sigma_n dz + O(m_{n,1}(\alpha) + m_{n,2}(\alpha) + m_{n,3}(\alpha)) \\
&+ \left( |\nabla \mathcal{H}_0(0)| \sigma_n + \sigma_n^2 \right) O((\sigma_n^{2\epsilon_0} + \lambda_n \sigma_n^2)^2) \\
&= I_{n,1} + I_{n,2} + O(m_{n,1}(\alpha)) + \left( |\nabla \mathcal{H}_0(0)| \sigma_n + \sigma_n^2 \right) O(\sigma_n^{4\epsilon_0}),
\end{aligned}$$

where

$$I_{n,1} = \int_{B(0, r\sigma_n^{-1})} \frac{\rho_n \bar{h}_1(0) |z|^{2\alpha} \left( \hat{\xi}_n - \frac{\omega_n}{2} \hat{\xi}_n^2 \right)}{(1 + \gamma_n |z|^{2+2\alpha})^2} \langle D\mathcal{H}_0(0) + \sigma_n D^2\mathcal{H}_0(0) \cdot z, z \rangle \sigma_n dz,$$

and

$$I_{n,2} = \int_{B(0, r\sigma_n^{-1})} \frac{\rho_n \bar{h}_1(0) |z|^{2\alpha} (\psi_{n,1}(z) + \langle D\mathcal{H}_0(0), z \rangle)}{(1 + \gamma_n |z|^{2+2\alpha})^2} \langle D\mathcal{H}_0(0), z \rangle \sigma_n^2 dz.$$

In view of (2.9), (2.10), (2.11) and (2.4), we have

$$\langle D\mathcal{H}_0(0), z \rangle = \partial_{x_1} \mathcal{H}_0(0) z_1 + O(\sigma_n^2)(z_1 + z_2) = a_{n,1} z_1 + O(\sigma_n^2)(z_1 + z_2),$$

and then, putting  $a_1 = \partial_{x_1} \mathcal{H}_0$  and  $\Lambda(z) = \rho_n \bar{h}_1(0) |z|^{2\alpha}$ , we conclude that

$$\begin{aligned}
\sigma_n^{-2} I_{n,2} &= \int_{B(0, r\sigma_n^{-1})} \frac{\Lambda(z) (\psi_{n,1}(z) + a_1 z_1 + O(\sigma_n^2)(z_1 + z_2))}{(1 + \gamma_n |z|^{2+2\alpha})^2} (a_1 z_1 + O(\sigma_n^2)(z_1 + z_2)) dz \\
&= \int_{B(0, r\sigma_n^{-1})} \frac{\Lambda(z)}{(1 + \gamma_n |z|^{2+2\alpha})^2} \left( -\frac{2(1+\alpha)a_1 z_1}{\alpha(1 + \gamma_n |z|^{2+2\alpha})} + a_1 z_1 + O(\sigma_n^2)(z_1 + z_2) \right) \\
&\quad \times (a_1 z_1 + O(\sigma_n^2)(z_1 + z_2)) dz \\
&= - \int_{B(0, r\sigma_n^{-1})} \frac{2(1+\alpha)\Lambda(z)a_1^2 z_1^2}{\alpha(1 + \gamma_n |z|^{2+2\alpha})^3} dz + \int_{B(0, r\sigma_n^{-1})} \frac{\Lambda(z)a_1^2 z_1^2}{(1 + \gamma_n |z|^{2+2\alpha})^2} dz + O(\sigma_n^2) \\
&= - \rho_n \bar{h}_1(0) \frac{a_1^2 \pi^2}{2(1+\alpha)^2 \gamma_n^{\frac{2+\alpha}{1+\alpha}} \sin \frac{\pi}{1+\alpha}} + \rho_n \bar{h}_1(0) \frac{a_1^2 \pi}{2\alpha \gamma_n^{\frac{2+\alpha}{1+\alpha}}} \Gamma\left(\frac{2+\alpha}{1+\alpha}\right) \Gamma\left(\frac{1+2\alpha}{1+\alpha}\right) \\
&\quad + O(\sigma_n^{2\alpha}) + O(\sigma_n^2) \\
&= O(\sigma_n^{2\alpha}) + O(\sigma_n^2),
\end{aligned}$$

where we used the properties of  $\Gamma(x)$ , and thus

$$I_{n,2} = O(\sigma_n^{2+\epsilon_0}). \quad (5.37)$$

On the other hand, in view of Lemma 4.1, for any fixed  $R \geq 1$  large, we have

$$\begin{aligned}
& \int_{B(0,R)} \frac{\Lambda(z)(\hat{\xi}_n - \frac{\omega_n}{2}\hat{\xi}_n^2)}{(1 + \gamma_n|z|^{2+2\alpha})^2} \left\langle D\mathcal{H}_0(0) + \sigma_n D^2\mathcal{H}_0(0) \cdot z, z \right\rangle \sigma_n dz \\
&= \int_{B(0,R)} \frac{\Lambda(z)(b_0\hat{\xi}_0(z) + o(1) + O(\sigma_n^{2\epsilon_0} + \lambda_n\sigma_n^2))}{(1 + \gamma_n|z|^{2+2\alpha})^2} \left\langle D\mathcal{H}_0(0) + \sigma_n D^2\mathcal{H}_0(0) \cdot z, z \right\rangle \sigma_n dz \\
&= \sigma_n^2 \int_{B(0,R)} \frac{\Lambda(z)(b_0\hat{\xi}_0(z) + O(\sigma_n^{2\epsilon_0} + \lambda_n\sigma_n^2))}{(1 + \gamma_n|z|^{2+2\alpha})^2} \left\langle D^2\mathcal{H}_0(0) \cdot z, z \right\rangle dz \\
&\quad + o(\sigma_n)|\nabla\mathcal{H}_0(0)| + O(\sigma_n^{2\epsilon_0} + \lambda_n\sigma_n^2) \left( |\nabla\mathcal{H}_0(0)|\sigma_n + \sigma_n^2 \right) \\
&= \sigma_n^2 \int_{B(0,R)} \frac{\Lambda(z)(b_0\hat{\xi}_0(z))}{(1 + \gamma_n|z|^{2+2\alpha})^2} \left\langle D^2\mathcal{H}_0(0) \cdot z, z \right\rangle dz + o(\sigma_n)|\nabla\mathcal{H}_0(0)| \\
&\quad + O(\sigma_n^{2\epsilon_0} + \lambda_n\sigma_n^2) \left( |\nabla\mathcal{H}_0(0)|\sigma_n + \sigma_n^2 \right) \\
&= 8\pi(1 + \alpha)\bar{h}_1(0)b_0\sigma_n^2 \int_{B(0,R)} \frac{|z|^{2\alpha}\hat{\xi}_0(z)}{(1 + \gamma|z|^{2+2\alpha})^2} \left\langle D^2\mathcal{H}_0(0) \cdot z, z \right\rangle dz \\
&\quad + o(\sigma_n)|\nabla\mathcal{H}_0(0)| + O(\sigma_n^{2\epsilon_0} + \lambda_n\sigma_n^2) \left( |\nabla\mathcal{H}_0(0)|\sigma_n + \sigma_n^2 \right).
\end{aligned}$$

Finally we have

$$\begin{aligned}
& \int_{B(0,R)} \frac{|z|^{2\alpha}\hat{\xi}_0(z)}{(1 + \gamma|z|^{2+2\alpha})^2} \left\langle D^2\mathcal{H}_0(0) \cdot z, z \right\rangle dz = \frac{\Delta\mathcal{H}_0(0)}{2} \int_{B(0,R)} \frac{1 - \gamma|z|^{2+2\alpha}}{(1 + \gamma|z|^{2+2\alpha})^3} |z|^{2\alpha+2} dz \\
&= -\frac{\Delta\mathcal{H}_0(0)}{2} \frac{\pi^2}{(1 + \alpha)^3 \gamma^{\frac{2+\alpha}{1+\alpha}} \sin \frac{\pi}{1+\alpha}} + O(R^{-2\alpha}) \\
&= -\frac{\pi}{2(1 + \alpha)^2 \bar{h}_1(0) \gamma^{\frac{1}{1+\alpha}} \sin \frac{\pi}{1+\alpha}} \Delta \log(h_*(0)) + O(R^{-2\alpha}).
\end{aligned}$$

On the other side, in view of (4.9), we also see that if  $R \leq |z| \leq r/\sigma_n$ , then it holds

$$\hat{\xi}_n(z) = -d_n + O(\lambda_n)|A_n| + O\left(\frac{1}{|z|}\right), \quad (5.38)$$

and thus

$$\hat{\xi}_n(z)^2 = d_n^2 + O(\lambda_n^2 + \frac{\lambda_n}{|z|})|A_n| + O\left(\frac{1}{|z|^2}\right).$$

As a consequence, by Lemma 3.1, we find that

$$\begin{aligned}
& \int_{B(0,r/\sigma_n) \setminus B(0,R)} \frac{\rho_n \bar{h}_1(0) |z|^{2\alpha} (\hat{\xi}_n - \frac{\omega_n}{2} \hat{\xi}_n^2)}{(1 + \gamma_n |z|^{2+2\alpha})^2} \left\langle D\mathcal{H}_0(0) + \sigma_n D^2\mathcal{H}_0(0) \cdot z, z \right\rangle \sigma_n dz \\
&= \left( -d_n - \frac{\omega_n(r)}{2} d_n^2 \right) \int_{B(0,r/\sigma_n) \setminus B(0,R)} \frac{\rho_n \bar{h}_1(0) |z|^{2\alpha}}{(1 + \gamma_n |z|^{2+2\alpha})^2} \left\langle D\mathcal{H}_0(0) + \sigma_n D^2\mathcal{H}_0(0) \cdot z, z \right\rangle \sigma_n dz \\
&\quad + \int_{B(0,r/\sigma_n) \setminus B(0,R)} \frac{\rho_n \bar{h}_1(0) |z|^{2\alpha} (O(\lambda_n) |A_n| + O(\frac{1}{|z|}))}{(1 + \gamma_n |z|^{2+2\alpha})^2} \left\langle D\mathcal{H}_0(0) + \sigma_n D^2\mathcal{H}_0(0) \cdot z, z \right\rangle \sigma_n dz \\
&= -b_0 \Delta \log h_*(0) \int_{B(0,r/\sigma_n) \setminus B(0,R)} \frac{\rho_n \bar{h}_1(0) |z|^{2\alpha+2}}{(1 + \gamma_n |z|^{2+2\alpha})^2} \sigma_n^2 dz + O(\sigma_n^2 (\sigma_n^{2\epsilon_0} + \lambda_n \sigma_n^2)) \\
&\quad + \left( O(|\lambda_n|) |A_n| + O\left(\frac{1}{R}\right) \right) (|\nabla \mathcal{H}_0(0)| \sigma_n + \sigma_n^2) \\
&= O(R^{-2\alpha}) \sigma_n^2 + \left( O(R^{-2\alpha}) + O(\lambda_n) |A_n| + O\left(\frac{1}{R}\right) \right) (|\nabla \mathcal{H}_0(0)| \sigma_n + \sigma_n^2) \\
&\quad + O(\sigma_n^{2+2\epsilon_0} + \lambda_n \sigma_n^4)
\end{aligned}$$

Collecting the above estimates we conclude that

$$\begin{aligned}
& \int_{B(0,r)} f_n^* \left\langle D(\log \bar{h}_1 + \varphi_n), x \right\rangle dx \\
&= -2b_0 \ell(p) \sigma_n^2 + o(\sigma_n) |\nabla \mathcal{H}_0(0)| + O(m_{n,1}(\alpha)) + O(\sigma_n^{2\epsilon_0} + \lambda_n \sigma_n^2) (|\nabla \mathcal{H}_0(0)| \sigma_n + \sigma_n^2) \\
&\quad + O(\sigma_n^{2+\epsilon_0}) + \left( O(R^{-2\alpha}) + O(\lambda_n) |A_n| + O\left(\frac{1}{R}\right) \right) (|\nabla \mathcal{H}_0(0)| \sigma_n + \sigma_n^2).
\end{aligned}$$

□

Recall that  $p = 0$ . Using the assumptions  $\ell(p) \neq 0$  and  $\nabla \mathcal{H}_0(0) = 0$  we can now prove that  $b_0 = 0$ .

**Lemma 5.4.**  $b_0 = 0$ .

*Proof.* By (5.3) and Lemmas 5.2-5.3, we have for any  $r \in (0, 1)$  and  $R > 1$ ,

$$\begin{aligned}
& -4(1 + \alpha) A_n - \frac{(8(1 + \alpha)^2)^3 b_0 e^{-\lambda_n}}{2\rho_n \bar{h}_1(0)} \int_{\Omega \setminus B(0,r)} |y|^{2\alpha} e^{\Phi(y,0)} dy + o(\sigma_n^2) + O(\sigma_n |A_n| + \frac{\sigma_n^3}{r^3}) \\
&= -\frac{128(1 + \alpha)^4 b_0 \pi e^{-\lambda_n}}{\rho_n \bar{h}_1(0) r^{2+2\alpha}} - \frac{32(1 + \alpha)^4 b_0 \pi e^{-\lambda_n}}{\rho_n \bar{h}_1(0) r^{2\alpha}} \Delta \log h_*(0) + O(r^{1-2\alpha} e^{-\lambda_n}) \\
&\quad - \frac{128(1 + \alpha)^5 b_0 e^{-\lambda_n}}{\rho_n \bar{h}_1(0)} \int_{\Omega \setminus B(0,r)} |y|^{2\alpha} e^{\Phi(y,0)} dy + \frac{o(e^{-\lambda_n})}{r^{2+2\alpha}} + 2b_0 \ell(p) \sigma_n^2 \\
&\quad + o(\sigma_n) |\nabla \mathcal{H}_0(0)| + O(m_{n,1}(\alpha)) + O(\sigma_n^{2\epsilon_0} + \lambda_n \sigma_n^2) (|\nabla \mathcal{H}_0(0)| \sigma_n + \sigma_n^2) \\
&\quad + O(\sigma_n^{2+\epsilon_0}) + \left( O(R^{-2\alpha}) + O(\lambda_n) |A_n| + O\left(\frac{1}{R}\right) \right) (|\nabla \mathcal{H}_0(0)| \sigma_n + \sigma_n^2).
\end{aligned}$$

Recall  $A_n = 0$ . Since  $\nabla \mathcal{H}_0(0) = 0$  by assumption, after some manipulations, for  $r \in (0, r_0)$  and any  $R > 1$ , we find that

$$\begin{aligned} b_0 \ell(p) \sigma_n^2 &= o(\sigma_n^2) + O(m_{n,1}(\alpha)) + O(R^{-2\alpha} + R^{-1}) \sigma_n^2 + O(\sigma_n^{2+2\epsilon_0} + \lambda_n \sigma_n^4) \\ &\quad + O\left(\frac{\sigma_n^3}{r^3}\right) + \frac{o(e^{-\lambda_n})}{r^{2+2\alpha}}, \end{aligned}$$

which implies

$$b_0 = 0.$$

provided  $\ell(p) \neq 0$ . Hence we finish the proof.  $\square$

*Proof of Theorem 1.1.* Let  $x_n^*$  be a maximum point of  $\tilde{\zeta}_n$ , then we have,

$$|\tilde{\zeta}_n(x_n^*)| = 1. \quad (5.39)$$

By Lemma 4.2 and Lemma 5.4 we have that  $x_n^* \rightarrow p$ . By Lemma 5.4, it holds that

$$\lim_{n \rightarrow +\infty} e^{\frac{\lambda_n^{(1)}}{2(1+\alpha)}} s_n = +\infty, \quad \text{where } s_n = |x_n^* - p|. \quad (5.40)$$

Setting  $\bar{\zeta}_n(x) = \tilde{\zeta}(s_n x + p)$ , then we have  $\bar{\zeta}_n$  satisfies

$$\begin{aligned} 0 &= \Delta \bar{\zeta}_n + \rho_n s_n^2 h(s_n x + p) c_n(s_n x + p) \bar{\zeta}_n \\ &= \Delta \bar{\zeta}_n + \frac{\rho_n \bar{h}_1(p) |x|^{2\alpha} s_n^{2+2\alpha} e^{\lambda_n^{(1)}} (1 + O(s_n |x|) + o(1)) \bar{\zeta}_n}{(1 + \frac{\rho_n \bar{h}_1(p)}{8(1+\alpha)^2} e^{\lambda_n^{(1)}} |s_n x|^{2+2\alpha})^2}. \end{aligned}$$

On the other hand, by (5.39), we also have

$$\left| \bar{\zeta}_n \left( \frac{x_n^* - p}{s_n} \right) \right| = |\tilde{\zeta}_n(x_n^*)| = 1. \quad (5.41)$$

In view of (5.40) and  $|\bar{\zeta}_n| \leq 1$  we see that  $\bar{\zeta}_n \rightarrow \bar{\zeta}_0$  on any compact subset of  $\mathbb{R}^2 \setminus \{0\}$ , where  $\bar{\zeta}_0$  satisfies  $\Delta \bar{\zeta}_0 = 0$  in  $\mathbb{R}^2 \setminus \{0\}$ . Since  $|\bar{\zeta}_0| \leq 1$ , we have  $\Delta \bar{\zeta}_0 = 0$  in  $\mathbb{R}^2$ , which implies  $\bar{\zeta}_0$  is a constant. At this point, since  $\frac{|x_n^* - p|}{s_n} = 1$  and in view of (5.41), we find  $\bar{\zeta}_0 = 1$  or  $\bar{\zeta}_0 = -1$ . From which we have  $|\bar{\zeta}_n(x)| \geq \frac{1}{2}$

when  $s_n \leq |x - p| \leq \frac{1}{2} s_n$ , which contradicts to (4.6)-(4.8) since  $e^{-\frac{\lambda_n^{(1)}}{2(1+\alpha)}} \ll s_n$  and  $\lim_{n \rightarrow +\infty} s_n = 0$  and  $b_0 = 0$ . This fact concludes the proof of Theorem 1.1.  $\square$

## 6. THE PROOF OF THEOREM 1.2

In this section we give the proof of the non-degeneracy result stated in Theorem 1.2. Since the argument is similar to the one yielding local uniqueness of bubbling solutions we will be sketchy to avoid repetitions, referring to [9] for full details.

Suppose by contradiction the linearized problem (1.2) admits a non-trivial solution  $\phi_n$ , where  $u_n$  is a singular 1-bubbling solution of  $(\mathbf{P}_{\rho_n})$  blowing up at the point



$p_i$  for some  $i \in \{1, \dots, N\}$ . We suppose with no loss of generality that  $p_i = 0 \in \Omega$ , set  $\alpha_i = \alpha$  and

$$\tilde{u}_n = u_n - \log \left( \int_{\Omega} h e^{u_n} dx \right), \quad \lambda_n = \max_{\Omega} \tilde{u}_n, \quad \sigma_n^{2(1+\alpha)} = e^{-\lambda_n},$$

Define

$$\Xi_n = \frac{\phi_n - \frac{\int_{\Omega} h e^{u_n} \phi_n dx}{\int_{\Omega} h e^{u_n} dx}}{\left\| \phi_n - \frac{\int_{\Omega} h e^{u_n} \phi_n dx}{\int_{\Omega} h e^{u_n} dx} \right\|_{L^\infty(\Omega)}},$$

which plays the role of the difference of two bubbling solutions, see (4.1) in the proof of Theorem 1.1. Then,  $\Xi_n$  satisfies

$$\begin{cases} \Delta \Xi_n + \rho_n h(x) c_n(x) \Xi_n(x) = 0 & \text{in } \Omega, \\ \Xi_n = -d_n & \text{on } \partial\Omega, \end{cases} \quad (6.1)$$

for some constant  $d_n$  satisfying  $|d_n| \leq 1$  and  $c_n(x) = e^{\tilde{u}_n(x)}$ .

**Step 1.** We start by considering the asymptotic behavior of  $\Xi_n$  near the blow up point  $p_i$ . After a suitable scaling,  $\Xi_n$  converges in  $C_{\text{loc}}^0(\mathbb{R}^2)$  to a solution  $\hat{\xi}$  of the linearized problem

$$\Delta \hat{\xi} + \frac{8\gamma(1+\alpha)^2 |z|^{2\alpha}}{(1+\gamma|z|^{2(1+\alpha)})^2} \hat{\xi} = 0 \text{ in } \mathbb{R}^2 \quad \text{and} \quad |\hat{\xi}(z)| \leq 1 \text{ in } \mathbb{R}^2,$$

where  $\gamma = \frac{\pi \bar{h}_1(0)}{1+\alpha}$ , see for example Lemma 4.1. It follows from [27, Corollary 2.2] that there exists a constant  $b_0 \in \mathbb{R}$  such that

$$\Xi_n(\sigma_n z) \rightarrow b_0 \frac{1 - \gamma |z|^{2+2\alpha}}{1 + \gamma |z|^{2+2\alpha}} \quad \text{in } C_{\text{loc}}^0(\mathbb{R}^2). \quad (6.2)$$

**Step 2.** We next consider the global behavior of  $\Xi_n$  away from the blow up point  $p_i$ . It follows from (2.2) that

$$c_n(x) \rightarrow 0 \quad \text{in } C_{\text{loc}}^0(\overline{\Omega} \setminus \{0\}).$$

Using then  $\|\Xi_n\|_{L^\infty(\Omega)} \leq 1$  and (6.1) it is not difficult to see that

$$\Xi_n \rightarrow \xi_0 \quad \text{in } C_{\text{loc}}^0(\overline{\Omega} \setminus \{0\}), \quad \Delta \xi_0 = 0 \quad \text{in } \Omega.$$

Therefore  $\xi_0 = -b$  in  $\Omega$  for some constant  $b$  and

$$\Xi_n \rightarrow -b \quad \text{in } C_{\text{loc}}^0(\overline{\Omega} \setminus \{0\}). \quad (6.3)$$

Finally, by an O.D.E. argument as in Lemma 4.2 one can show  $b = b_0$ .

**Step 3.** We then study the asymptotic in the Pohozaev-type identity given by Lemma 5.1 (with suitable minor modifications, see for example [9]). Using the assumption  $\nabla \mathcal{H}_{p_i}(p_i) = 0$  it is possible to prove that

$$b_0 \ell(p_i) = o(1) \quad \text{for } n \text{ large,}$$

see section 5. Since by assumption  $\ell(p_i) \neq 0$  we deduce  $b_0 = 0$ .

**Step 4.** The contradiction is then obtained by a blow up argument using  $b = b_0 = 0$  jointly with (6.2) and (6.3) exactly as in the proof of Theorem 1.1, see the end of section 5. The proof of Theorem 1.2 is completed.

## REFERENCES

- [1] D. Bartolucci, *Global bifurcation analysis of mean field equations and the Onsager microcanonical description of two-dimensional turbulence*, Calc. Var. P.D.E. (2019), 58:18.
- [2] D. Bartolucci, C.C. Chen, C.S. Lin, G. Tarantello, *Profile of Blow Up Solutions To Mean Field Equations with Singular Data*, Comm. in P. D. E. **29**(7-8) (2004), 1241-1265.
- [3] D. Bartolucci, F. De Marchis, *On the Ambjorn-Olesen electroweak condensates*, Jour. Math. Phys. **53** 073704 (2012).
- [4] D. Bartolucci, F. De Marchis, *Supercritical Mean Field Equations on convex domains and the Onsager's statistical description of two-dimensional turbulence*, Archive for Rational Mechanics and Analysis, **217**/2 (2015), 525-570.
- [5] D. Bartolucci, F. De Marchis, A. Malchiodi, *Supercritical conformal metrics on surfaces with conical singularities*, Int. Math. Res. Not. 2011, (2011)(**24**), 5625-5643.
- [6] D. Bartolucci, C. Gui, Y. Hu, A. Jevnikar, W. Yang, *Mean field equation on torus: existence and uniqueness of evenly symmetric blow-up solutions*, Preprint (2019); arXiv:1902.06934.
- [7] D. Bartolucci, C. Gui, A. Jevnikar, A. Moradifam, *A singular Sphere Covering Inequality: uniqueness and symmetry of solutions to singular Liouville-type equations*, Math. Ann. (2018); DOI:10.1007/s00208-018-1761-1.
- [8] D. Bartolucci, A. Jevnikar, Y. Lee, W. Yang, *Uniqueness of bubbling solutions of mean field equations*, J. Math. Pures Appl. **123** (2019), 78-126.
- [9] D. Bartolucci, A. Jevnikar, Y. Lee, W. Yang, *Non degeneracy, Mean Field Equations and the Onsager theory of 2D turbulence*, Arch. Rat. Mech. Anal. (ARMA) **230**(1) (2018), 397-426.
- [10] D. Bartolucci, A. Jevnikar, Y. Lee, W. Yang, *Local uniqueness of m-bubbling sequences for the Gel'fand equation*, Comm. P. D. E. (2019); DOI:10.1080/03605302.2019.1581801.
- [11] D. Bartolucci, A. Jevnikar, C.S. Lin, *Non-degeneracy and uniqueness of solutions to singular mean field equations on bounded domains*, J. Diff. Eq. **266**(1) (2019), 716-741.
- [12] D. Bartolucci, C.S. Lin, *Uniqueness Results for Mean Field Equations with Singular Data*, Comm. in P. D. E. **34**(7) (2009), 676-702.
- [13] D. Bartolucci, C.S. Lin, *Existence and uniqueness for Mean Field Equations on multiply connected domains at the critical parameter*, Math. Ann. **359** (2014), 1-44.
- [14] D. Bartolucci, C.S. Lin, G. Tarantello, *Uniqueness and symmetry results for solutions of a mean field equation on  $S^2$  via a new bubbling phenomenon*, Comm. Pure Appl. Math. **64**(12) (2011), 1677-1730.
- [15] D. Bartolucci, A. Malchiodi, *An improved geometric inequality via vanishing moments, with applications to singular Liouville equations*, Comm. Math. Phys. **322** (2013), 415-452.
- [16] D. Bartolucci, E. Montefusco, *Blow up analysis, existence and qualitative properties of solutions for the two dimensional Emden-Fowler equation with singular potential*, M<sup>2</sup>.A.S. **30**(18) (2007), 2309-2327.
- [17] D. Bartolucci, G. Tarantello, *Liouville type equations with singular data and their applications to periodic multivortices for the electroweak theory*, Comm. Math. Phys. **229** (2002), 3-47.
- [18] D. Bartolucci, G. Tarantello, *Asymptotic blow-up analysis for singular Liouville type equations with applications*, J. Diff. Eq. **262** (2017), 3887-3931.
- [19] H. Brezis, F. Merle, *Uniform estimates and blow-up behaviour for solutions of  $-\Delta u = V(x)e^u$  in two dimensions*, Comm. in P.D.E., **16**(8,9) (1991), 1223-1253.
- [20] E. Caglioti, P.L. Lions, C. Marchioro, M. Pulvirenti, *A special class of stationary flows for two dimensional Euler equations: a statistical mechanics description. II*, Comm. Math. Phys. **174** (1995), 229-260.
- [21] A. Carlotto, A. Malchiodi, *Weighted barycentric sets and singular Liouville equations on compact surfaces*, J. Funct. Anal. **262**(2) (2012), 409-450.
- [22] C.C. Chai, C.S. Lin, C.L. Wang, *Mean field equations, hyperelliptic curves, and modular forms: I*, Camb. J. Math. **3**(1-2) (2015), 127-274.
- [23] S.Y.A. Chang, C.C. Chen, C.S. Lin, *Extremal functions for a mean field equation in two dimension*, Lecture on Partial Differential Equations, New Stud. Adv. Math. **2** Int. Press, Somerville, MA, 2003, 61-93.
- [24] Z.J. Chen, T.J. Kuo, C.S. Lin, *Hamiltonian system for the elliptic form of Painlevé VI equation*, J. Math. Pure App. **106**(3) (2016), 546-581.
- [25] C. C. Chen, C.S. Lin, *Sharp estimates for solutions of multi-bubbles in compact Riemann surface*. Comm. Pure Appl. Math. **55** (2002), 728-771.

- [26] C.C. Chen, C.S. Lin, *Topological degree for a mean field equation on Riemann surfaces*, Comm. Pure Appl. Math. **56** (2003), 1667-1727.
- [27] C.C. Chen, C.S. Lin, *Mean field equations of Liouville type with singular data: sharper estimates*, Discr. Cont. Dyn. Syst. **28**(3) (2010), 1237-1272.
- [28] C.C. Chen, C.S. Lin, *Mean field equation of Liouville type with singular data: topological degree*, Comm. Pure Appl. Math. **68**(6) (2015), 887-947.
- [29] F. De Marchis, *Generic multiplicity for a scalar field equation on compact surfaces*, J. Funct. An. **259** (2010), 2165-2192.
- [30] W. Ding, J. Jost, J. Li, G. Wang, *Existence results for mean field equations*, Ann. Inst. H. Poincaré Anal. Non Linéaire **16** (1999), 653-666.
- [31] Z. Djadli, *Existence result for the mean field problem on Riemann surfaces of all genres*, Comm. Contemp. Math. **10**(2) (2008), 205-220.
- [32] P. Esposito, M. Grossi, A. Pistoia, *On the existence of blowing-up solutions for a mean field equation*, Ann. Inst. H. Poincaré Anal. Non Linéaire **22**(2) (2005), 227-257.
- [33] C. Gui, A. Moradifard, *The Sphere Covering Inequality and Its Applications*, Invent. Math. **214**(3) (2018), 1169-1204.
- [34] C. Gui, A. Moradifard, *Uniqueness of solutions of mean field equations in  $\mathbb{R}^2$* , Proc. Am. Math. Soc. **146** (2018), 1231-1242.
- [35] J.L. Kazdan, F.W. Warner, *Curvature functions for compact 2-manifolds*, Ann. Math. **99** (1974), 14-74.
- [36] M. Kowalczyk, M. Musso, M. del Pino, *Singular limits in Liouville-type equations*, Calc. Var. P.D.E. **24**(1) (2005), 47-81.
- [37] T.J. Kuo, C.S. Lin, *Estimates of the mean field equations with integer singular sources: non-simple blow up*, Jour. Diff. Geom. **103** (2016), 377-424.
- [38] Y.Y. Li, *Harnack type inequality: the method of moving planes*, Comm. Math. Phys., **200** (1999), 421-444.
- [39] Y.Y. Li, I. Shafrir, *Blow-up analysis for Solutions of  $-\Delta u = V(x)e^u$  in dimension two*, Ind. Univ. Math. J., **43**(4) (1994), 1255-1270.
- [40] C.S. Lin, *Uniqueness of solutions to the mean field equation for the spherical Onsager Vortex*, Arch. Rat. Mech. An. **153** (2000), 153-176.
- [41] C.S. Lin, M. Lucia, *Uniqueness of solutions for a mean field equation on torus*, J. Diff. Eq. **229**(1) (2006), 172-185.
- [42] C.S. Lin, C.L. Wang, *Elliptic functions, Green functions and the mean field equations on tori*, Ann. of Math. **172**(2) (2010), 911-954.
- [43] C.S. Lin, S. Yan, *On the mean field type bubbling solutions for Chern-Simons-Higgs equation*, Advances in Mathematics **338** (2018), 1141-1188.
- [44] L. Ma, J. Wei, *Convergence for a Liouville equation*, Comment. Math. Helv. **76** (2001), 506-514.
- [45] A. Poliakovsky, G. Tarantello, *On a planar Liouville-type problem in the study of selfgravitating strings*, J. Diff. Eq. **252** (2012), 3668-3693.
- [46] J. Prajapat, G. Tarantello, *On a class of elliptic problems in  $\mathbb{R}^2$ : Symmetry and uniqueness results*, Proc. Roy. Soc. Edinburgh Sect. A. **131** (2001), 967-985.
- [47] J. Spruck, Y. Yang, *On Multivortices in the Electroweak Theory I: Existence of Periodic Solutions*, Comm. Math. Phys. **144** (1992), 1-16.
- [48] T. Suzuki, *Global analysis for a two-dimensional elliptic eigenvalue problem with the exponential nonlinearity*, Ann. Inst. H. Poincaré Anal. Non Linéaire **9**(4) (1992), 367-398.
- [49] G. Tarantello, *Multiple condensate solutions for the Chern-Simons-Higgs theory*, J. Math. Phys. **37** (1996), 3769-3796.
- [50] M. Troyanov, *Prescribing curvature on compact surfaces with conical singularities*, Trans. Amer. Math. Soc. **324** (1991), 793-821.
- [51] J. Wei, L. Zhang, *Estimates for Liouville equation with quantized singularities*, Preprint (2019); arxiv: 1905.04123.
- [52] G. Wolansky, *On steady distributions of self-attracting clusters under friction and fluctuations*, Arch. Rational Mech. An. **119** (1992), 355-391.
- [53] Y. Yang, "Solitons in Field Theory and Nonlinear Analysis", Springer Monographs in Mathematics, Springer, New York, 2001.
- [54] L. Zhang, *Asymptotic behavior of blowup solutions for elliptic equations with exponential nonlinearity and singular data*, Commun. Contemp. Math. **11** (2009), 395-411.

DANIELE BARTOLUCCI, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ROME "*Tor Vergata*",  
VIA DELLA RICERCA SCIENTIFICA N.1, 00133 ROMA, ITALY.

*E-mail address:* bartoluc@mat.uniroma2.it

ALEKS JEVNIKAR, SCUOLA NORMALE SUPERIORE, PIAZZA DEI CAVALIERI 7, 56126, PISA, ITALY.

*E-mail address:* aleks.jevnikar@sns.it

YOUNGAE LEE, DEPARTMENT OF MATHEMATICS EDUCATION, TEACHERS COLLEGE, KYUNGPOOK  
NATIONAL UNIVERSITY, DAEGU, SOUTH KOREA

*E-mail address:* youngaelee@knu.ac.kr

WEN YANG, WUHAN INSTITUTE OF PHYSICS AND MATHEMATICS, CHINESE ACADEMY OF SCI-  
ENCES, P.O. BOX 71010, WUHAN 430071, P. R. CHINA

*E-mail address:* wyang@wipm.ac.cn