
ATTI ACCADEMIA NAZIONALE DEI LINCEI
CLASSE SCIENZE FISICHE MATEMATICHE NATURALI
RENDICONTI

PAOLO LIPPARINI

Some results about compact logics

*Atti della Accademia Nazionale dei Lincei. Classe di Scienze Fisiche,
Matematiche e Naturali. Rendiconti, Serie 8, Vol. 72 (1982), n.6, p. 308–311.*

Accademia Nazionale dei Lincei

<http://www.bdim.eu/item?id=RLINA_1982_8_72_6_308_0>

L'utilizzo e la stampa di questo documento digitale è consentito liberamente per motivi di ricerca e studio. Non è consentito l'utilizzo dello stesso per motivi commerciali. Tutte le copie di questo documento devono riportare questo avvertimento.

*Articolo digitalizzato nel quadro del programma
bdim (Biblioteca Digitale Italiana di Matematica)*

SIMAI & UMI

<http://www.bdim.eu/>

Logica matematica. — *Some results about compact logics.* Nota di PAOLO LIPPARINI, presentata (*) dal Socio G. ZAPPA.

RIASSUNTO. — Nell'ambito della Teoria dei Modelli Astratta è possibile dimostrare che una logica compatta L è univocamente determinata dalla sua relazione di L -elementare equivalenza (Teorema 1). Si enunciano poi alcuni risultati sulle logiche massime correlate a certe relazioni di equivalenza e sulle logiche compatte generate da qualche sistema di Fraïssé-Ehrenfeucht.

The aim of this note is to announce some results of the author in Soft Model Theory: their proofs will appear in [3].

Soft (or Abstract) Model Theory studies the extensions of first-order logic ($L_{\omega\omega}$): in particular two kinds of problems are taken into account: i) the research for extensions of first-order logic having some of its good model-theoretical properties (for example satisfying compactness, Craig interpolation theorem, Löwenheim-Skolem theorems and so on); and ii) the characterization of existing logics as maximal with respect to certain properties.

For example, the first non trivial theorem of Abstract Model Theory, proved by Lindström in 1966 [2], says that $L_{\omega\omega}$ is the only compact logic satisfying the downward Löwenheim-Skolem theorem, so that it is maximal among those having these properties. Naturally, this gives negative answers to problem i); nevertheless, by a result of Shelah [11], compact proper extensions of $L_{\omega\omega}$ do exist.

These are only examples of the nature of Soft Model Theory: many other interesting results and applications can be found in the literature (see [1]). In particular some logical concepts are given a purely algebraic characterization: in [6] and [4] it is proved that: i) compactness is equivalent to the joint embedding property, and also to the amalgamation property for L -elementary embeddings; ii) compactness+Craig interpolation is equivalent to the Robinson property. (In passing we remark that the Robinson property has many useful consequences, see [7], [8], [9]).

The strength of these characterizations can be seen from the following example: Lindström proved that $L_{\omega\omega}$ is the only compact logic L generating \equiv as L -elementary equivalence. But any logic generating \equiv must satisfy the joint embedding property in view of i) above. So, the quoted proposition can be improved to: $L_{\omega\omega}$ is the only logic generating \equiv .

(*) Nella seduta del 25 giugno 1982.

In [7] similar theorems results are obtained for compact logics satisfying Craig in place of $L_{\omega\omega}$; but perhaps these results can be applied only to $L_{\omega\omega}$, since, as far as we know, Friedman's fourth problem is still open (the research for compact proper extensions of $L_{\omega\omega}$ satisfying Craig). Nevertheless our Theorem 1 says that the Craig assumption is unnecessary, and this makes the result 'non-empty'. For example, a concrete consequence is that there is no logic $L \neq L_{\omega\omega} (Q^{ef\omega})$ such that $\equiv_L = \equiv_{L_{\omega\omega} (Q^{ef\omega})}$.

The fact that algebraic properties of equivalence relations can be useful for the study of logics has led to the following problem: characterize those equivalence relations which are generated by some logic (see [7], [10]). A rather complete answer for Robinson logics is given in [9]; in proposition 1 we extend some of the results of the latter paper to any countably generated logic.

Theorems 1 and 3 support the author's belief that, with few more technical efforts, many theorems about compact logics satisfying Craig can be proved assuming only compactness.

All notations and definitions are taken from [5], [9], [7]; we use the definition of logic of this last paper.

My thanks go to P. Mangani, my Research Director, and to D. Mundici, for his valuable help.

THEOREM 1. *If L, M are logics such that $\text{Stn}_L(\tau)$ is a set for any τ , \equiv_M is coarser than \equiv_L , and L is compact, then $M \leq L$. Moreover, if $\equiv_M = \equiv_L$, then $M = L$.*

DEFINITION 1. Let \sim be any equivalence relation on the class of all structures. We say that a logic L respects \sim iff whenever $\mathfrak{A} \sim \mathfrak{B}$ are two models of type τ , and $\varphi \in \text{Stn}_L$ has type $\tau \cup \{c_1, \dots, c_n\}$, if we add to \mathfrak{A} and \mathfrak{B} a new relation R defined by:

$$R(c_1, \dots, c_n) \iff \varphi,$$

then the two expanded models are also \sim -equivalent.

THEOREM 2. *Let \sim be an equivalence relation finer than \equiv and closed under renaming and reduct. If L and M respect \sim , then $L \cup M$ respects \sim . Thus, there exists the largest logic respecting \sim . Moreover, if $L = L_{\omega\omega} (Q_i)_{i \in I}$, then there exists the largest logic M of the form $M = L_{\omega\omega} (Q_j)_{j \in J}$ such that $\equiv_L = \equiv_M$.*

It can be proved that if \sim has Robinson, then L respects \sim if \equiv_L is coarser than \sim , so Theorem 2 generalizes parts of Theorem 5.5 of [7] (see also Remarks 6.10).

DEFINITION 2. A FE-system satisfies the *expansion property* if, whenever $\mathfrak{A} \simeq_{\tau}^{m+n} \mathfrak{B}$, $\tau' = \tau \cup \{\bar{c}\}$, U is a union of classes of $\simeq_{\tau'}^n$, then $(\mathfrak{A}, R) \simeq_{\tau \cup \{R\}} (\mathfrak{B}, R)$, where R is defined in \mathfrak{A} by

$$R\bar{a} \iff \langle \mathcal{A}, \bar{a} \rangle \in U,$$

and similarly for \mathfrak{B} .

PROPOSITION 1. For any \sim , there is a FE-system satisfying expansion and generating \sim if there is a countably generated logic L with $\equiv_L = \sim$.

THEOREM 3. For any \sim , the following are equivalent:

- i) $\sim = \equiv_L$ for some countably generated compact logic;
- ii) $\sim = \equiv_L$ for exactly one logic L , which is compact and countably generated;
- iii) \sim has the joint embedding property and is generated by some FE-system satisfying expansion;
- iv) \sim has the joint embedding property and is generated by exactly one FE-system, which satisfies expansion.

ADDED IN PROOF (DECEMBER 1982). Other results are:

i) $L \cup M$ is $(\lambda, \mu)_\nu$ -compact iff $S_\mu(\lambda)$ is not ν -atomically characterizable by $\equiv_L \cap \equiv_M$ (cf. [3]); $(\lambda, \mu)_\nu$ -compactness is defined as $(\lambda, \mu)^*$ -compactness in [4], but requiring $|\tau(\Sigma \cup \Sigma_1)| < \nu$.

ii) If $\equiv_L \subseteq \equiv_M$, $OC(M) \leq \nu$, L is (λ, ω) -compact and $|\text{Stn}_L(\tau)| \leq \lambda$ for $|\tau| < \nu$, then $M \leq L$.

iii) If $OC(L) \leq \lambda$, and \equiv_L satisfies the Robinson property ([8]) with the restriction that $|\tau| \leq \lambda$ and $|\tau' \setminus \tau|, |\tau'' \setminus \tau| < \nu$, then L is $(\lambda, \omega)_\nu$ -compact.

iv) If L is compact and generated by a set of quantifiers, then L satisfies Beth's Definability Theorem if \equiv_L has the definability property ([3]).

v) If D is an ultrafilter, α an ordinal, and $\mathfrak{A} \sim \mathfrak{B}$ iff $\prod_D^\beta \mathfrak{A} \cong \prod_D^\gamma \mathfrak{B}$ for some $\beta, \gamma < \omega^\alpha$ (where $\prod_D^0 \mathfrak{A} = \mathfrak{A}$, and $\prod_D^\beta \mathfrak{A} = \bigcup_{\gamma \in \beta} \prod_D (\prod_D^\gamma \mathfrak{A})$), then \sim is a Robinson equivalence relation. If ω is the only measurable cardinal and $\sim = \equiv_L$, then $\sim = \cong$.

BIBLIOGRAPHY

- [1] K. J. BARWISE (1974) - *Axioms for abstract model theory*, «Ann. Math. Logic», 7, 221-265.
- [2] P. LINDSTROM (1966) - *First order predicate logic with generalized quantifiers*, «Theoria», 32, 186-195.
- [3] P. LIPPARINI - *Duality for compact logics and substitution in abstract model theory*, preprint.
- [4] J. A. MAKOWSKY and S. SHELAH, *Positive results in abstract model theory*, preprint.
- [5] J. A. MAKOWSKY, S. SHELAH and S. STAVI (1976) - Δ -logics and generalized quantifiers, «Ann. Math. Logic.», 10, 155-192.
- [6] D. MUNDICI (1982) - *Interpolation, compactness and JEP in soft model theory*, «Arch. Math. Logik», 22, 61-67.

-
- [7] D. MUNDICI (1982) – *Duality between logics and equivalence relations*, «Trans. Amer. Math. Soc.», 270, 111–129.
 - [8] D. MUNDICI (1981) – *An algebraic result about soft model theoretical equivalence relations with an application to H. Friedman's fourth problem*, «J. Symbolic Logic», 46, 523–530.
 - [9] D. MUNDICI (1981) – *Variations on Friedman's third and fourth problem*, in Proceedings of the International Conference: «Open Days for Model Theory and Set Theory» (H. RASIOWA and W. MAREK Chair.), Warsaw Sept.
 - [10] M. E. NADEL (1980) – *An arbitrary equivalence relation as elementary equivalence in abstract logic*, «Z. Math. Logik Grundlagen Math.», 26, 103–109.
 - [11] S. SHELAH (1975) – *Generalized quantifiers and compact logic*, «Trans. Amer. Math. Soc.», 204, 342–364.