



**ISTITUTO NAZIONALE DI FISICA NUCLEARE**  
Laboratori Nazionali di Frascati

**FRASCATI PHYSICS SERIES**



**Vulcano 2016**

**Frontier Objects  
in Astrophysics and Particle Physics**

**Editors:**

R. Fusco Femiano, G. Mannocchi, A. Morselli, G. Trincherò

## FRASCATI PHYSICS SERIES

Series Editor

*Rossana Centioni*

Technical Editor

*Debora Bifaretti*

*Cover by Claudio Federici*

---

Volume LXIV

---

*Istituto Nazionale di Fisica Nucleare – Laboratori Nazionali di Frascati  
Divisione Ricerca  
SIDS – Servizio Informazione e Documentazione Scientifica  
Ufficio Biblioteca e Pubblicazioni  
P.O. Box 13, I-00044 Frascati Roma Italy  
email: [sis.publications@lnf.infn.it](mailto:sis.publications@lnf.infn.it)*

Copyright © 2017 by INFN

*All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior permission of the copyright owner.*

ISBN 978-88-86409-65-0

Frascati Physics Series Vol. 64 (2016)  
FRONTIER OBJECTS IN ASTROPHYSICS AND PARTICLE PHYSICS  
May 22-28, 2016

**LISA PATHFINDER  
FIRST STEP TOWARD A GRAVITATIONAL WAVE SPACE  
OBSERVATORY**

Massimo Bassan  
*Dipartimento di Fisica Università di Roma Tor Vergata  
& INFN Roma Tor Vergata*

On behalf of the LISA Pathfinder collaboration (\*)

**Abstract**

We briefly review the concept of a space-based gravitational wave interferometer, and the science it can explore in the milliHertz frequency region. Then we discuss the LISA Pathfinder technology demonstrator mission that is currently flying and will soon deliver the first results.

**1 Gravitational astronomy from space**

Gravitational waves have finally been detected <sup>1)</sup>, one century after being predicted by A. Einstein <sup>2)</sup>. The very first signal has already shown the amazing power of gravitational astronomy, offering detailed information about the merging of two black holes of unexpected mass. In the years to come, Earth-based Advanced Interferometers will provide a great wealth of information about compact objects like neutron stars and black holes in their final stages of life

before merging.

What is the role then of a space observatory like LISA, expected to operate no sooner than fifteen years from now ?

Theoretical predictions suggest that a wide variety of astrophysical sources emit gravitational waves (g.w.) in that part of the frequency spectrum, from 0.1 mHz to about 100 mHz, that is only accessible by a space interferometer. Signals gathered in this band can provide crucial insights about binary stars formation, Extreme Mass Ratio Inspirals, SuperMassive Black Holes and galaxy mergers. An observatory in that frequency range has moreover guaranteed signals for calibration, the so called Verification Binaries, of which all parameters are known by e.m. observations and whose g.w. flux is precisely predicted.

## 2 The LISA mission concept

The frequency range below 1 Hz, where all these sources are to be observed, is only accessible by a space-borne detector, that is immune from Earth-generated, low frequency disturbances, like seismic and Newtonian noise. LISA is a space mission concept, developed over two decades through several different configurations, that addresses this request. LISA's basic scientific and technological fundamentals are well established since many years<sup>3)</sup>. Its actual design has undergone several revisions, but has reached a stable configuration since 1998<sup>4)</sup>; it relies on a few pillars:

- *The orbital configuration:* a redundant set of three interferometers with arm length of 1-5 Gm (million kilometers), hosted by a constellation of three satellites arranged at the vertices of a gigantic triangle. Each spacecraft (S/C) passively move on a different "smart" heliocentric orbit<sup>5)</sup>: while keeping the relative distance roughly constant, the triangular pattern rotates around its center, that moves on the ecliptic, trailing the Earth by some  $20^\circ$ . This motion modulates the g.w. signals, giving them a unique signature.
- *The optical transponders:* the divergence of a laser beam is such that, even using a large telescope, only a few parts in  $10^9$  of the incoming beam can be received at the far end of such a long interferometric arm, thus making it unfeasible to reflect the light back with a mirror. A transponder<sup>6)</sup> is a

phase-preserving amplifier that will regenerate the beam to its full power (1-5 W) and send it back to the other S/C for phase comparison.

- *Time delay interferometry:* The distances between S/C's vary in time, due to Keplerian dynamics, of 1-2 %, i.e. up to  $10^5$  km: the unequal arm length gives rise to frequency noise in the interferometer. This noise can be depressed, by several order of magnitude, by recording the phase signal of each arm and then, off-line, synthesizing linear combinations of the signals emitted at different times<sup>7)</sup>, before comparing them, thus canceling most of the the frequency noise. So, the usual optical path difference with equal arms  $X(t) = y_1(t) - y_2(t)$  is substituted by:

$$X(t) = y_1(t) - y_2(t) - y_1(t - 2L_2/c) + y_2(t - 2L_1/c) \quad (1)$$

More complex relations (second order TDIs) would apply if one also wants to cancel the Doppler effect of different S/C velocities .

- *Test masses in free fall:* aboard each S/C, two test masses (TMs), one for each interferometer arm ending at that node, are in geodesic motion, shielded by the spacecraft from any external disturbance, and responding only to space-time perturbation. A feedback loop, acting on the S/C microthrusters, recenters the S/C on the TM along the sensitive axis. All other Degrees of Freedom (DoFs) are stabilized by electrostatic forces. Each interferometric arm acts then as a huge differential accelerometer.

In 2011, financial constraints forced NASA to withdraw its support for the LISA mission. European scientists undertook an intense effort to "rescope" the mission so that it could fit in the "ESA only" budget: the ambitious goal was to achieve the same physics with half the budget. This study produced a new configuration, called "eLISA": still with three spacecrafts, but with only one interferometer (two arms, therefore giving up redundancy and polarization detection capabilities), a reduced (1 or 2 Gm) arm length (thus making do with less laser power and without a pointing mechanism for the telescopes) and a shorter mission duration. This project, presented to ESA in 2013<sup>8)</sup>, gained for the theme *The Gravitational Universe* the selection to third large ESA mission of the Cosmic Vision Plan (L3), for launch in 2034. Nevertheless, recent re-analyses of the science, technology and finances for the mission make

us hopeful that the full, three arm configuration can be regained before the detailed mission formulation.




Figure 1: The *smart orbits* of the space g.w. observatory: each spacecraft moves on an independent, passive, Keplerian orbit, and the constellation maintains a triangular formation, while rotating around its center. Picture from ref. 8)

### 3 LISA Pathfinder, the technology demonstrator

The technical difficulty of the LISA project made it sensible to test as much as possible of the enabling technology in a dedicated mission. Particularly challenging is the assessment of the quality of free fall that can be achieved in space, despite extensive laboratory investigations, carried out with torsion pendulums<sup>9)</sup>, of the residual forces that can locally disturb the inertial motion of the TM. Other items that need to be space-tested are the interferometric read-out of the TM positions, the Gravitational Reference Sensor (GRS) that senses motion in the other degrees of freedom and applies electrostatic feedback, its front-end electronics (FEE) the microthrusters, TM decaging and release (with minimum velocity) and more.

Therefore ESA has approved and realized a dedicated space mission to verify all the flight hardware that can be tested in a single-spacecraft mission, and to evaluate the level of geodesic motion that current technology can achieve

and measure. Such mission, called LISA Pathfinder <sup>10</sup>), is active and operational now.

In LISA Pathfinder, a single spacecraft holds two independent TMs: once spaceborne, each TM is released within its GRS and their relative distance is measured by a differential interferometer: in many respects, this is a LISA arm shrunk from few Gm down to 38 cm. Feedback loops act on the spacecraft and on the second TM to make sure that they both follow the first TM in its free fall. Most of the hardware (with few exceptions like the telescope, the pointing mechanism and the TDI) are therefore tested in an environment that is representative of the final LISA mission. Moreover, a large number of noise sources, i.e. all *local* sources, like thermal, magnetic, laser shot and radiation pressure, long term changes in local gravity etc. can be accurately assessed and measured, so that a reliable noise budget can be formulated for the LISA mission.

LPF was launched with the sixth VEGA launcher on Dec 3rd 2015, one




Figure 2: Schematics of the LISA Pathfinder payload: two Test Masses, each surrounded by its GRS (electrostatic sensors and actuators) and, in between them, the monolithic Optical Bench, with all the optics needed to operate 4 different interferometers

century after publication of Einstein's first paper on the *General Theory of Relativity* and few weeks after the detection of the first g.w. signal by LIGO, thus marking an unforgettable year for Gravitational research. After a few

orbits around the Earth, raising each time its apogee, it has started its journey toward the first Lagrangian point (L1) of the Sun-Earth system, where the gravitational gradient is a minimum, about 1.5 Gm from the Earth. It has now reached its target orbit, a large, slow orbit around L1 where it has undergone testing and commissioning. At the time of this talk, operations are under way and the first results are extremely promising. However, no data has yet been published, and expectations are high for the first release due in June.

#### 4 Post-conference update: first results from LISA Pathfinder




Figure 3: Spectrum of the differential acceleration noise of the two Test Masses of LISA Pathfinder. We also show the requirements, well exceeded the present mission and closely approached for the future LISA observatory. Some known sources of noise, measured and subtracted, are also indicated. The rise on the right is readout noise (white in displacement), that grows as  $\omega^2$  when converted to acceleration. Figure from ref. <sup>11</sup>).

On June 7th 2016, about two weeks after the workshop, the first LPF data were released <sup>11</sup>). The results are excellent, beyond the most optimistic expectations: the interferometer readout noise is  $35 fm/\sqrt{Hz}$ , that is 100 times better than the requirements and than noise measured on ground, where mirrors could be carefully hand-aligned. The purity of free-fall is gauged by the differ-

ential acceleration of the two test masses, and is measured to be  $5.2 \cdot 10^{-15} m s^{-2}$  i.e. below the femto-g range. This is also lower, by a factor 5, than the mission requirements and closely approaches the tougher spec required for LISA. Even more striking, the specs are exceeded not only on the required frequency region, i.e. down to 1 mHz, but on a band extending almost down to 0.1 mHz, i.e. on the full LISA sensitivity band. Figure 3 shows the residual acceleration noise, together with the LISA Pathfinder requirements and the LISA requirements, that are approached to within a factor 1.5. Moreover, the noise in the "flat" region 2-8 mHz, that is mainly of Brownian origin, appears to be decreasing with time: a probable reason for this is the continuing decrease in pressure due to venting to open space.

The tests and investigations on LISA Pathfinder will continue for the next several months, with the aim of characterizing all components of the residual noise, but we can already confidently state that LISA Pathfinder is doing its job and the road to LISA looks brighter than ever.

## References

1. B.P.Abbott et al. Phys Rev Lett **116**, 061102 (2016)
2. A. Einstein, Sitzungsber. K. Preuss. Akad. Wiss. **1**, 688 (1916).
3. See e.g. K.Danzmann, A.Rüdiger, Class. Quantum Grav. **20** (2003) S1S9
4. LISA Pre-Phase A Report, MPQ 233, July 1998
5. F.Hechler and W.M.Folkner Adv. Space Res. **32**, No. 7, 1277,(2003)
6. Yinan Yu, Shawn Mitryk, and Guido Mueller Phys. Rev. D **90**, 062005
7. M.Tinto, S. V. Dhurandar: Living Rev. Relativity, **17**, 6, (2014)
8. The eLISA Consortium, arXiv:1305.5720.
9. L Carbone et al. Phys Rev Lett **91**, 151101, (2003)  
M. Bassan et al. Phys Rev Lett **116**, 051104 (2016) and references therein
10. P. McNamara, S. Vitale, and K. Danzmann, Classical Quantum Gravity **25**, 114034 (2008).
11. M. Armano et al. Phys. Rev. Lett **116**, 231101 (2016).

**(\*) The LISA Pathfinder collaboration:**

M Armano<sup>a</sup>, H Audley<sup>b</sup>, G Auger<sup>c</sup>, J T Baird<sup>d</sup>, M Bassan<sup>e</sup>, P Binetruy<sup>c</sup>, M Born<sup>b</sup>, D Bortoluzzi<sup>f</sup>,  
N Brandt<sup>g</sup>, M Caleno<sup>h</sup>, L Carbone<sup>i</sup>, A Cavalleri<sup>bb</sup>, A Cesarini<sup>i</sup>, G Ciani<sup>i,dd</sup>, G Congedo<sup>i,cc</sup>,  
A M Cruise<sup>j</sup>, K Danzmann<sup>b</sup>, M de Deus Silva<sup>a</sup>, R De Rosa<sup>y</sup>, L Di Fiore<sup>y</sup>, M Diaz-Aguilón<sup>n</sup>,  
I Diepholz<sup>b</sup>, G Dixon<sup>j</sup>, R Dolesi<sup>i</sup>, N Dunbar<sup>k</sup>, L Ferraioli<sup>l</sup>, V Ferroni<sup>i</sup>, E D Fitzsimons<sup>m</sup>,  
R Flatscher<sup>g</sup>, M Freschi<sup>a</sup>, A Garcia Marín<sup>b,ff</sup>, C García Marrero<sup>h</sup>, R Gerndt<sup>g</sup>, L Gesa<sup>n</sup>,  
F Gibert<sup>i</sup>, D Giardini<sup>l</sup>, R Giusteri<sup>i</sup>, A Grado<sup>x</sup>, C Grimani<sup>o</sup>, A Grynagier<sup>gg</sup>, J Grzymisch<sup>h</sup>,  
F Guzman<sup>b</sup>, I Harrison<sup>p</sup>, G Heinzl<sup>b</sup>, M Hewitson<sup>b</sup>, D Hollington<sup>d</sup>, D Hoyland<sup>j</sup>, M Hueller<sup>i</sup>,  
H Inchauspé<sup>c</sup>, O Jennrich<sup>h</sup>, P Jetzer<sup>q</sup>, B Johlander<sup>h</sup>, N Karnesis<sup>b</sup>, B Kaune<sup>b</sup>, N Korsakova<sup>b</sup>,  
C J Killow<sup>r</sup>, J A Lobo<sup>n,ii</sup>, I Lloro<sup>n</sup>, L Liu<sup>i</sup>, J P López-Zaragoza<sup>n</sup>, R Maarschalkerweerd<sup>p</sup>,  
D Mance<sup>l</sup>, V Martín<sup>n</sup>, L Martin-Polo<sup>a</sup>, J Martino<sup>c</sup>, F Martin-Porqueras<sup>a</sup>, S Madden<sup>h</sup>,  
I Mateos<sup>n</sup>, P W McNamara<sup>h</sup>, J Mendes<sup>p</sup>, L Mendes<sup>a</sup>, A Monsky<sup>b</sup>, D Nicolodi<sup>i,ee</sup>, M Nofrarias<sup>n</sup>,  
S Paczkowski<sup>b</sup>, M Perreur-Lloyd<sup>r</sup>, A Petiteau<sup>c</sup>, P Pivato<sup>i</sup>, E Plagnol<sup>c</sup>, P Prat<sup>c</sup>, U Ragnit<sup>h</sup>,  
B Raïs<sup>c</sup>, J Ramos-Castro<sup>s</sup>, J Reiche<sup>b</sup>, D I Robertson<sup>r</sup>, H Rozemeijer<sup>h</sup>, F Rivas<sup>n</sup>, G Russano<sup>i</sup>,  
J Sanjuan<sup>n,hh</sup>, P Sarra<sup>aa</sup>, A Schleicher<sup>g</sup>, D Shaul<sup>d</sup>, J Slutsky<sup>u</sup>, C F Sopena<sup>n</sup>, R Stanga<sup>z</sup>,  
F Steier<sup>b,ff</sup>, T Sumner<sup>d</sup>, D Texier<sup>a</sup>, J I Thorpe<sup>u</sup>, C Trenkel<sup>k</sup>, M Tröbs<sup>b</sup>, H B Tu<sup>i</sup>, D Vetrugno<sup>i</sup>,  
S Vitale<sup>i</sup>, V Wand<sup>b,jj</sup>, G Wanner<sup>b</sup>, H Ward<sup>r</sup>, P J Wass<sup>d</sup>, D Wealthy<sup>k</sup>, W J Weber<sup>i</sup>, L Wissel<sup>b</sup>,  
A Wittchen<sup>b</sup>, A Zambotti<sup>f</sup>, C Zanon<sup>f</sup>, T Ziegler<sup>g</sup>, P Zweifel<sup>l</sup>

<sup>a</sup>European Space Astronomy Centre, European Space Agency, Villanueva de la Cañada,  
28692 Madrid, Spain

<sup>b</sup>Albert-Einstein-Institut, Max-Planck-Institut für Gravitationsphysik und Universität Han-  
nover, Callinstrasse 38, 30167 Hannover, Germany

<sup>c</sup>APC, Univ Paris Diderot, CNRS/IN2P3, CEA/lrfu, Obs de Paris, Sorbonne Paris Cité,  
France

<sup>d</sup>High Energy Physics Group, Physics Department, Imperial College London, Blackett Lab-  
oratory, Prince Consort Road, London, SW7 2BW, UK

<sup>e</sup>Dipartimento di Fisica, Università di Roma “Tor Vergata”, and INFN, sezione Roma Tor  
Vergata, I-00133 Roma, Italy

<sup>f</sup>Department of Industrial Engineering, University of Trento, via Sommarive 9, 38123 Trento,  
and Trento Institute for Fundamental Physics and Application / INFN

<sup>g</sup>Airbus Defence and Space, Claude-Dornier-Strasse, 88090 Immenstaad, Germany

<sup>h</sup>European Space Technology Centre, European Space Agency, Keplerlaan 1, 2200 AG No-  
ordwijk, The Netherlands

<sup>i</sup>Dipartimento di Fisica, Università di Trento and Trento Institute for Fundamental Physics  
and Application / INFN, 38123 Povo, Trento, Italy

<sup>j</sup>The School of Physics and Astronomy, University of Birmingham, Birmingham, UK  
<sup>k</sup>Airbus Defence and Space, Gunness Wood Road, Stevenage, Hertfordshire, SG1 2AS, UK  
<sup>l</sup>Institut für Geophysik, ETH Zürich, Sonneggstrasse 5, CH-8092, Zürich, Switzerland  
<sup>m</sup>The UK Astronomy Technology Centre, Royal Observatory, Edinburgh, Blackford Hill, Edinburgh, EH9 3HJ, UK  
<sup>n</sup>Institut de Ciències de l'Espai (CSIC-IEEC), Campus UAB, Carrer de Can Magrans s/n, 08193 Cerdanyola del Vallès, Spain  
<sup>o</sup>DISPEA, Università di Urbino "Carlo Bo" & INFN, Via S. Chiara, 27 61029 Urbino, Italy  
<sup>p</sup>European Space Operations Centre, European Space Agency, 64293 Darmstadt, Germany  
<sup>q</sup>Physik Institut, Universität Zürich, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland  
<sup>r</sup>SUPA, Institute for Gravitational Research, School of Physics and Astronomy, University of Glasgow, Glasgow, G12 8QQ, UK  
<sup>s</sup>Depart. d'Enginyeria Electrònica, Universitat Politècnica de Catalunya, 08034 Barcelona, Spain  
<sup>t</sup>Inst. d'Estudis Espacials de Catalunya (IEEC), C/ Gran Capità 2-4, 08034 Barcelona, Spain  
<sup>u</sup>Gravitational Astrophysics Lab, NASA Goddard Space Flight Center, 8800 Greenbelt Road, Greenbelt, MD 20771 USA  
<sup>x</sup>INAF Osservatorio Astronomico di Capodimonte, I-80131 Napoli, Italy and INFN Sezione di Napoli, I-80126 Napoli, Italy  
<sup>y</sup>Dipartimento di Fisica, Università di Napoli "Federico II", I-80126, Napoli, Italy and INFN - Sezione di Napoli, I-80126, Napoli, Italy  
<sup>z</sup>Dipartimento di Fisica ed Astronomia, Università degli Studi di Firenze and INFN - Sezione di Firenze, I-50019 Firenze, Italy  
<sup>u</sup>Center for Space Science & Technology, University of Maryland Baltimore County, 1000 Hilltop Circle, Baltimore, MD 21250, USA  
<sup>aa</sup>CGS S.p.A, Compagnia Generale per lo Spazio, Via Gallarate, 150 - 20151 Milano, Italy  
<sup>bb</sup>Istituto di Fotonica e Nanotecnologie, CNR-Fondazione Bruno Kessler, I-38123 Povo, Italy  
<sup>cc</sup>University of Florida, Gainesville, FL 32611, USA  
<sup>dd</sup>Department of Physics, University of Oxford, Keble Road, Oxford, OX1 3RH  
<sup>dd</sup>Institute of Geodesy and Geophysics, CAS, No. 340 Xudong Street, Wuhan 430077, China  
<sup>ee</sup>National Institute for Standards and Technology, 325 Broadway, Boulder, CO 80305, USA  
<sup>ff</sup>OHB System AG, Universitätsallee 27-29, D-28359 Bremen, Germany  
<sup>gg</sup>Thales Alenia Space, 5 Allée des Gabians, BP 99 - 06156 Cannes, France  
<sup>hh</sup>Deutsches Zentrum für Luft-Raumfahrt, Robert-Hooke-Str.7, 28359 Bremen, Germany  
<sup>ii</sup> Deceased 30 September 2012  
<sup>jj</sup>Jena-Optronik GmbH, Otto-Eppenstein-Strasse 3, 07745 Jena