

The role of HECT-type E3 ubiquitin ligases in DNA damage response and repair

Received: 11 September 2025

Revised: 31 October 2025

Accepted: 26 November 2025

Cite this article as: **Giovannini, S., Fiorilli, C., Moriconi, V. *et al.*** The role of HECT-type E3 ubiquitin ligases in DNA damage response and repair. *Cell Death Discov.* (2025). <https://doi.org/10.1038/s41420-025-02911-0>

Sara Giovannini, Claudia Fiorilli, Valeria Moriconi, Yufang Shi, Eleonora Candi, Gerry Melino & Francesca Bernassola

We are providing an unedited version of this manuscript to give early access to its findings. Before final publication, the manuscript will undergo further editing. Please note there may be errors present which affect the content, and all legal disclaimers apply.

If this paper is publishing under a Transparent Peer Review model then Peer Review reports will publish with the final article.

The role of HECT-type E3 ubiquitin ligases in DNA damage response and repair

Sara Giovannini¹®, Claudia Fiorilli¹, Valeria Moriconi¹, Yufang Shi², Eleonora Candi^{1,3}, Gerry Melino¹, Francesca Bernassola¹®.

¹Department of Experimental Medicine, TOR, University of Rome Tor Vergata, 00133 Rome, Italy. ²The Third Affiliated Hospital of Soochow University, Institutes for Translational Medicine, Soochow University, Suzhou 215000, China. ³Biochemistry Laboratory, Istituto Dermopatico Immacolata (IDI-IRCCS), 00100 Rome, Italy.

@Correspondence: SG (sara.giovannini@uniroma2.it), FB (bernasso@uniroma2.it)

ORCID & Email contacts

Bernassola F	0000-0002-8883-8654	bernasso@uniroma2.it
Candi E	0000-0001-8332-4825	candi@uniroma2.it
Fiorilli C	0009-0005-6442-7083	claudiafiorilli98@gmail.com
Giovannini S	0000-0002-8916-476X	sara.giovannini@uniroma2.it
Melino G	0000-0001-9428-5972	melino@uniroma2.it
Moriconi V	0009-0002-0307-0680	moriconivaleria1@gmail.com
Shi Y	0000-0001-8964-319X	yufangshi@sibs.ac.cn

Abstract

The post-translational modification ubiquitination consists in a three-step reaction triggered by E1 ubiquitin activating enzymes, E2 ubiquitin conjugating enzymes, and E3 ubiquitin ligases. The latter enzymes, providing substrate specificity, play an important role in determining the fate of the substrate proteins, by regulating their level and function.

Efficient DNA damage response (DDR) is necessary to detect and signal DNA damage, thus favoring DNA damage repair to prevent genomic instability and tumorigenesis.

Differently from RING (really interesting new gene)-type E3s, the ones belonging to the Homologous to E6AP C-terminus (HECT) family have an intrinsic catalytic activity, which enables them to directly transfer ubiquitin molecules to their substrates. They participate in the regulation of numerous processes, from cell proliferation to apoptosis. Nevertheless, their role in DDR and repair is less known.

Recent evidence reports of the HECT E3s involvement in the regulation of DNA damage signaling, chromatin remodeling, repair pathway choice and DNA damage resolution. Further elucidating their functions in DDR and repair may provide new insights into the processes aimed at the preservation of genome integrity, putatively uncovering HECT E3s as therapeutic targets in tumors and defective DNA repair pathologies.

Key words

Ubiquitination, HECT E3s, DNA repair, Genomic instability, DDR.

Facts

- Counteracting DNA damage, DNA damage response and repair mechanisms play a crucial role in preserving genome integrity.
- Ubiquitination regulates the protein level and function of substrates, involved in multiple biological processes, thereby significantly affecting DNA damage repair.

- While the contribution of RING-type E3s to DNA damage repair signaling pathways is extensively reported, the role of HECT-type E3s in the regulation of DNA damage cell response remains less studied.
- Recent evidence unveils a role for HECT E3s in protecting from genome instability and suggests them as putative targets for new cancer therapies.

Open questions

- Which substrates are modified by HECT E3s in the regulation of DNA damage response and repair?
- How do HECT E3s cooperate with each other, or with members of other E3 families to maintain genome stability?
- Will future screenings and structural studies help in designing more selective inhibitors, able to overcome chemotherapy resistance and reduce off-target effects?

Background

Ubiquitination regulates cellular processes like cell cycle progression, cell death and DDR (1,2). It occurs through a cascade of enzymatic reactions: ubiquitin-activation by E1 enzymes, ubiquitin-conjugation by E2 enzymes and ubiquitin ligation by E3 ligases (3) (**BOX 1**). E3s confer substrate specificity and facilitate regulation and the attachment of ubiquitin molecules to target proteins (4). The HECT family E3 members are characterized by a unique catalytic mechanism. Unlike other E3s acting as scaffolds, HECT E3s form a thioester intermediate with ubiquitin before its direct transfer to the substrate protein. HECT E3s are involved in many cellular functions, including regulation of cell cycle, stress response, and several modalities of cell death, including apoptosis (5–7), autophagy (8–10), and ferroptosis (11–13), thus influencing tumorigenesis (8,14–17).

Evidence suggests an important contribution of HECT-type E3s in DDR and repair. Upon DNA damage, cells activate DDR pathways to detect lesions, signal their presence, and coordinate repair mechanisms (18). Importantly, ubiquitination influences the protein stability and function of various proteins involved in DDR. By controlling their ubiquitination status, HECT E3s can then influence the choice of the repair pathway, the efficiency of repair, as well as the overall response to DNA damage, to finally prevent mutagenesis. Consequently, dysregulation of HECT E3s could lead to defective DNA repair, accumulation of mutations, cancer and even increased resistance to therapies (15,16).

Compared to other E3 families, like RING E3s, the role of HECT E3s in DDR and repair is less characterized. A better understanding of the mechanisms ensuring genome integrity, by further investigations, may uncover new therapeutic approaches to possibly enhance DNA repair or to sensitize resistant tumors to treatments.

An overview of DDR and repair mechanisms

In response to the presence of DNA damage, signaling pathways, collectively known as DDR, are activated to maintain genome integrity (19,20). These signaling cascades trigger cell cycle checkpoints, induce cell cycle arrest and initiate transcriptional events that promote the expression or regulation of DNA repair factors (18,21). Defined as the guardian of the genome (22), the transcription factor p53 is the main player of DDR (23–25), as well as other related mechanisms (26–29). It is involved in cell cycle arrest and activation of transcriptional, repair and apoptotic events, therefore finely controlled in protein level, via an autoregulatory negative feedback loop. Its degradation depends on ubiquitination catalyzed by the RING E3 MDM2

(Mouse double minute 2), one of its principal transcription targets (30). Depending on the type of DNA damage, DDR signaling induces specific repair mechanisms, such as the double-strand breaks (DSBs) repair pathways, homologous recombination (HR) and non-homologous end joining (NHEJ), as well as the single-stranded DNA repair pathways, nucleotide excision repair (NER), base excision repair (BER), and mismatch repair (MMR) (17,31–34) (**BOX 2**). Multiple DNA repair proteins, belonging to different pathways, cooperate to remove specific types of DNA damage. As discussed in BOX 2, interstrand crosslinks (ICLs) activate Fanconi anemia (FA), HR and NER factors, while oxidative damage induces both BER and NER (35). Moreover, different repair systems can reciprocally influence each other, as for instance, defects in FA pathway could impair Alt (alternative)-NHEJ by favoring C (canonical)-NHEJ (36). Importantly, a different type of DNA damage may arise consequently to the intervention of a repair system, as revealed by single-strand breaks (SSBs) induced by BER glycosylases, or by unrepaired SSBs colliding with the replication fork, and finally leading to DSBs (37).

The interplay between different repair pathways is further supported by the synthetic lethal interaction observed between BER and HR defects, suggesting new therapeutic strategies, as the use of Poly(ADP-ribose)polymerase inhibitors (PARPis) to target HR-deficient tumors. In the presence of DNA breaks, PARP1 catalyzes its own ADP-ribosylation and the modification of other proteins, thus modifying chromatin structure at the sites of damage. This modification induces the recruitment of additional repair factors, including the BER scaffold protein XRCC1 (X-ray repair cross-complementing protein 1) (38). The aberrant activity of PARP1 can impair transcriptional recovery during repair by excessively recruiting the ubiquitin protease USP3, which reduces ubiquitinated histone levels (39). Moreover, PARP1 activity is regulated by ubiquitination-mediated degradation (40).

XRCC1 itself is subject to various post-translational modifications, such as poly-ADP-ribosylation, phosphorylation, and ubiquitination as well (41,42). Sensitivity to PARPis has also been observed in non-canonical HR-defective cells, which often show impaired responses to diverse DNA-damaging agents (43). These phenotypes are linked to altered DNA damage bypass mechanisms that depend on mono-ubiquitinated PCNA (Proliferating cell nuclear antigen) (44). This evidence highlights the role of oxidative DNA lesions processing in influencing the sensitivity of both canonical and non-canonical HR-deficient cells to PARPis (43) and supports the importance of the cooperation among different repair pathways to maintain genome stability.

A severe risk of genomic instability may derive from chromosomal breakage and rearrangements occurring during the resolution of HR intermediates, specifically double Holliday junctions (dHJs). This is exemplified by the elevated crossover events seen in cells from Bloom syndrome (BS) patients (45). As well as FA, ataxia telangiectasia (A-T) and Nijmegen breakage syndrome (NBS), BS is indeed classified as a severe chromosome instability disorder, characterized by predisposition to cancer (46). Mutations in the RecQ helicase Bloom (BLM), a component of the BTR complex with Topoisomerase III α (TOP3A), RMI1 (RecQ-mediated genome instability protein 1) and RMI2, impair the dissolution of dHJs into non-crossover products, leading to increased crossover events. The regulation of BLM's anti-crossover activity has been extensively studied. Phosphorylation events trigger protein-protein interactions and create a positive feedback loop aimed at suppressing crossover recombination and limiting genomic instability (45). Interestingly, ubiquitination also plays a crucial role in regulating BLM function, being necessary for its recruitment to stalled replication forks (47).

The presence of SSBs and DSBs clearly promotes post-translational modifications such as phosphorylation and ubiquitination. Ubiquitination of chromatin and associated proteins at damage sites coordinates the DDR by signaling the recruitment of repair factors or promoting proteasomal degradation to regulate repair factor levels (48). Ubiquitination also modulates the activity of the repair mechanism. For example, the FA pathway is responsible for repairing DNA ICLs that stall replication forks, primarily functioning through the ubiquitination activity of the RING E3 FA core complex (49). Overall, DDR coordinates cell cycle check point activation and repair

processes, displaying a crucial role to preserve genome stability, as sustained by cancer predispositions in DDR-defective syndromes (46) and by the highly frequent mutations in DDR genes reported in multiple cancers (50,51). A fine-tuned regulation of signaling recruitment and activation of specific factors or pathways is indispensable for a properly functional DDR.

Role of ubiquitination in DNA damage signaling and repair

Ubiquitination coordinates DDR and repair at multiple steps (52–55), from histone modification for chromatin remodeling-dependent recruitment of repair factors, to repair protein ubiquitination for degradation and regulation of their activity. Chromatin remodeling involves a variety of histone modifications, including ubiquitination, which collectively facilitate DDR activation and the repair mechanisms (56–58). Depending on the specific E3 catalyzing H2A modification, different DNA repair pathways may be activated (59,60). The specific lysine residue on histone H2A also influences the progression of the response to DNA damage (56). In addition, H2B ubiquitination is crucial for damage checkpoint activation and initiation of repair (60), while ubiquitination of H3 and H4 plays an important role in response to UV-induced DNA damage (61).

Ubiquitination significantly influences the levels of DDR factors by targeting them for proteasomal degradation. For instance, regulation of p53 by the RING E3 MDM2 through ubiquitination ensures proper turnover of key DNA repair proteins, maintaining a balance between activation and resolution of repair pathways. Additionally, in the presence of mismatches, the levels of hMut (mutator) α protein represent a limiting factor for MMR activity. As for other repair proteins (62), the stability of hMut α is regulated by a RING E3-mediated ubiquitination event (63). In response to UV rays-induced bulking lesions, as well as to oxidative damage causing helix distorting lesions, the NER is activated. At the early stages, the sensor factor XPC (Xeroderma pigmentosum C) is ubiquitinated by the DDB1 (DNA damage-binding protein 1) - CRL4 (Cul (Cullin) 4A-RING E3 ligase) complex, which enhances its affinity binding to the damaged DNA site (64). The role of NER proteins in oxidative damage response is quite well described (35,65), particularly with respect to TC TCR (transcription-coupled)-NER Cockayne syndrome proteins, CSA and CSB (66), which are directly involved in ubiquitin/proteasome degradation events, finally regulating transcription, cell division (67) and DNA repair. CSA is indeed part of the CRL4 complex (CSA/DDB1/Cul4A), while CSB, harbors a ubiquitin-binding domain (68). Their role in recognizing the stalled RNA polymerase II (RNAPII) and in promoting its ubiquitination-mediated proteasomal degradation, as well as that of the cyclic AMP-dependent transcription factor ATF3 (Activating transcription factor 3), is crucial for the proper recruitment and activation of other NER factors (68,69). Therefore, both DDB2 (or XPE) and CSA proteins have been shown to associate with DDB1 and CUL4A, in complexes displaying E3 activity (70). The recruitment of XPC during GG (global genome)-NER relies on chromatin remodeling ubiquitination events mediated by the CUL4/DDB1/ROC1 (regulator of cullins 1) E3 complex, including the ubiquitination of histones H3 and H4 (61,71). DDB1 poly-ubiquitinates XPC and XPE, to increase XPC affinity to DNA or to induce the degradation of XPE (64,72). Finally, CSB is targeted for degradation by the E3 activity of CSA/DDB1/CUL4A complex (73). Interestingly, CSA and CSB are also reported to interact with the tumor suppressor p53, and to regulate its ubiquitination by MDM2 (74). Given the involvement of the tumor suppressor in the regulation of the aging process (26,75), a defect in the ubiquitination activity of MDM2, mediated by dysfunctional CSA and CSB (74), might potentially contribute to the development of the premature aging features observed in CS patients (26,68,74,75).

BER activation in response to oxidative stress is regulated by ubiquitination, involving both RING- and HECT-type E3s, which target multiple DNA glycosylases, finally influencing their activity. As an example, the RING E3 tripartite motif 26 (TRIM26) regulates glycosylase endonuclease III homologue (NTH1) protein levels (76). Moreover, in cooperation with a HECT-type E3, TRIM26 mediates the ubiquitination and degradation of the DNA glycosylase endonuclease VIII-like protein 1 (NEIL1) (77). Additionally, the main BER glycosylase OGG1 (8-

oxoguanine DNA glycosylase-1) is ubiquitinated for proteasomal degradation by a HECT E3 (78). The function of the main DNA breaks sensor, PARP1, determining the recruitment of the BER factor XRCC1 to the sites of the lesions, is critically regulated by post-translational modifications. In particular, ubiquitination events target PARP1 for proteasomal degradation (40). In a similar manner, XRCC1 undergoes multiple modifications, such as poly-ADP-ribosylation and phosphorylation. Importantly, ubiquitination regulates its function and stability, consequently influencing repair efficiency (41,42).

When DDR is needed to be activated, as by the most deleterious form of damage, DSBs, cell cycle checkpoint activation is coordinated by cyclin-dependent kinases (CDKs), which in complex with their cyclin partners, phosphorylate target proteins involved in DDR and repair, finally facilitating DNA resection (79). CDKs activity is firstly kept low in the G1 phase, to progressively increase during the cell cycle (80). The availability of repair factors belonging to one or another pathway is also regulated during cell cycle, contributing to affect their competition, finally aimed at activating a specific repair mechanism. Further elucidating the role of RING-type E3s in DDR, RNF (RING finger protein) 8 and RNF168 are mainly involved in early stages of DSBs response being, together with ATM (ataxia telangiectasia mutated), MDC1 (mediator of DNA damage checkpoint 1) and MRN (MRE11, Meiotic recombination 11, RAD50, and NBS1, Nijmegen breakage syndrome protein 1) complex, among the earliest factors found in DNA damage-induced foci (81). Pointing to their involvement in initiating DSBs-dependent DDR, after the key event of histone H2A variant X (H2AX) phosphorylation (γ -H2AX) (82), RNF8 ubiquitinates γ -H2AX and H2A (83–85), as well as histone H1, thus amplifying ubiquitin-mediated signaling (86). RNF168 is consequently recruited and further propagates histone ubiquitination (83–85). RNF8 and RNF168 form K63-linked ubiquitin chains, which convert the ubiquitinated targets into recruitment platforms for downstream factors. RNF8 and RNF168 sequentially act with E2 ubiquitin-conjugating enzyme Ubc13, to generate histone K63-linked ubiquitin chains (85,87,88). RNF8 induces histones ubiquitination and favors the recruitment of repair proteins (83). It targets E2 Ubc13 to DSBs sites and stimulates poly-ubiquitination through modulating the conformation of ubiquitin linked to Ubc13 (88).

By setting a recruitment platform for DSBs repair factors, RNF8 and RNF168 influence the repair pathway choice in response to DSBs. Chromatin-targeted RNF168 rescues 53BP1 (p53 binding protein 1) recruitment for NHEJ(60), but not BRCA1 (Breast cancer gene 1) for HR (88). Moreover, RNF8 regulates the abundance of NHEJ factors at the sites of damage. In particular, it targets Ku80 for degradation via the formation of K48-linked ubiquitin chains. The depletion of RNF8 affects NHEJ efficiency, causing the prolonged retention of Ku80 at damaged DNA (89). Additionally, RNF8 and RNF168 play an important role at the latest stages of DSB repair, being the E3s which trigger BLM recruitment to sites of replication fork stalling through its ubiquitination, thus impeding excessive HR and protecting from genome instability (47).

As in BLM patients' cells, defects in the FA pathway are related to sensitivity to DNA damage and repair impairment. This mechanism interestingly involves ubiquitination for proper functioning. The RING-type E3 FANCL mono-ubiquitinates the FANCI-FANCD2 (ID2) complex to trigger ICLs removal (49) and favor the recruitment of HR factors to finalize replication after repair (49,90).

Ubiquitination is evidently and strictly involved in the regulation of DSBs repair pathway choice. For example, the DDB1-CUL4 E3 facilitates histone mono-ubiquitination to promote DNA end resection and HR (91,92). Ubiquitination of H2A by BRCA1/BARD1 (BRCA1 associated RING domain 1) promotes HR (59) while ubiquitination by RNF168 favors NHEJ (60). RNF138 also favors end resection and recombination, by regulating Ku70-Ku80 proteins displacement from the DSBs site, thus impairing NHEJ activity (93).

The tumor suppressor protein BRCA1 promotes DNA resection in HR, interacting with CtIP and by opposing to 53BP1 (94,95). In addition, it aids RAD51 loading through interaction with PALB2-BRCA2 (96,97). The RING-type E3s BRCA1 and BARD1 act in a complex, which

regulates histone (H2A and H2AX) ubiquitination and DNA repair factors recruitment on the sites of lesions (52,88,96,98). Its E3 activity involves various factors, including RNAPII, TFIIE (Transcription factor II E), NPM1 (nucleophosmin 1), CtIP (CtBP (C terminus-binding protein)-interacting protein), gamma-tubulin, ER- α (estrogen receptor alpha) and claspin (99). Finally, in the context of the competition and mutual regulation of HR and NHEJ factors, BRCA1 ligase activity antagonizes 53BP1 in the DSBs repair pathway choice by controlling end resection (59). Chromatin and chromatin-associated proteins' ubiquitination is finally essential for both the recruitment of DSBs repair factors and the regulation of their function. Among the RING-type E3s involved in the DSBs response, RNF4 plays a key role by regulating the turnover of the checkpoint mediator MDC1, and of RPA (Replication protein A) at DNA damage sites. In particular, RNF4-depleted cells are reported to fail the effective replacement of RPA by BRCA2 and RAD51 on resected DNA, a critical HR step (100).

Given the involvement of ubiquitination in the regulation of DDR and repair processes, research has focused on the contributions of the E3s, which enable the substrate specificity, and are categorized into distinct families. (101). Scientific reports mainly deal with the contribution of RING-type E3 ligases, whereas the role of HECT-type E3s remains less well understood and requires further investigation.

The HECT E3 family

The HECT E3 family consists of 28 members that are characterized by the presence of a 350 amino acids long C-terminal catalytic domain, namely the HECT domain (16). Unlike RING-type E3s, which directly transfer the ubiquitin molecules from the E2 to the target protein (102), HECT E3s form a covalent thioester intermediate with ubiquitin prior to transferring the ubiquitin molecule to the substrate. The HECT domain itself is structurally divided into two lobes, the N-terminal lobe (N-lobe) and the C-terminal lobe (C-lobe), which are connected by a flexible hinge region, crucial for the catalytic function (Figure 1). The N-lobe interacts with the ubiquitin-bound E2, while the C-lobe harbors the catalytic cysteine responsible for the formation of the thioester intermediate (16,103). The flexible hinge region allows the relative repositioning of the C- and the N-lobe, to bring the catalytic cysteine in proximity to the E2 and the substrate. The structural model from multiple HECT E3s, such as NEDD4 (Neural precursor cell expressed developmentally downregulated 4) (PDB: 2XBF) (104), ITCH (Itchy E3 ubiquitin protein ligase) (PDB: 3TUG) (105), WWP1 (WW domain-containing E3 ubiquitin ligase 1) (PDB: 1ND7) (106), WWP2 (PDB: 4Y07) (107), and HUWE1 (HECT, UBA and WWE domain containing E3 ubiquitin protein ligase 1) (PDB: 3G1N) (108), reveals such conformational flexibility (Figure 1).

HECT E3s-dependent catalytic mechanism and conformational dynamics offer interesting possibilities to the development of selective inhibitors, targeting the catalytic cysteine or disrupting the E2-E3 interaction. Nevertheless, specificity remains a significant hurdle due to the conserved nature of the catalytic domain across HECT ligases. Accordingly, a small molecule, heclin (HECT ligase inhibitor), was found to broadly inhibit HECT E3s by inducing oxidation of the active-site cysteine, without blocking E2 binding (109). Recent advances have identified small-molecule inhibitors of HECT family members, including NEDD4, WWP1-2, SMURF (SMAD ubiquitination regulatory factor) 1-2, and HUWE1, targeting allosteric sites and exploiting the conformational dynamics of the hinge region. Bicyclic peptides selectively target SMURF2, NEDD4, WWP1, and HUWE1, inhibiting their activity *in vitro* (109–111). Interestingly, ITCH-selective inhibitors that interfere with the intramolecular domain arrangement necessary for ubiquitin transfer were developed (112). These inhibitors exhibit promising specificity by stabilizing an inactive conformation of the HECT domain, effectively preventing substrate ubiquitination without affecting other ubiquitin ligase families.

Involvement of HECT-type E3s in DNA damage response

Depending on the presence of structural domains, which are responsible for the recruitment of the substrates, the HECT-type E3 family members are divided into three subfamilies: the NEDD4 (neural precursor cell expressed, developmentally downregulated 4)-like subfamily, characterized by the presence of WW (tryptophan-tryptophan) domains (Figure 2A), the HERC (HECT and RCC1, regulator of chromosome condensation 1) subfamily, characterized by the presence of RLD (RCC1-like domain) domains (Figure 2B) and the “other” HECT-type subfamily, which lacks both RLD and WW domains (Figure 2C) (16).

The fate of the substrate depends on the type of ubiquitin linkages of the polyubiquitin chain. K48 and K11 linkages are mostly involved in targeting substrates for proteasomal degradation, whereas K29, K33 and K63 ubiquitin chains are implicated in autophagic processes (113,114). While the functions of homotypic chains are well-defined, the structure of heterotypic chains has more recently emerged, and their functions need to be further explored (115). Importantly, specific ubiquitin chains (K6, K27 and K63) are reported to affect DDR and repair, influencing chromatin conformation and protein complexes formation, as well as the recruitment of repair factors (116)(Figure 1C).

HECT-type E3s contribute to tumorigenesis by ubiquitinating substrates that function as either tumor suppressors or oncogenes (15,16,103). Being involved in the modification of histones and DDR factors, they are increasingly emerging as key regulators of effective DNA damage responses and genome stability. Nevertheless, our knowledge regarding their specific functions in these mechanisms is relatively limited compared to the extensively studied RING-type E3s. Filling this gap may offer great potential to reveal interesting novel targets for cancer therapy. In the following section, we describe the specific contribution of HECT-type E3s in the DDR and activation of repair pathways.

The NEDD4-like HECT E3 subfamily

The members of the NEDD4-like subfamily present a forty amino acids-long conserved region containing multiple WW domains that recognize substrates proline-rich sequences (PPXY, PPLP, or PPR) (Figure 2A). This family is composed by nine members: NEDD4 (NEDD4-1), NEDD4L (NEDD4-like, NEDD4-2), ITCH (AIP4), WWP1 (AIP5), WWP2 (AIP2), SMURF1, SMURF2, NEDL1 (HECW1, HECT, C2 and WW domain containing E3 ubiquitin protein ligase), and NEDL2 (HECW2) (117). As detailed in the following paragraphs, NEDD4-like E3s play pivotal roles in maintaining genome stability by regulating DDR and repair mechanisms (Figure 3).

NEDD4

NEDD4 is involved in maintaining genome stability by regulating DDR proteins (Figure 3A). It directly or indirectly affects the degradation of tumor suppressors, such as p53 and phosphatase and tensin homolog (PTEN), belonging to the PI3K/AKT (phosphatidyl 3-kinase/protein kinase B, PKB) signaling pathway (118). NEDD4 physically interacts with MDM2, facilitating its stabilization through K63-poly-ubiquitination, in order to reduce p53 stability and activity (119,120).

NEDD4 ubiquitinates PTEN, leading to its degradation and consequent protein level reduction (121).

PTEN participates in the stabilization of RAD51 and BRCA1, as well as in the regulation of the expression of ATM, Chk1 (Checkpoint kinase 1) and 53BP1. Therefore, an altered regulation of its levels by NEDD4 may result in a defective response to DSBs (120–122).

NEDD4 is also involved in RNAPII ubiquitination, thus regulating transcription-coupled repair processes. Accordingly, it favors the ubiquitination of NER proteins within the RING E3 CS complex, further contributing to DDR and protecting from genome stability (123).

NEDD4L

NEDD4L mainly participates in DDR by regulating the stability of DNA repair proteins, thus affecting the final response to the presence of DNA damage (Figure 3B). For instance, NEDD4L-

dependent ubiquitination of 53BP1, induces its degradation. In this way, NEDD4L triggers the shifting of the repair balance toward the more error-free pathway HR (124). NEDD4L also plays a role in controlling apoptosis during DDR. In response to genotoxic stress, it ubiquitinates SMAD2/3, thereby downregulating its protein levels and influencing TGF- β signaling pathways (125).

In addition, NEDD4L mediates the poly-ubiquitination of the glycosylase OGG1 at lysine 341, which leads to its proteasomal degradation and to the impairment of its activity during BER (78). Impacting on BER efficiency, NEDD4L affects DNA repair, particularly under oxidative stress conditions (78,126). On the other hand, in the absence of NEDD4L, prolonged OGG1 stability causes an excessive excision activity, enhancing the formation of SSBs as BER-intermediates (78). Therefore, the E3-dependent regulation of OGG1 protein levels might ensure its balance to prevent genome instability. On the other hand, the expression of NEDD4L itself can be regulated by factors of other repair pathways, such as DDB2, an NER factor with transcriptional activity, shown to downregulate the expression of NEDD4L (78,127). Such example of reciprocal regulation between E3s and repair enzymes underlines the importance of their cooperation for safeguarding genome stability.

ITCH

ITCH participates in the control of transcriptional and apoptotic events, therefore significantly influencing DDR, by modulating the levels and functions of DNA repair and apoptotic factors (Figure 3C). In this regard, ITCH catalyzes the ubiquitination of p73, a member of the tumor suppressor p53 family, for proteasomal degradation (103,128), consequently preventing excessive pro-apoptotic signaling. Its function additionally affects the level of the other family member, p63 (129), involved in development and homeostasis processes of epithelial tissues (130). By modulating the degradation of this substrate, ITCH affects cell proliferation, preventing tumorigenesis. Alternatively, dysregulation of such pathway may result in skin-related diseases and cancer development.

ITCH influences cell survival also by regulating the protein levels of c-FLIP (cellular FLICE-like inhibitory protein), a caspase-8 inhibitor. c-FLIP prevents caspase-mediated apoptosis by interfering with the death-inducing signaling complex (DISC). ITCH-dependent degradation of c-FLIP, restores caspase-8 activity, thereby balancing apoptotic and survival responses to DNA damage (5).

Furthermore, ITCH modifies the regulator of oxidative metabolism thioredoxin-interacting protein (TXNIP) through the formation of K63-linked poly-ubiquitin chains. Notably, the subsequent degradation of TXNIP involves K48-linked ubiquitin chains, indicating a cooperative mechanism among different E3s that modulate oxidative stress responses during DDR (131).

ITCH activity is interestingly modulated by upstream DDR factors, such as ATM, that phosphorylates ITCH in response to DSBs, enhancing its catalytic activity (132).

Furthermore, AKT phosphorylates ITCH, whose nuclear localization, with the subsequent ubiquitination of the linker histone H1.2, impairs RNF8/RNF168-dependent formation of 53BP1 foci (133). In this way, ITCH contributes to chromatin remodeling during DDR, influencing the repair pathway choice in response to DSBs, to finally favor HR.

Given its contribution to preserving genome stability, ITCH emerges as an interesting therapeutic target. Accordingly, a screening study aimed at identifying ITCH inhibitors has discovered a selective compound which blocks autophagic flux and enhances the effectiveness of chemotherapeutic agents, confirming its potential use as target in combined anticancer therapies (112).

WWP1

WWP1 has been shown to contribute to oncogenesis by regulating tumor suppressor proteins and oncoproteins, thereby affecting cell growth, survival, and differentiation (10,16). Its E3 activity

has been linked to multiple effects on stability, function or degradation of substrates with tumor suppressive activities, such as $p27^{Kip1}$ and p53 (10,134,135) (Figure 3D). For example, WWP1 catalyzes Lys-48-linked poly-ubiquitination of $p27^{Kip1}$, favoring its degradation and promoting cell proliferation (136). It has been reported to determine proteasomal degradation of the p53 family member p63 and regulating apoptosis (6). On the other hand, it ubiquitinates the tumor suppressor p53, influencing its activity in the regulation of cell cycle arrest, apoptosis, and repair. In this regard, it has been shown that WWP1-mediated ubiquitination of p53 promotes its export to the cytosol, which results in its decreased transcriptional activities (135). This dysregulation of p53 reduces the expression of its downstream target p21(135).

Another WWP1-mediated nonproteolytic ubiquitination mechanism targets the p53 family member p63 isoform, $\Delta Np63$. Catalyzing the formation of K63-based poly-ubiquitin chain on this substrate, WWP1 serves as modulator of cell proliferation (134). In this way, the oncogenic activity of WWP1 might lead to tumorigenesis and chemoresistance. Furthermore, the downregulation of WWP1 regulates the expression of the apoptotic factors Bcl-2 (B-cell leukemia/lymphoma 2 protein) and BAX (Bcl-2-associated X protein). WWP1 also affects the expression of β -catenin, E-cadherin, MMP (Matrix metalloproteinase) -2, and MMP-9, influencing cell adhesion, migration and invasion, which are critical for metastasis events (137,138).

In addition, WWP1 regulates the cellular redox state, and consequently the basal levels of DNA damage through its interaction with TXNIP. Because TXNIP inhibits the disulfide reductase enzymatic activity of thioredoxin and can affect its antioxidant function, WWP1-mediated ubiquitination of TXNIP modulates redox signaling, which is important for cellular survival under oxidative stress conditions (139). A recent study by Fierro et al. (140) reported that WWP1 is also a critical regulator of DDR, implying that its dysregulation may significantly contribute to therapy resistance, and highlighting its potential as a therapeutic target in cancer. The Authors showed that WWP1 modulates chromatin remodeling and subsequent repair of DNA breaks via modification of the histone demethylase JARID1B, which is involved in the recruitment of DNA repair factors in response to DNA damage (141–143). WWP1 determines the formation of K63-linked poly-ubiquitin chains on JARID1B to positively regulate its stability and potentiating its demethylase activity during DDR-associated chromatin remodeling. Conversely, WWP1 silencing impairs JARID1B-dependent chromatin accessibility of HR and NHEJ factors, to finally delay the repair processes and sensitize cancer cells to chemotherapy(140).

WWP2

WWP2 also contributes to DDR and genomic stability by regulating the levels and function of multiple factors (Figure 3E). WWP2 catalyzes the formation of K48-linked ubiquitin chains on the large subunit of RNAPII, RPB1. During DDR this mechanism importantly results in a transcriptional arrest at the damage site. This event, triggered by the DNA-dependent protein kinase, DNA-PK, impedes transcription to interfere with DNA repair mechanisms (144). Conversely, defects in WWP2 significantly impact on the recruitment of DNA-PK, XRCC4 and DNA Ligase IV, thereby compromising DSBs repair by NHEJ (144).

In response to the presence of DNA breaks, the activity of PARP1 is regulated by WWP2, which performs its ubiquitination for proteasomal degradation (40). By reducing PARP1 levels and, consequently, its PARylation activity, WWP2 prevents a putatively excessive signaling, which can be deleterious, as during isoproterenol-induced cardiac remodeling (40). Coherently, WWP2 downregulation is associated to increased PARylation and aberrant DNA damage signaling (40).

The role of WWP1 in coordinating transcription and repair during DDR, as well as the one in modulating PARP1 activation in response to DNA breaks, points to underlying its critical involvement in preventing genome instability.

SMURF1 and SMURF2

SMURF1 and SMURF2 participate in DDR regulation (Figure 3F-G), particularly downstream of the ATR (ataxia telangiectasia and Rad3-related)/Chk1 signaling pathway, being involved in the regulation of p53-dependent apoptosis in an indirect way. Their activity favors the stabilization of MDM2, which can then ubiquitinate its substrate p53 to promote its proteasomal degradation (145). SMURF1 and SMURF2 additionally influence DDR and apoptosis in a direct way. In response to DNA damage, ATR/Chk1 activation leads to the phosphorylation of SMURF1, which then catalyzes its self-ubiquitination for degradation (146). The reduced levels of SMURF1 lead to stabilization and accumulation of its tumor suppressor substrate RhoB, thereby promoting DNA damage induced apoptotic events (146).

Beyond its role in the regulation of p53 during DDR, (Figure 3G) (145), SMURF2 indirectly influences damage-induced chromatin remodeling. After phosphorylation by ATM, SMURF2 ubiquitinates the RING-type E3 RNF20, which is responsible for H2B ubiquitination (147). Blank and colleagues have reported a colocalization of SMURF2 and RNF20 at γ H2AX foci at DSBs and demonstrated SMURF2-dependent degradation of RNF20, consequently affecting the mono-ubiquitination of H2B (147). On the other hand, SMURF2 counteracts the activity of the RING-type E3 RNF168, promoting the destabilization of H2AX through ubiquitin-mediated proteasomal degradation (148). Therefore, SMURF2 plays a dual role in the modulation of chromatin dynamics during DDR. It promotes chromatin compaction, to protect DNA from damage, but also hinders the recruitment of DNA repair factors. Being activated by ATM, which is involved in DSBs-induced signaling to finally promote repair, by affecting the recruitment of repair factors, SMURF2 participates in a regulating axis which might then exert a negative feedback control of DDR and DSBs repair (149).

NEDL1 and NEDL2

Although the roles of NEDL1 and NEDL2 in DDR are still largely unexplored, current results indicate that NEDL1 interacts with p53, to promote its transcriptional activity and subsequent apoptotic processes (Figure 3H) (150). This evidence suggests a potential role of this HECT E3 in the regulation of the tumor suppressor mediated response to DNA damage.

NEDL2 has been reported to regulate DDR via its interaction with p73 (Figure 3I). Indeed, Miyazaki and colleagues demonstrated that NEDL2 ubiquitinates p73 leading to its stabilization and to the enhancement of its transcriptional activity. In the absence of NEDL2, p73 undergoes faster decay, sustaining that NEDL2 positively regulates p73 stability (151).

The involvement of NEDL1 and NEDL2 in the modulation of the activity of p53-family members highlights them as interesting candidates for further investigations in the DDR context.

The HERC subfamily

The HERC subfamily is composed of six members, divided in large and small E3s. Whereas large HERCs (HERC1 and HERC2) present multiple RLD domains, small HERCs (HERC3-6) possess a single one (152) (Figure 2B). Among the HERC family members, HERC1 and HERC2 are the most extensively studied for their roles in maintaining genome stability through chromatin remodeling, cell cycle regulation, and modulation of DNA repair pathways (Figure 4).

HERC1

HERC1 was the first member of HERC family to be identified, in a study aimed at finding human oncogenic mutations in human breast cancer cells (152). *Herc1* gene is often mutated in metastatic triple negative breast tumors and leukemia (153,154). Upon UV-induced DNA damage, in the presence of the HPV5 β virus E6 protein, HERC1 impedes apoptosis by promoting the degradation of BAK (Bcl-2 homologous antagonist/killer) (7) (Figure 4A).

Considering a direct involvement of this E3 in the regulation of DNA damage repair, it has been shown to impact on MMR, by regulating the protein levels of MSH2 (MutS Homolog 2) (Figure 4A). Given the key role of MSH2 in the detection of replication-induced mismatches and

insertion/deletion (155), the reduction of its levels significantly affects genomic stability. Diouf and colleagues (154) observed that the downregulation of HERC1 is associated with a marked reduction in MSH2 protein levels, suggesting a role in their stabilization, either directly or indirectly. Although the specific mechanism has not been elucidated, they hypothesized that HERC1 may regulate MSH2 through ubiquitination, either preventing its degradation or facilitating its proper cellular localization (154). Given the critical role of MSH2 in maintaining genomic fidelity, HERC1 dysregulation might lead to defective MMR repair, thereby promoting mutations, genomic instability and tumorigenesis.

HERC2

HERC2 plays a key role in modifying chromatin architecture at damage sites and helps signaling to facilitate repair. Of note, HERC2 serves as a regulator of p53 (Figure 4B), by selectively recruiting ATM and ATR kinases, thereby modulating MDM2 and p53 stability throughout the DDR. In addition, the interaction between HERC2 and p53 is necessary for the oligomerization of this transcription factor (156). Despite this evidence, the specific mechanism by which ATM- and ATR-dependent phosphorylation of HERC2 determines the stabilization of p53, or how HERC2 marks p53 for degradation by MDM2, has not been elucidated yet (126).

HERC2 also interacts with RPA, which plays the necessary role of protecting ssDNA from nuclease digestion during repair (157). Under replication stress, RPA2 is phosphorylated by ATR and ubiquitinated by HERC2 for final proteasomal degradation (Figure 4B) (158). This mechanism is essential for the subsequent recruitment of BLM and WRN (Werner) helicases to the RPA complex, thus finally ensuring genomic stability (157,158).

HERC2 is phosphorylated by ATM, ATR or DNA-PK, to promote the formation of a complex with MDC1 and RNF8. The latter is subsequently induced to recruit RNF168 and the E2 enzyme UBE2N to the DSBs. In their proximity, this complex finally targets H2A and H2AX histones with K63-linked ubiquitin chains to further mark damaged DNA. These markers facilitate the recruitment of the DNA repair factors BRCA1, BARD1, RAP80 (Receptor associated protein 80), and 53BP1, ensuring a complete damage response (126,152). Additionally, the E3 activity of HERC2 can target BRCA1 for degradation (Figure 4B), counteracting the stabilizing effect of BARD1. Elevated HERC2 expression in breast tumors suggests that this regulatory mechanism may contribute to carcinogenesis (159).

HERC2 also participates in regulating NER in response to UV-induced DNA lesions. The downregulation of HERC2 results in the stabilization Xeroderma pigmentosum A factor (XPA) and enhances repair activity following cisplatin treatment, indicating that HERC2 post-translationally modulates XPA levels through ubiquitination-mediated proteasomal degradation (Figure 4B) (152,160). Conversely, phosphorylation of XPA by ATM prevents its HERC2-mediated degradation, thereby maintaining a balance between repair and cell cycle progression after damage (161). Dysregulation of this process may compromise DNA repair fidelity, increase mutation rates, and heighten cancer susceptibility (160).

HERC3-6

Currently, our understanding of the roles played by the small HERC family members in the context of DDR and repair mechanisms remains limited. Despite this, their pathological significance and potential functional involvement in DNA repair pathways have been the subject of growing scientific interest, particularly considering their participation in various essential cellular processes such as cell growth regulation, immune responses, and maintenance of cellular homeostasis (152).

Interestingly, mutations in genes which encode for HERC family member proteins have been identified. Frameshift mutations in the HERC3 gene were found in gastric and colorectal carcinomas characterized by microsatellite instability (Figure 4C) (162). In invasive breast and lung cancer, HERC4 was found to be overexpressed, suggesting it as a putative marker for these

types of tumors (163,164). Its overexpression is associated with cell proliferation and growth signaling, particularly through the regulation of SAV1 (Salvador family WW domain containing protein 1), belonging to the Hippo pathway and involved in tumor suppression (Figure 4D) (165). Even though the exact process determining HERC4-mediated regulation of carcinogenesis has not been deeply investigated yet, these findings suggest it has an oncogenic factor, and possible cancer therapeutic target.

HERC5 has also been proposed as a prognostic marker in patients with non-small-cell lung cancer (NSCLC) and hepatocellular carcinoma (HCC) (166,167). Both HERC5 and HERC6 were identified as ISG15 (Interferon-stimulated gene 15) E3s, catalyzing the conjugation of ISG15, a ubiquitin-like modifier, in response to viral infections during antiviral immune responses (Figure 4E-F) (168,169). This function suggests that small HERCs could potentially play interesting roles in safeguarding genomic integrity by modulating DDR, especially when induced by infections-associated DNA damage (170,171).

The “Other” subfamily

The “Other” HECT subfamily is composed of thirteen members, which are: E6AP, E6-associated protein, or UBE3A, ubiquitin protein ligase E3A; UBE3B; UBE3C; HACE1, HECT domain and ankyrin repeat containing E3 ubiquitin protein ligase 1; HECTD1, HECT domain E3 ubiquitin protein ligase 1; HECTD2; HECTD3; HECTD4; AREL1, apoptosis-resistant E3 ubiquitin protein ligase 1; HUWE1, HECT, UBA and WWE domain containing E3 ubiquitin protein ligase 1; G2E3, G2/M-phase-specific E3 ubiquitin protein ligase; UBR5, ubiquitin protein ligase E3 component N-recogin 5; and TRIP12, thyroid hormone receptor-interacting protein 12. Differently from the members of the other families, these E3s lack both WW and RLD domains (Figure 2C). Among these thirteen enzymes, five have been primarily implicated in the regulation of apoptosis and DNA repair pathways (Figure 5).

AREL1

AREL1 is an anti-apoptotic E3 capable of assembling K33-, K48-, and K63-linked poly-ubiquitin chains (116,172). It was reported to be overexpressed in NSCLC and downregulated in multiple cancer cell lines, where it was demonstrated to favor apoptosis. AREL1 presents various ubiquitination substrates, especially among IAP (inhibitor of apoptosis) antagonists, such as SMAC (Second mitochondria-derived activator of caspases), HtrA2 (High-temperature-required protein A2), and ARTS (apoptosis-related protein in the TGF- β signaling pathway). The ubiquitination process catalyzed by AREL1 induces the degradation of such substrates, thus impairing their pro-apoptotic functions and finally aiding cancer cells in evading programmed cell death (Figure 5A) (173).

HUWE1

HUWE1 catalyzes the formation of K6-, K11-, K48-, and K48/K63-linked poly-ubiquitin chains on different substrates involved in the regulation of the balance between cell death and proliferation (174,175). For instance, HUWE1 mediates K63-linked poly-ubiquitination of Dishevelled, modulating Wnt signaling, and of c-Myc, promoting tumor proliferation (Figure 5B) (176,177). Additionally, HUWE1 catalyzes the ubiquitination of the histone deacetylase HDAC2, of the tumor suppressor p53, of Mcl-1 (Myeloid cell leukemia 1) and Cdc6 (Cell division control protein 6) (Figure 5B) (178–181). The interactions between HUWE1 and the anti-apoptotic factor Mcl-1 occurs through a region similar to the Bcl-2 homology 3 (BH3) domain. The subsequent ubiquitination and degradation of Mcl-1 promotes DNA damage-induced apoptosis. Conversely, the downregulation of HUWE1 stabilizes Mcl-1, thereby attenuating apoptosis (179). Following DNA damage induced by UV or alkylating agents, HUWE1 influences the levels of Cdc6, a critical component of pre-replication complexes, thereby impacting on DNA replication and repair processes (180).

HUWE1 also targets HDAC2 for ubiquitination and subsequent degradation. Zhang et al. demonstrated that loss of HUWE1 leads to HDAC2 accumulation, which impairs p53 acetylation, consequently compromising p53-dependent transcriptional activation, stabilization, and apoptotic responses upon DNA damage (Figure 5B) (178). In p53 wild-type cells, inactivation of HUWE1 results in p53 stabilization and activation, indicating that HUWE1 is a key regulator of p53 activity, functioning alongside the canonical MDM2 pathway and ARF (ADP-ribosylation factor)-mediated activation (181).

HUWE1 also targets H2AX (182) and BRCA1 for degradation (Figure 5B) (183). Affecting H2AX dynamics during DDR, it might influence the recruitment of repair factors at the sites of damage. Additionally, HUWE1-mediated proteasomal degradation of BRCA1 might affect HR response to DSBs. Moreover, in the context of BER-dependent response to oxidative damage, together with the RING-E3 TRIM26, HUWE1 targets the BER glycosylase NEIL1 for ubiquitination-dependent degradation (Figure 5B) (77).

G2E3

G2E3 has been identified as a mitotic regulatory protein that is specifically active during the G2/M phase and functions in response to DNA damage (184). Studies using G2E3 knockout mice demonstrated its role in inhibiting apoptosis during early embryonic development, highlighting its importance in cell survival (185). In terms of cellular localization, G2E3 mostly accumulates in the nucleolus but relocates to the nucleoplasm under genotoxic stress conditions, suggesting a direct involvement in DDR (184). A comprehensive siRNA screening targeting 327 human E3s revealed G2E3 as a novel factor involved in damage response. Depletion of G2E3 led to a reduction in γ H2AX levels and impaired phosphorylation of Chk1 following cisplatin treatment, indicating a compromised DNA damage signaling (Figure 5C). Additionally, loss of G2E3 induced apoptosis and resulted in decreased proliferation of cancer cells. Treatment with the nucleoside analogue gemcitabine further revealed that G2E3 depletion causes the accumulation of ssDNA, indicative of replication stress (Figure 5C). Importantly, endogenous G2E3 levels are downregulated in cancer cells following chemotherapy exposure (186). Collectively, these findings suggest that G2E3 participate in DDR, protecting from replication stress and genomic instability (186,187).

TRIP12

TRIP12 was initially identified as a binding partner of the thyroid hormone receptor (188), as well as a factor indirectly involved in p53 degradation (189). Its expression was found to be altered in multiple diseases, underlying it as a potential therapeutic target (190). Regarding its role in DDR, Altmeyer's laboratory demonstrated that TRIP12 binds PARP1 via a PAR-binding domain and catalyzes its poly-ubiquitination for proteasomal degradation (Figure 5D) (191). By regulating PARP1 levels, TRIP12 limits the PARP1-trapping effect of PARPi at sites of DNA breaks, whereas the downregulation of TRIP12 enhances PARPi efficacy, leading to increased DNA replication stress and damage (191). According to such evidence, TRIP12 might be a promising candidate for therapeutic strategies aimed at overcoming resistance mechanisms in BRCA-deficient tumors.

TRIP12 also acts as a tumor suppressor, by modulating the ARF-p53 pathway (192,193). Chen and colleagues demonstrated that knocking down TRIP12 stabilizes ARF in normal human cells, resulting in ARF-dependent p53-mediated growth arrest (Figure 5D) (192).

Additionally, Gudjonsson and collaborators revealed that TRIP12, along with UBR5, regulates the levels of RNF168 (193). Their findings indicate that RNF168 excessively accumulates as DSBs levels increase, especially in dependence on the depletion of TRIP12 and UBR5, thus resulting in the excessive recruitment of 53BP1 and BRCA1 (Figure 5D) (193).

Poulsen and colleagues demonstrated an interesting functional correlation between TRIP12 and HUWE1 in the ubiquitin fusion degradation pathway. Their double knockdown caused an additive stabilization of shared substrates, suggesting that these enzymes operate in parallel

(194). The functional redundancy between TRIP12 and HUWE1 suggests that they may share additional common substrates involved in DDR, further supporting the importance of ubiquitin-mediated events in the maintenance of genome integrity.

UBR5

In addition to its roles in the modulation of RNF168 levels and in histone ubiquitination for DNA damage-associated chromatin remodeling, together with TRIP12 (Figures 5D-E) (193), UBR5 participates in the ubiquitin-proteasome system in tumors. UBR5 is indeed highly expressed in colorectal cancer, where it shows oncogenic functions (195).

Importantly, UBR5 participates in the early stages of DNA repair. The peroxisome proliferator-activated receptor- γ (PPAR γ) enhances UBR5-mediated ubiquitination and subsequent degradation of the ATM-interacting protein ATMIN (Figure 5E) (196). Since ATMIN is essential for the proper localization of 53BP1 under replicative stress, its regulation by UBR5 influences NHEJ efficiency in response to DSBs (Figure 5E) (196,197).

Furthermore, UBR5 collaborates with other E3s to assemble branched ubiquitin chains. It has been reported to facilitate ITCH in forming K48/K63-branched ubiquitin chains on TXNIP, thereby targeting it for degradation (Figure 5E) (131). This suggests a potential indirect role of UBR5 in oxidative damage response regulation. Mechanistically, TXNIP is initially targeted with K63-linked ubiquitin chains, which may serve as a substrate-specific signal to recruit other E3s, such as UBR5 and WWP1, that catalyze K48 linkages (131,139,193). While K63-linked ubiquitination of TXNIP is necessary for its recognition, it is not sufficient alone for proteasomal degradation (131), which can instead be directly induced by the k48-linked ubiquitination activity of WWP1 (139). These findings further sustain the important role of the crosstalk among different E3s in regulating DNA repair, particularly in response to oxidative stress.

Final Remarks and Future Directions

In conclusion, although being less studied than the members of other E3 families, HECT E3s have emerged as regulators of DDR (198), underlying the importance of ubiquitination, as well as of the crosstalk among different E3 families, in the regulation of processes aimed at the maintenance of genome integrity. By regulating the protein levels and functions of their DDR substrates, HECT E3s orchestrate the balance between cell survival and cell death upon genotoxic stress, thus preventing tumorigenesis. Furthermore, most of them show altered expression levels in association with the onset of multiple cancers and therapeutic resistance acquisition.

Notably, proteasome inhibitors show efficacy in cancer therapies (199), confirming the important role of ubiquitin-related enzymes in multiple biological mechanisms, both in physiological and pathological contexts. Proteasomal function is regulated by ubiquitination events, mediated by proteasomal-associated HECT E3s, which alter its substrates recognition and proteolytic activity, importantly, also in response to genotoxic stress (200,201). Investigating the involvement of specific HECT E3s in proteasomal degradation, assembly and function, would further uncover their role in DDR.

Despite the critical roles of HECT E3s, the mechanisms dictating substrate specificity, spatiotemporal regulation, and crosstalk with other E3s or with different post-translational modifying enzymes remain poorly explored. The interplay among members of different subfamilies, determining a variety of particularly characteristic, mixed and branched ubiquitin chains, also offers an interesting opening to additional investigations. Structural studies and high-throughput proteomic approaches could help uncover these regulatory layers. Screenings for HECT E3 inhibitors will putatively aim at identifying molecules, able to selectively target specific ligases. By affecting specific DDR mechanisms, they have good potential to therapeutically achieve improvements for patients' therapies. The integration of screening platforms with

structural-modeling approaches will possibly help to isolate selective inhibitors, finally reducing the risk of off-target effects and associated toxicity.

The development of PROTACs (Proteolysis Targeting Chimeras) represents an innovative strategy in which the targeted degradation of proteins is accomplished by recruiting specific E3s (202,203). Despite such a promising approach, the big obstacle of achieving specificity remains for both inhibitor and PROTAC development, due to the significant structural similarities among various HECT E3s.

The refinement of small molecule inhibitors and PROTACs will possibly enhance specificity and help to develop more effective and combined therapies to overcome resistance. Further investigations on the multiple roles of HECT E3s will then be particularly helpful to improve cancer therapies, especially where resistance remains a persistent hurdle (204).

Targeting such enzymes may modulate DDR and DNA repair pathways in cancer and diseases characterized by genomic instability, to possibly sensitize tumors to genotoxic agents or restoring DNA repair in inherited deficiencies. For instance, targeting HERC2 might affect both BRCA1 and XPA levels, influencing PARPi- or cisplatin-resistant tumors response to therapy. Inhibitors of TRIP12 or WWP2 might positively affect the sensitivity to PARPi of HR defective tumors. Potential inhibitors of WWP1 might also be deeply investigated as therapeutic agents to modulate chromatin accessibility and DDR functionality in cancer. Overall, the combined use of HECT E3s inhibitors with PARPi, oxidative treatments or radiotherapy may help in sensitizing tumors with altered DDR or repair efficiency. Finally, targeting these enzymes, or impeding the interaction with their substrates, may allow the modulation of specific pathways, thus reducing off-target effects.

Acknowledgements

The Research leading to these results has received funding from AIRC to FB (IG 2019 – 23232), to GM (IG 2022 ID 27366; 2023-2027), to EC (IG#31044; 2024-2029). This work has been also supported by the MUR-PNRR M4C2I1.3 PE6 project PE00000019 Heal Italia (CUP: E83C22004670001) to GM, EC, FB; Ministry of Health - HUB LIFE SCIENCE – Advanced Diagnostic- Italian network of excellence for advanced diagnosis (INNOVA) (PNC-E3-2022-23683266) to GM, EC; and Fondazione Luigi Maria Monti IDI-IRCCS (R.C. to EC).

Authors' contributions

S.G. and F.B. conceived the manuscript; S.G. C.F. and V.M wrote the manuscript; F.B., Y.S., E.C. and G.M. revised the manuscript. All the Authors have reviewed and approved this version.

Availability of data and materials

Availability of data and materials is not applicable in this study.

Ethics declarations

Not applicable to this study.

Consent for publication

The authors consent to publication.

Conflict of Interest Statement

There is **NO** conflict of interest.

References

1. Zhang R, Liu B, Tian Y, Xin M, Li Q, Huang X, et al. A chromosome-coupled ubiquitin-proteasome pathway is required for meiotic surveillance. *Cell Death Differ.* 2024 Dec;31(12):1730–45.
2. Sun H, Liu C, Han F, Lin X, Cao L, Liu C, et al. The regulation loop of MARVELD1 interacting with PARP1 in DNA damage response maintains genome stability and promotes therapy resistance of cancer cells. *Cell Death Differ.* 2023 Apr;30(4):922–37.
3. Hershko A, Ciechanover A. The ubiquitin system. *Annu Rev Biochem.* 1998;67:425–79.
4. Oberst A, Malatesta M, Aqeilan RI, Rossi M, Salomoni P, Murillas R, et al. The Nedd4-binding partner 1 (N4BP1) protein is an inhibitor of the E3 ligase Itch. *Proc Natl Acad Sci U S A.* 2007 Jul 3;104(27):11280–5.
5. Le Clorennec C, Lazrek Y, Dubreuil O, Sampaio C, Larbouret C, Lanotte R, et al. ITCH-dependent proteasomal degradation of c-FLIP induced by the anti-HER3 antibody 9F7-F11 promotes DR5/caspase 8-mediated apoptosis of tumor cells. *Cell Communication and Signaling.* 2019 Dec 23;17(1):106.
6. Li Y, Zhou Z, Chen C. WW domain-containing E3 ubiquitin protein ligase 1 targets p63 transcription factor for ubiquitin-mediated proteasomal degradation and regulates apoptosis. *Cell Death Differ.* 2008 Dec 19;15(12):1941–51.
7. Holloway A, Simmonds M, Azad A, Fox JL, Storey A. Resistance to UV-induced apoptosis by β -HPV5 E6 involves targeting of activated BAK for proteolysis by recruitment of the HERC1 ubiquitin ligase. *Int J Cancer.* 2015 Jun 15;136(12):2831–43.
8. Melino G, Cecconi F, Pelicci PG, Mak TW, Bernassola F. Emerging roles of HECT-type E3 ubiquitin ligases in autophagy regulation. *Mol Oncol.* 2019 Oct;13(10):2033–48.
9. Chhangani D, Upadhyay A, Amanullah A, Joshi V, Mishra A. Ubiquitin ligase ITCH recruitment suppresses the aggregation and cellular toxicity of cytoplasmic misfolded proteins. *Sci Rep.* 2014 May 28;4(1):5077.
10. Sanarico AG, Ronchini C, Croce A, Memmi EM, Cammarata UA, De Antoni A, et al. The E3 ubiquitin ligase WWP1 sustains the growth of acute myeloid leukaemia. *Leukemia.* 2018 Apr;32(4):911–9.
11. Meng Y, Sun H, Li Y, Zhao S, Su J, Zeng F, et al. Targeting Ferroptosis by Ubiquitin System Enzymes: A Potential Therapeutic Strategy in Cancer. *Int J Biol Sci.* 2022;18(14):5475–88.
12. Wu Y, Jiao H, Yue Y, He K, Jin Y, Zhang J, et al. Ubiquitin ligase E3 HUWE1/MULE targets transferrin receptor for degradation and suppresses ferroptosis in acute liver injury. *Cell Death Differ.* 2022 Sep 8;29(9):1705–18.
13. Yang Y, Luo M, Zhang K, Zhang J, Gao T, Connell DO, et al. Nedd4 ubiquitylates VDAC2/3 to suppress erastin-induced ferroptosis in melanoma. *Nat Commun.* 2020 Jan 23;11(1):433.
14. Scheffner M, Nuber U, Huibregtse JM. Protein ubiquitination involving an E1–E2–E3 enzyme ubiquitin thioester cascade. *Nature.* 1995 Jan;373(6509):81–3.
15. Bernassola F, Karin M, Ciechanover A, Melino G. The HECT Family of E3 Ubiquitin Ligases: Multiple Players in Cancer Development. *Cancer Cell.* 2008 Jul;14(1):10–21.
16. Bernassola F, Chillemi G, Melino G. HECT-Type E3 Ubiquitin Ligases in Cancer. *Trends Biochem Sci.* 2019 Dec;44(12):1057–75.
17. Feng P, Wang Y, Liu N, Chen Y, Hu Y, Huang Z, et al. High expression of PPP1CC promotes NHEJ-mediated DNA repair leading to radioresistance and poor prognosis in nasopharyngeal carcinoma. *Cell Death Differ.* 2024 May;31(5):683–96.

18. Jackson SP, Bartek J. The DNA-damage response in human biology and disease. *Nature*. 2009 Oct 22;461(7267):1071–8.
19. Chuah YH, Tay EXY, Grinchuk O V, Yoon J, Feng J, Kannan S, et al. CAMK2D serves as a molecular scaffold for RNF8-MAD2 complex to induce mitotic checkpoint in glioma. *Cell Death Differ*. 2023 Aug;30(8):1973–87.
20. Tran AP, Tralie CJ, Reyes J, Moosmüller C, Belkhatir Z, Kevrekidis IG, et al. Long-term p21 and p53 dynamics regulate the frequency of mitosis events and cell cycle arrest following radiation damage. *Cell Death Differ*. 2023 Mar;30(3):660–72.
21. Ciccia A, Elledge SJ. The DNA Damage Response: Making It Safe to Play with Knives. *Mol Cell*. 2010 Oct;40(2):179–204.
22. Lane DP. Cancer. p53, guardian of the genome. *Nature*. 1992 Jul 2;358(6381):15–6.
23. Rodriguez-Pastrana I, Birli E, Coutts AS. p53-dependent DNA repair during the DNA damage response requires actin nucleation by JMY. *Cell Death Differ*. 2023 Jul;30(7):1636–47.
24. Peng Q, Shi X, Li D, Guo J, Zhang X, Zhang X, et al. SCML2 contributes to tumor cell resistance to DNA damage through regulating p53 and CHK1 stability. *Cell Death Differ*. 2023 Jul;30(7):1849–67.
25. Fischer M, Sammons MA. Determinants of p53 DNA binding, gene regulation, and cell fate decisions. *Cell Death Differ*. 2024 Jul;31(7):836–43.
26. Nicolai S, Rossi A, Di Daniele N, Melino G, Annicchiarico-Petruzzelli M, Raschellà G. DNA repair and aging: the impact of the p53 family. *Aging*. 2015 Dec;7(12):1050–65.
27. Billon N, Terrinoni A, Jolicoeur C, McCarthy A, Richardson WD, Melino G, et al. Roles for p53 and p73 during oligodendrocyte development. *Development*. 2004 Mar;131(6):1211–20.
28. Rotblat B, Southwell AL, Ehrnhoefer DE, Skotte NH, Metzler M, Franciosi S, et al. HACE1 reduces oxidative stress and mutant Huntingtin toxicity by promoting the NRF2 response. *Proc Natl Acad Sci U S A*. 2014 Feb 25;111(8):3032–7.
29. Savini I, Catani MV, Rossi A, Duranti G, Melino G, Avigliano L. Characterization of keratinocyte differentiation induced by ascorbic acid: protein kinase C involvement and vitamin C homeostasis. *J Invest Dermatol*. 2002 Feb;118(2):372–9.
30. Wu X, Bayle JH, Olson D, Levine AJ. The p53-mdm-2 autoregulatory feedback loop. *Genes Dev*. 1993 Jul;7(7A):1126–32.
31. Hoeijmakers JHJ. DNA Damage, Aging, and Cancer. *New England Journal of Medicine*. 2009 Oct 8;361(15):1475–85.
32. Zhang H, Jiang L, Du X, Qian Z, Wu G, Jiang Y, et al. The cGAS-Ku80 complex regulates the balance between two end joining subpathways. *Cell Death Differ*. 2024 Jun;31(6):792–803.
33. Miao H, Meng H, Zhang Y, Chen T, Zhang L, Cheng W. FSP1 inhibition enhances olaparib sensitivity in BRCA-proficient ovarian cancer patients via a nonferroptosis mechanism. *Cell Death Differ*. 2024 Apr;31(4):497–510.
34. He L, Liang J, Chen C, Chen J, Shen Y, Sun S, et al. C9orf72 functions in the nucleus to regulate DNA damage repair. *Cell Death Differ*. 2023 Mar;30(3):716–30.
35. Brooks PJ. The 8,5'-cyclopurine-2'-deoxynucleosides: Candidate neurodegenerative DNA lesions in xeroderma pigmentosum, and unique probes of transcription and nucleotide excision repair. *DNA Repair (Amst)*. 2008 Jul;7(7):1168–79.

36. Kee Y, D'Andrea AD. Expanded roles of the Fanconi anemia pathway in preserving genomic stability. *Genes Dev.* 2010 Aug 15;24(16):1680–94.
37. Giovannini S, Weller MC, Repmann S, Moch H, Jiricny J. Synthetic lethality between BRCA1 deficiency and poly(ADP-ribose) polymerase inhibition is modulated by processing of endogenous oxidative DNA damage. *Nucleic Acids Res.* 2019 Sep 26;47(17):9132–43.
38. Caldecott KW. Protein ADP-ribosylation and the cellular response to DNA strand breaks. *DNA Repair (Amst).* 2014 Jul;19:108–13.
39. Adamowicz M, Hailstone R, Demin AA, Komulainen E, Hanzlikova H, Brazina J, et al. XRCC1 protects transcription from toxic PARP1 activity during DNA base excision repair. *Nat Cell Biol.* 2021 Dec 22;23(12):1287–98.
40. Zhang N, Zhang Y, Qian H, Wu S, Cao L, Sun Y. Selective targeting of ubiquitination and degradation of PARP1 by E3 ubiquitin ligase WWP2 regulates isoproterenol-induced cardiac remodeling. *Cell Death Differ.* 2020 Sep 5;27(9):2605–19.
41. Wei L, Nakajima S, Hsieh CL, Kanno S, Masutani M, Levine AS, et al. Damage response of XRCC1 at sites of DNA single strand breaks is regulated by phosphorylation and ubiquitylation after degradation of poly(ADP-ribose). *J Cell Sci.* 2013 Oct 1;126(19):4414–23.
42. Parsons JL, Tait PS, Finch D, Dianova II, Allinson SL, Dianov GL. CHIP-Mediated Degradation and DNA Damage-Dependent Stabilization Regulate Base Excision Repair Proteins. *Mol Cell.* 2008 Feb;29(4):477–87.
43. Giovannini S, Weller MC, Hanzlíková H, Shiota T, Takeda S, Jiricny J. ATAD5 deficiency alters DNA damage metabolism and sensitizes cells to PARP inhibition. *Nucleic Acids Res.* 2020 May 21;48(9):4928–39.
44. Lee K young, Yang K, Cohn MA, Sikdar N, D'Andrea AD, Myung K. Human ELG1 Regulates the Level of Ubiquitinated Proliferating Cell Nuclear Antigen (PCNA) through Its Interactions with PCNA and USP1. *Journal of Biological Chemistry.* 2010 Apr;285(14):10362–9.
45. Balbo Pogliano C, Ceppi I, Giovannini S, Petroulaki V, Palmer N, Uliana F, et al. The CDK1-TOPBP1-PLK1 axis regulates the Bloom's syndrome helicase BLM to suppress crossover recombination in somatic cells. *Sci Adv.* 2022 Feb 4;8(5).
46. Taylor AMR, Rothblum-Oviatt C, Ellis NA, Hickson ID, Meyer S, Crawford TO, et al. Chromosome instability syndromes. *Nat Rev Dis Primers.* 2019 Sep 19;5(1):64.
47. Tikoo S, Madhavan V, Hussain M, Miller ES, Arora P, Zlatanou A, et al. Ubiquitin-dependent recruitment of the Bloom Syndrome helicase upon replication stress is required to suppress homologous recombination. *EMBO J.* 2013 May 24;32(12):1778–92.
48. Dantuma NP, van Attikum H. Spatiotemporal regulation of posttranslational modifications in the <sc>DNA</sc> damage response. *EMBO J.* 2016 Jan 4;35(1):6–23.
49. Smogorzewska A, Matsuoka S, Vinciguerra P, McDonald ER, Hurov KE, Luo J, et al. Identification of the FANCI Protein, a Monoubiquitinated FANCD2 Paralog Required for DNA Repair. *Cell.* 2007 Apr;129(2):289–301.
50. Groelly FJ, Fawkes M, Dagg RA, Blackford AN, Tarsounas M. Targeting DNA damage response pathways in cancer. *Nat Rev Cancer.* 2023 Feb;23(2):78–94.
51. Giovannini S, Smirnov A, Concetti L, Scimeca M, Mauriello A, Bischof J, et al. A comprehensive molecular characterization of a claudin-low luminal B breast tumor. *Biol Direct.* 2024 Aug 16;19(1):66.

52. Koo SY, Park EJ, Noh HJ, Jo SM, Ko BK, Shin HJ, et al. Ubiquitination Links DNA Damage and Repair Signaling to Cancer Metabolism. *Int J Mol Sci.* 2023 May 8;24(9):8441.
53. Cheng M, Lu Y, Wang J, Wang H, Sun Y, Zhao W, et al. The E3 ligase ASB3 downregulates antiviral innate immunity by targeting MAVS for ubiquitin-proteasomal degradation. *Cell Death Differ.* 2024 Dec;31(12):1746–60.
54. Zhang R, Liu B, Tian Y, Xin M, Li Q, Huang X, et al. A chromosome-coupled ubiquitin-proteasome pathway is required for meiotic surveillance. *Cell Death Differ.* 2024 Dec;31(12):1730–45.
55. Bian S, Ni W, Zhou L, Tong Y, Dai C, Zhao X, et al. Ubiquitin-specific protease 1 facilitates hepatocellular carcinoma progression by modulating mitochondrial fission and metabolic reprogramming via cyclin-dependent kinase 5 stabilization. *Cell Death Differ.* 2024 Sep;31(9):1202–18.
56. Uckelmann M, Sixma TK. Histone ubiquitination in the DNA damage response. *DNA Repair (Amst).* 2017 Aug;56:92–101.
57. Kolobynina KG, Rapp A, Cardoso MC. Chromatin Ubiquitination Guides DNA Double Strand Break Signaling and Repair. *Front Cell Dev Biol.* 2022 Jul 5;10.
58. Giovannini S, Weibel L, Schitteck B, Sinnberg T, Schaller M, Lemberg C, et al. Skin Cancer Induction by the Antimycotic Drug Voriconazole Is Caused by Impaired DNA Damage Detection Due to Chromatin Compaction. *Journal of Investigative Dermatology.* 2024 Nov;144(11):2465–76.
59. Densham RM, Garvin AJ, Stone HR, Strachan J, Baldock RA, Daza-Martin M, et al. Human BRCA1-BARD1 ubiquitin ligase activity counteracts chromatin barriers to DNA resection. *Nat Struct Mol Biol.* 2016 Jul;23(7):647–55.
60. Schwertman P, Bekker-Jensen S, Mailand N. Regulation of DNA double-strand break repair by ubiquitin and ubiquitin-like modifiers. *Nat Rev Mol Cell Biol.* 2016 Jun 23;17(6):379–94.
61. Wang H, Zhai L, Xu J, Joo HY, Jackson S, Erdjument-Bromage H, et al. Histone H3 and H4 Ubiquitylation by the CUL4-DDB-ROC1 Ubiquitin Ligase Facilitates Cellular Response to DNA Damage. *Mol Cell.* 2006 May;22(3):383–94.
62. Sweder K, Madura K. Regulation of repair by the 26 S proteasome. *Biomed Res Int.* 2002 Jan 10;2(2):94–105.
63. Hernandez-Pigeon H, Laurent G, Humbert O, Salles B, Lautier D. Degradation of mismatch repair hMutS α heterodimer by the ubiquitin-proteasome pathway. *FEBS Lett.* 2004 Mar 26;562(1–3):40–4.
64. Sugasawa K, Okuda Y, Saijo M, Nishi R, Matsuda N, Chu G, et al. UV-Induced Ubiquitylation of XPC Protein Mediated by UV-DDB-Ubiquitin Ligase Complex. *Cell.* 2005 May;121(3):387–400.
65. Pascucci B, D'Errico M, Parlanti E, Giovannini S, Dogliotti E. Role of nucleotide excision repair proteins in oxidative DNA damage repair: an updating. *Biochemistry (Moscow).* 2011 Jan 16;76(1):4–15.
66. Pascucci B, Lemma T, Iorio E, Giovannini S, Vaz B, Iavarone I, et al. An altered redox balance mediates the hypersensitivity of Cockayne syndrome primary fibroblasts to oxidative stress. *Aging Cell.* 2012 Jun 5;11(3):520–9.
67. Paccosi E, Costanzo F, Costantino M, Balzerano A, Monteonofrio L, Soddu S, et al. The Cockayne syndrome group A and B proteins are part of a ubiquitin-proteasome

- degradation complex regulating cell division. *Proceedings of the National Academy of Sciences*. 2020 Dec 16;117(48):30498–508.
68. Costanzo F, Paccosi E, Proietti-De-Santis L, Egly JM. CS proteins and ubiquitination: orchestrating DNA repair with transcription and cell division. *Trends Cell Biol*. 2024 Jun;
 69. Epanchintsev A, Costanzo F, Rauschendorf MA, Caputo M, Ye T, Donnio LM, et al. Cockayne's Syndrome A and B Proteins Regulate Transcription Arrest after Genotoxic Stress by Promoting ATF3 Degradation. *Mol Cell*. 2017 Dec;68(6):1054-1066.e6.
 70. Groisman R, Polanowska J, Kuraoka I, Sawada J ichi, Saijo M, Drapkin R, et al. The Ubiquitin Ligase Activity in the DDB2 and CSA Complexes Is Differentially Regulated by the COP9 Signalosome in Response to DNA Damage. *Cell*. 2003 May;113(3):357–67.
 71. Sekiguchi M, Matsushita N. DNA Damage Response Regulation by Histone Ubiquitination. *Int J Mol Sci*. 2022 Jul 25;23(15):8187.
 72. Nospikel T. DNA Repair in Mammalian Cells. *Cellular and Molecular Life Sciences*. 2009 Mar 21;66(6):994–1009.
 73. Groisman R, Kuraoka I, Chevallier O, Gaye N, Magnaldo T, Tanaka K, et al. CSA-dependent degradation of CSB by the ubiquitin–proteasome pathway establishes a link between complementation factors of the Cockayne syndrome. *Genes Dev*. 2006 Jun 1;20(11):1429–34.
 74. Latini P, Frontini M, Caputo M, Gregan J, Cipak L, Filippi S, et al. CSA and CSB proteins interact with p53 and regulate its Mdm2-dependent ubiquitination. *Cell Cycle*. 2011 Nov 1;10(21):3719–30.
 75. Wu D, Prives C. Relevance of the p53–MDM2 axis to aging. *Cell Death Differ*. 2018 Jan 1;25(1):169–79.
 76. Williams SC, Parsons JL. NTH1 Is a New Target for Ubiquitylation-Dependent Regulation by TRIM26 Required for the Cellular Response to Oxidative Stress. *Mol Cell Biol*. 2018 Jun 15;38(12).
 77. Edmonds MJ, Carter RJ, Nickson CM, Williams SC, Parsons JL. Ubiquitylation-dependent regulation of NEIL1 by Mule and TRIM26 is required for the cellular DNA damage response. *Nucleic Acids Res*. 2017 Jan 25;45(2):726–38.
 78. Hughes JR, Parsons JL. The E3 Ubiquitin Ligase NEDD4L Targets OGG1 for Ubiquitylation and Modulates the Cellular DNA Damage Response. *Front Cell Dev Biol*. 2020;8:607060.
 79. Falck J, Forment J V, Coates J, Mistrik M, Lukas J, Bartek J, et al. CDK targeting of NBS1 promotes DNA-end resection, replication restart and homologous recombination. *EMBO Rep*. 2012 Jun 8;13(6):561–8.
 80. Hohegger H, Takeda S, Hunt T. Cyclin-dependent kinases and cell-cycle transitions: does one fit all? *Nat Rev Mol Cell Biol*. 2008 Nov 24;9(11):910–6.
 81. Bekker-Jensen S, Mailand N. Assembly and function of DNA double-strand break repair foci in mammalian cells. *DNA Repair (Amst)*. 2010 Dec;9(12):1219–28.
 82. Kinner A, Wu W, Staudt C, Iliakis G. -H2AX in recognition and signaling of DNA double-strand breaks in the context of chromatin. *Nucleic Acids Res*. 2008 Aug 21;36(17):5678–94.
 83. Mailand N, Bekker-Jensen S, Fastrup H, Melander F, Bartek J, Lukas C, et al. RNF8 Ubiquitylates Histones at DNA Double-Strand Breaks and Promotes Assembly of Repair Proteins. *Cell*. 2007 Nov;131(5):887–900.

84. Huen MSY, Grant R, Manke I, Minn K, Yu X, Yaffe MB, et al. RNF8 Transduces the DNA-Damage Signal via Histone Ubiquitylation and Checkpoint Protein Assembly. *Cell*. 2007 Nov;131(5):901–14.
85. Mattioli F, Vissers JHA, van Dijk WJ, Ikpa P, Citterio E, Vermeulen W, et al. RNF168 Ubiquitinates K13-15 on H2A/H2AX to Drive DNA Damage Signaling. *Cell*. 2012 Sep;150(6):1182–95.
86. Thorslund T, Ripplinger A, Hoffmann S, Wild T, Uckelmann M, Villumsen B, et al. Histone H1 couples initiation and amplification of ubiquitin signalling after DNA damage. *Nature*. 2015 Nov 19;527(7578):389–93.
87. Kolas NK, Chapman JR, Nakada S, Ylanko J, Chahwan R, Sweeney FD, et al. Orchestration of the DNA-Damage Response by the RNF8 Ubiquitin Ligase. *Science* (1979). 2007 Dec 7;318(5856):1637–40.
88. Hodge CD, Ismail IH, Edwards RA, Hura GL, Xiao AT, Tainer JA, et al. RNF8 E3 Ubiquitin Ligase Stimulates Ubc13 E2 Conjugating Activity That Is Essential for DNA Double Strand Break Signaling and BRCA1 Tumor Suppressor Recruitment. *Journal of Biological Chemistry*. 2016 Apr;291(18):9396–410.
89. Feng L, Chen J. The E3 ligase RNF8 regulates KU80 removal and NHEJ repair. *Nat Struct Mol Biol*. 2012 Feb 22;19(2):201–6.
90. Tan W, Deans AJ. The ubiquitination machinery of the Fanconi Anemia DNA repair pathway. *Prog Biophys Mol Biol*. 2021 Aug;163:5–13.
91. Bothmer A, Robbiani DF, Feldhahn N, Gazumyan A, Nussenzweig A, Nussenzweig MC. 53BP1 regulates DNA resection and the choice between classical and alternative end joining during class switch recombination. *Journal of Experimental Medicine*. 2010 Apr 12;207(4):855–65.
92. Bunting SF, Callén E, Wong N, Chen HT, Polato F, Gunn A, et al. 53BP1 Inhibits Homologous Recombination in Brca1-Deficient Cells by Blocking Resection of DNA Breaks. *Cell*. 2010 Apr;141(2):243–54.
93. Ismail IH, Gagné JP, Genois MM, Strickfaden H, McDonald D, Xu Z, et al. The RNF138 E3 ligase displaces Ku to promote DNA end resection and regulate DNA repair pathway choice. *Nat Cell Biol*. 2015 Nov 26;17(11):1446–57.
94. Jiang Q, Greenberg RA. Deciphering the BRCA1 Tumor Suppressor Network. *Journal of Biological Chemistry*. 2015 Jul;290(29):17724–32.
95. Panier S, Boulton SJ. Double-strand break repair: 53BP1 comes into focus. *Nat Rev Mol Cell Biol*. 2014 Jan 11;15(1):7–18.
96. Densham RM, Morris JR. The BRCA1 Ubiquitin ligase function sets a new trend for remodelling in DNA repair. *Nucleus*. 2017 Mar 4;8(2):116–25.
97. Zhang F, Fan Q, Ren K, Andreassen PR. PALB2 Functionally Connects the Breast Cancer Susceptibility Proteins BRCA1 and BRCA2. *Molecular Cancer Research*. 2009 Jul 15;7(7):1110–8.
98. Jo EH, Kim MY, Lee HJ, Park HS. Ubiquitin E3 ligases in cancer: somatic mutation and amplification. *BMB Rep*. 2023 May 31;56(5):265–74.
99. Wu W, Koike A, Takeshita T, Ohta T. The ubiquitin E3 ligase activity of BRCA1 and its biological functions. *Cell Div*. 2008 Dec 7;3(1):1.
100. Galanty Y, Belotserkovskaya R, Coates J, Jackson SP. RNF4, a SUMO-targeted ubiquitin E3 ligase, promotes DNA double-strand break repair. *Genes Dev*. 2012 Jun 1;26(11):1179–95.

101. Nakayama KI, Nakayama K. Ubiquitin ligases: cell-cycle control and cancer. *Nat Rev Cancer*. 2006 May 1;6(5):369–81.
102. Plechanovová A, Jaffray EG, Tatham MH, Naismith JH, Hay RT. Structure of a RING E3 ligase and ubiquitin-loaded E2 primed for catalysis. *Nature*. 2012 Sep 29;489(7414):115–20.
103. Delvecchio VS, Fierro C, Giovannini S, Melino G, Bernassola F. Emerging roles of the HECT-type E3 ubiquitin ligases in hematological malignancies. *Discover Oncology*. 2021 Oct 8;12(1):39.
104. Maspero E, Mari S, Valentini E, Musacchio A, Fish A, Pasqualato S, et al. Structure of the HECT:ubiquitin complex and its role in ubiquitin chain elongation. *EMBO Rep*. 2011 Apr 11;12(4):342–9.
105. Zhu K, Shan Z, Chen X, Cai Y, Cui L, Yao W, et al. Allosteric auto-inhibition and activation of the Nedd4 family E3 ligase Itch. *EMBO Rep*. 2017 Sep 26;18(9):1618–30.
106. Verdecia MA, Joazeiro CAP, Wells NJ, Ferrer JL, Bowman ME, Hunter T, et al. Conformational Flexibility Underlies Ubiquitin Ligation Mediated by the WWP1 HECT Domain E3 Ligase. *Mol Cell*. 2003 Jan;11(1):249–59.
107. Gong W, Zhang X, Zhang W, Li J, Li Z. Structure of the HECT domain of human WWP2. *Acta Crystallogr F Struct Biol Commun*. 2015 Oct 1;71(10):1251–7.
108. Pandya RK, Partridge JR, Love KR, Schwartz TU, Ploegh HL. A Structural Element within the HUWE1 HECT Domain Modulates Self-ubiquitination and Substrate Ubiquitination Activities. *Journal of Biological Chemistry*. 2010 Feb;285(8):5664–73.
109. Mund T, Lewis MJ, Maslen S, Pelham HR. Peptide and small molecule inhibitors of HECT-type ubiquitin ligases. *Proceedings of the National Academy of Sciences*. 2014 Nov 25;111(47):16736–41.
110. Rothman AMK, Florentin A, Zink F, Quigley C, Bonneau O, Hemmig R, et al. Therapeutic potential of allosteric HECT E3 ligase inhibition. *Cell*. 2025 May;188(10):2603-2620.e18.
111. Dudev AP, Rigby JM, Hughes GR, Stephenson GR, Storr TE, Chantry A, et al. Expanding the inhibitor space of the WWP1 and WWP2 HECT E3 ligases. *J Enzyme Inhib Med Chem*. 2024 Dec 31;39(1).
112. Rossi M, Rotblat B, Ansell K, Amelio I, Caraglia M, Misso G, et al. High throughput screening for inhibitors of the HECT ubiquitin E3 ligase ITCH identifies antidepressant drugs as regulators of autophagy. *Cell Death Dis*. 2014 May 1;5(5):e1203–e1203.
113. Kwon YT, Ciechanover A. The Ubiquitin Code in the Ubiquitin-Proteasome System and Autophagy. *Trends Biochem Sci*. 2017 Nov;42(11):873–86.
114. Wickliffe KE, Williamson A, Meyer HJ, Kelly A, Rape M. K11-linked ubiquitin chains as novel regulators of cell division. *Trends Cell Biol*. 2011 Nov;21(11):656–63.
115. Gonzalez-Santamarta M, Bouvier C, Rodriguez MS, Xolalpa W. Ubiquitin-chains dynamics and its role regulating crucial cellular processes. *Semin Cell Dev Biol*. 2022 Dec;132:155–70.
116. Singh S, Ng J, Sivaraman J. Exploring the “Other” subfamily of HECT E3-ligases for therapeutic intervention. *Pharmacol Ther*. 2021 Aug;224:107809.
117. Scheffner M, Kumar S. Mammalian HECT ubiquitin-protein ligases: Biological and pathophysiological aspects. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research*. 2014 Jan;1843(1):61–74.

118. Ferretti GDS, Quaas CE, Bertolini I, Zuccotti A, Saatci O, Kashatus JA, et al. HSP70-mediated mitochondrial dynamics and autophagy represent a novel vulnerability in pancreatic cancer. *Cell Death Differ.* 2024 Jul;31(7):881–96.
119. Xu C, Fan CD, Wang X. Regulation of Mdm2 protein stability and the p53 response by NEDD4-1 E3 ligase. *Oncogene.* 2015 Jan 15;34(3):281–9.
120. Lu X, Xu H, Xu J, Lu S, You S, Huang X, et al. The regulatory roles of the E3 ubiquitin ligase NEDD4 family in DNA damage response. *Front Physiol.* 2022 Aug 26;13.
121. Wang X, Trotman LC, Koppie T, Alimonti A, Chen Z, Gao Z, et al. NEDD4-1 is a proto-oncogenic ubiquitin ligase for PTEN. *Cell.* 2007 Jan 12;128(1):129–39.
122. Song MS, Salmena L, Pandolfi PP. The functions and regulation of the PTEN tumour suppressor. *Nat Rev Mol Cell Biol.* 2012 Apr 4;13(5):283–96.
123. Anindya R, Aygün O, Svejstrup JQ. Damage-Induced Ubiquitylation of Human RNA Polymerase II by the Ubiquitin Ligase Nedd4, but Not Cockayne Syndrome Proteins or BRCA1. *Mol Cell.* 2007 Nov;28(3):386–97.
124. Aminu B, Fux J, Mallette E, Petersen N, Zhang W. Targeted Degradation of 53BP1 Using Ubiquitin Variant Induced Proximity. *Biomolecules.* 2022 Mar 22;12(4):479.
125. Gao S, Alarcón C, Sapkota G, Rahman S, Chen PY, Goerner N, et al. Ubiquitin Ligase Nedd4L Targets Activated Smad2/3 to Limit TGF- β Signaling. *Mol Cell.* 2009 Nov;36(3):457–68.
126. Mathieu NA, Levin RH, Spratt DE. Exploring the Roles of HERC2 and the NEDD4L HECT E3 Ubiquitin Ligase Subfamily in p53 Signaling and the DNA Damage Response. *Front Oncol.* 2021 Mar 31;11.
127. Zhao R, Cui T, Han C, Zhang X, He J, Srivastava AK, et al. DDB2 modulates TGF- β signal transduction in human ovarian cancer cells by downregulating NEDD4L. *Nucleic Acids Res.* 2015 Sep 18;43(16):7838–49.
128. Rossi M, De Laurenzi V, Munarriz E, Green DR, Liu YC, Vousden KH, et al. The ubiquitin–protein ligase Itch regulates p73 stability. *EMBO J.* 2005 Feb 23;24(4):836–48.
129. Rossi M, Aqeilan RI, Neale M, Candi E, Salomoni P, Knight RA, et al. The E3 ubiquitin ligase Itch controls the protein stability of p63. *Proceedings of the National Academy of Sciences.* 2006 Aug 22;103(34):12753–8.
130. Li Y, Giovannini S, Wang T, Fang J, Li P, Shao C, et al. p63: a crucial player in epithelial stemness regulation. *Oncogene.* 2023 Nov 10;42(46):3371–84.
131. Ohtake F, Tsuchiya H, Saeki Y, Tanaka K. K63 ubiquitylation triggers proteasomal degradation by seeding branched ubiquitin chains. *Proceedings of the National Academy of Sciences.* 2018 Feb 13;115(7).
132. Santini S, Stagni V, Giambruno R, Fianco G, Di Benedetto A, Mottolese M, et al. ATM kinase activity modulates ITCH E3-ubiquitin ligase activity. *Oncogene.* 2014 Feb 27;33(9):1113–23.
133. Chang L, Shen L, Zhou H, Gao J, Pan H, Zheng L, et al. ITCH nuclear translocation and H1.2 polyubiquitination negatively regulate the DNA damage response. *Nucleic Acids Res.* 2019 Jan 25;47(2):824–42.
134. Peschiaroli A, Scialpi F, Bernassola F, El Sherbini ES, Melino G. The E3 ubiquitin ligase WWP1 regulates Δ Np63-dependent transcription through Lys63 linkages. *Biochem Biophys Res Commun.* 2010 Nov 12;402(2):425–30.
135. Laine A, Ronai Z. Regulation of p53 localization and transcription by the HECT domain E3 ligase WWP1. *Oncogene.* 2007 Mar 1;26(10):1477–83.

136. Cao X, Xue L, Han L, Ma L, Chen T, Tong T. WW Domain-containing E3 Ubiquitin Protein Ligase 1 (WWP1) Delays Cellular Senescence by Promoting p27Kip1 Degradation in Human Diploid Fibroblasts. *Journal of Biological Chemistry*. 2011 Sep;286(38):33447–56.
137. Wu Z, Zan P, Li S, Liu J, Wang J, Chen D, et al. Knockdown of WWP1 inhibits growth and invasion, but induces apoptosis of osteosarcoma cells. *Int J Clin Exp Pathol*. 2015;8(7):7869–77.
138. Hu X, Yu J, Lin Z, Feng R, Wang Z wei, Chen G. The emerging role of WWP1 in cancer development and progression. *Cell Death Discov*. 2021 Jun 21;7(1):163.
139. Giovannini S, Li Y, Pecorari R, Fierro C, Fiorilli C, Corigliano F, et al. Thioredoxin-interacting protein (<scp>TXNIP</scp>) is a substrate of the <scp>NEDD4</scp> -like <scp>E3</scp> ubiquitin-protein ligase <scp>WWP1</scp> in cellular redox state regulation of acute myeloid leukemia cells. *Mol Oncol*. 2025 Jan 4;19(1):133–50.
140. Fierro C, Giovannini S, Moriconi V, Fiorilli C, Panatta E, Frezza V, et al. The WWP1–JARID1B axis sustains acute myeloid leukemia chemoresistance. *Proceedings of the National Academy of Sciences*. 2025 Jul 15;122(28).
141. Li X, Liu L, Yang S, Song N, Zhou X, Gao J, et al. Histone demethylase KDM5B is a key regulator of genome stability. *Proceedings of the National Academy of Sciences*. 2014 May 13;111(19):7096–101.
142. Bayo J, Tran TA, Wang L, Peña-Llopis S, Das AK, Martinez ED. Jumonji Inhibitors Overcome Radioresistance in Cancer through Changes in H3K4 Methylation at Double-Strand Breaks. *Cell Rep*. 2018 Oct;25(4):1040-1050.e5.
143. Xu W, Zhou B, Zhao X, Zhu L, Xu J, Jiang Z, et al. KDM5B demethylates H3K4 to recruit XRCC1 and promote chemoresistance. *Int J Biol Sci*. 2018;14(9):1122–32.
144. Caron P, Pankotai T, Wiegant WW, Tollenaere MAX, Furst A, Bonhomme C, et al. WWP2 ubiquitylates RNA polymerase II for DNA-PK-dependent transcription arrest and repair at DNA breaks. *Genes Dev*. 2019 Jun 1;33(11–12):684–704.
145. Nie J, Xie P, Liu L, Xing G, Chang Z, Yin Y, et al. Smad Ubiquitylation Regulatory Factor 1/2 (Smurf1/2) Promotes p53 Degradation by Stabilizing the E3 Ligase MDM2. *Journal of Biological Chemistry*. 2010 Jul;285(30):22818–30.
146. Wang M, Guo L, Wu Q, Zeng T, Lin Q, Qiao Y, et al. ATR/Chk1/Smurf1 pathway determines cell fate after DNA damage by controlling RhoB abundance. *Nat Commun*. 2014 Sep 24;5(1):4901.
147. Blank M, Tang Y, Yamashita M, Burkett SS, Cheng SY, Zhang YE. A tumor suppressor function of Smurf2 associated with controlling chromatin landscape and genome stability through RNF20. *Nat Med*. 2012 Feb 8;18(2):227–34.
148. Du C, Hansen LJ, Singh SX, Wang F, Sun R, Moure CJ, et al. A PRMT5-RNF168-SMURF2 Axis Controls H2AX Proteostasis. *Cell Rep*. 2019 Sep;28(12):3199–3211.e5.
149. Tang LY, Thomas A, Zhou M, Zhang YE. Phosphorylation of SMURF2 by ATM exerts a negative feedback control of DNA damage response. *Journal of Biological Chemistry*. 2020 Dec;295(52):18485–93.
150. Li Y, Ozaki T, Kikuchi H, Yamamoto H, Ohira M, Nakagawara A. A novel HECT-type E3 ubiquitin protein ligase NEDL1 enhances the p53-mediated apoptotic cell death in its catalytic activity-independent manner. *Oncogene*. 2008 Jun 12;27(26):3700–9.

151. Miyazaki K, Ozaki T, Kato C, Hanamoto T, Fujita T, Irino S, et al. A novel HECT-type E3 ubiquitin ligase, NEDL2, stabilizes p73 and enhances its transcriptional activity. *Biochem Biophys Res Commun*. 2003 Aug;308(1):106–13.
152. Sánchez-Tena S, Cubillos-Rojas M, Schneider T, Rosa JL. Functional and pathological relevance of HERC family proteins: a decade later. *Cellular and Molecular Life Sciences*. 2016 May 22;73(10):1955–68.
153. Craig DW, O’Shaughnessy JA, Kiefer JA, Aldrich J, Sinari S, Moses TM, et al. Genome and Transcriptome Sequencing in Prospective Metastatic Triple-Negative Breast Cancer Uncovers Therapeutic Vulnerabilities. *Mol Cancer Ther*. 2013 Jan 1;12(1):104–16.
154. Diouf B, Cheng Q, Krynetskaia NF, Yang W, Cheok M, Pei D, et al. Somatic deletions of genes regulating MSH2 protein stability cause DNA mismatch repair deficiency and drug resistance in human leukemia cells. *Nat Med*. 2011 Oct 25;17(10):1298–303.
155. Modrich P, Lahue R. MISMATCH REPAIR IN REPLICATION FIDELITY, GENETIC RECOMBINATION, AND CANCER BIOLOGY. *Annu Rev Biochem*. 1996 Jun;65(1):101–33.
156. Cubillos-Rojas M, Amair-Pinedo F, Peiró-Jordán R, Bartrons R, Ventura F, Rosa JL. The E3 Ubiquitin Protein Ligase HERC2 Modulates the Activity of Tumor Protein p53 by Regulating Its Oligomerization. *Journal of Biological Chemistry*. 2014 May;289(21):14782–95.
157. Sala-Gaston J, Martinez-Martinez A, Pedrazza L, Lorenzo-Martín LF, Caloto R, Bustelo XR, et al. HERC Ubiquitin Ligases in Cancer. *Cancers (Basel)*. 2020 Jun 22;12(6):1653.
158. Wu W, Rokutanda N, Takeuchi J, Lai Y, Maruyama R, Togashi Y, et al. HERC2 Facilitates BLM and WRN Helicase Complex Interaction with RPA to Suppress G-Quadruplex DNA. *Cancer Res*. 2018 Nov 15;78(22):6371–85.
159. Wu W, Sato K, Koike A, Nishikawa H, Koizumi H, Venkitaraman AR, et al. HERC2 Is an E3 Ligase That Targets BRCA1 for Degradation. *Cancer Res*. 2010 Aug 1;70(15):6384–92.
160. Kang TH, Reardon JT, Sancar A. Regulation of nucleotide excision repair activity by transcriptional and post-transcriptional control of the XPA protein. *Nucleic Acids Res*. 2011 Apr 1;39(8):3176–87.
161. Lee TH, Park JM, Leem SH, Kang TH. Coordinated regulation of XPA stability by ATR and HERC2 during nucleotide excision repair. *Oncogene*. 2014 Jan 2;33(1):19–25.
162. Yoo NJ, Park SW, Lee SH. Frameshift mutations of ubiquitination-related genes HERC2, HERC3, TRIP12, UBE2Q1 and UBE4B in gastric and colorectal carcinomas with microsatellite instability. *Pathology*. 2011 Dec;43(7):753–5.
163. Zhou H, Shi R, Wei M, Zheng WL, Zhou JY, Ma WL. The expression and clinical significance of HERC4 in breast cancer. *Cancer Cell Int*. 2013;13(1):113.
164. Zeng WL, Chen YW, Zhou H, Zhou JY, Wei M, Shi R. Expression of HERC4 in Lung Cancer and its Correlation with Clinicopathological Parameters. *Asian Pacific Journal of Cancer Prevention*. 2015 Feb 25;16(2):513–7.
165. Aerne BL, Gailite I, Sims D, Tapon N. Hippo Stabilises Its Adaptor Salvador by Antagonising the HECT Ubiquitin Ligase Herc4. *PLoS One*. 2015 Jun 30;10(6):e0131113.
166. Wrage M, Haggmann W, Kemming D, Uzunoglu FG, Riethdorf S, Effenberger K, et al. Identification of HERC5 and its potential role in NSCLC progression. *Int J Cancer*. 2015 May 15;136(10):2264–72.

167. Xue F, Higgs BW, Huang J, Morehouse C, Zhu W, Yao X, et al. HERC5 is a prognostic biomarker for post-liver transplant recurrent human hepatocellular carcinoma. *J Transl Med.* 2015 Dec 11;13(1):379.
168. Dastur A, Beaudenon S, Kelley M, Krug RM, Huibregtse JM. Herc5, an Interferon-induced HECT E3 Enzyme, Is Required for Conjugation of ISG15 in Human Cells. *Journal of Biological Chemistry.* 2006 Feb;281(7):4334–8.
169. Versteeg GA, Hale BG, van Boheemen S, Wolff T, Lenschow DJ, García-Sastre A. Species-Specific Antagonism of Host ISGylation by the Influenza B Virus NS1 Protein. *J Virol.* 2010 May 15;84(10):5423–30.
170. Chumduri C, Gurumurthy RK, Zietlow R, Meyer TF. Subversion of host genome integrity by bacterial pathogens. *Nat Rev Mol Cell Biol.* 2016 Oct 18;17(10):659–73.
171. Weitzman MD, Weitzman JB. What's the Damage? The Impact of Pathogens on Pathways that Maintain Host Genome Integrity. *Cell Host Microbe.* 2014 Mar;15(3):283–94.
172. Kristariyanto YA, Choi SY, Rehman SAA, Ritorto MS, Campbell DG, Morrice NA, et al. Assembly and structure of Lys33-linked polyubiquitin reveals distinct conformations. *Biochemical Journal.* 2015 Apr 15;467(2):345–52.
173. Kim J bin, Kim SY, Kim BM, Lee H, Kim I, Yun J, et al. Identification of a Novel Anti-apoptotic E3 Ubiquitin Ligase That Ubiquitinates Antagonists of Inhibitor of Apoptosis Proteins SMAC, HtrA2, and ARTS. *Journal of Biological Chemistry.* 2013 Apr;288(17):12014–21.
174. Lin IH, Li YR, Chang CH, Cheng YW, Wang YT, Tsai YS, et al. Regulation of primary cilia disassembly through HUWE1-mediated TTBK2 degradation plays a crucial role in cerebellar development and medulloblastoma growth. *Cell Death Differ.* 2024 Oct;31(10):1349–61.
175. Huang X, Huo L, Xiao B, Ouyang Y, Chen F, Li J, et al. Activating STING/TBK1 suppresses tumor growth via degrading HPV16/18 E7 oncoproteins in cervical cancer. *Cell Death Differ.* 2024 Jan;31(1):78–89.
176. Adhikary S, Marinoni F, Hock A, Hulleman E, Popov N, Beier R, et al. The Ubiquitin Ligase HectH9 Regulates Transcriptional Activation by Myc and Is Essential for Tumor Cell Proliferation. *Cell.* 2005 Nov;123(3):409–21.
177. de Groot REA, Ganji RS, Bernatik O, Lloyd-Lewis B, Seipel K, Šedová K, et al. Huwe1-Mediated Ubiquitylation of Dishevelled Defines a Negative Feedback Loop in the Wnt Signaling Pathway. *Sci Signal.* 2014 Mar 18;7(317).
178. Zhang J, Kan S, Huang B, Hao Z, Mak TW, Zhong Q. Mule determines the apoptotic response to HDAC inhibitors by targeted ubiquitination and destruction of HDAC2. *Genes Dev.* 2011 Dec 15;25(24):2610–8.
179. Zhong Q, Gao W, Du F, Wang X. Mule/ARF-BP1, a BH3-Only E3 Ubiquitin Ligase, Catalyzes the Polyubiquitination of Mcl-1 and Regulates Apoptosis. *Cell.* 2005 Jul;121(7):1085–95.
180. Hall JR, Kow E, Nevis KR, Lu CK, Luce KS, Zhong Q, et al. Cdc6 Stability Is Regulated by the Huwe1 Ubiquitin Ligase after DNA Damage. *Mol Biol Cell.* 2007 Sep;18(9):3340–50.
181. Chen D, Kon N, Li M, Zhang W, Qin J, Gu W. ARF-BP1/Mule Is a Critical Mediator of the ARF Tumor Suppressor. *Cell.* 2005 Jul;121(7):1071–83.

182. Atsumi Y, Fujimori H, Fukuda H, Inase A, Shinohe K, Yoshioka Y, et al. Onset of Quiescence Following p53 Mediated Down-Regulation of H2AX in Normal Cells. *PLoS One*. 2011 Aug 12;6(8):e23432.
183. Wang X, Lu G, Li L, Yi J, Yan K, Wang Y, et al. HUWE1 interacts with BRCA1 and promotes its degradation in the ubiquitin–proteasome pathway. *Biochem Biophys Res Commun*. 2014 Feb;444(3):290–5.
184. Brooks WS, Banerjee S, Crawford DF. G2E3 is a nucleo-cytoplasmic shuttling protein with DNA damage responsive localization. *Exp Cell Res*. 2007 Feb;313(4):665–76.
185. Brooks WS, Helton ES, Banerjee S, Venable M, Johnson L, Schoeb TR, et al. G2E3 Is a Dual Function Ubiquitin Ligase Required for Early Embryonic Development. *Journal of Biological Chemistry*. 2008 Aug;283(32):22304–15.
186. Schmidt F, Kunze M, Looock AC, Dobbstein M. Screening analysis of ubiquitin ligases reveals G2E3 as a potential target for chemosensitizing cancer cells. *Oncotarget*. 2015 Jan 20;6(2):617–32.
187. Schmidt F, Karnitz LM, Dobbstein M. G2E3 attenuating replicative stress. *Aging*. 2015 Aug 18;7(8):527–8.
188. Lee JW, Choi HS, Gyuris J, Brent R, Moore DD. Two classes of proteins dependent on either the presence or absence of thyroid hormone for interaction with the thyroid hormone receptor. *Molecular Endocrinology*. 1995 Feb;9(2):243–54.
189. Huibregtse JM, Scheffner M, Beaudenon S, Howley PM. A family of proteins structurally and functionally related to the E6-AP ubiquitin-protein ligase. *Proc Natl Acad Sci U S A*. 1995 May 23;92(11):5249.
190. Brunet M, Vargas C, Larriou D, Torrisani J, Dufresne M. E3 Ubiquitin Ligase TRIP12: Regulation, Structure, and Physiopathological Functions. *Int J Mol Sci*. 2020 Nov 12;21(22).
191. Gatti M, Imhof R, Huang Q, Baudis M, Altmeyer M. The Ubiquitin Ligase TRIP12 Limits PARP1 Trapping and Constrains PARP Inhibitor Efficiency. *Cell Rep*. 2020 Aug;32(5):107985.
192. Chen D, Shan J, Zhu WG, Qin J, Gu W. Transcription-independent ARF regulation in oncogenic stress-mediated p53 responses. *Nature*. 2010 Mar 7;464(7288):624–7.
193. Gudjonsson T, Altmeyer M, Savic V, Toledo L, Dinant C, Grøfte M, et al. TRIP12 and UBR5 Suppress Spreading of Chromatin Ubiquitylation at Damaged Chromosomes. *Cell*. 2012 Aug;150(4):697–709.
194. Poulsen EG, Steinhauer C, Lees M, Lauridsen AM, Ellgaard L, Hartmann-Petersen R. HUWE1 and TRIP12 Collaborate in Degradation of Ubiquitin-Fusion Proteins and Misframed Ubiquitin. *PLoS One*. 2012 Nov 27;7(11):e50548.
195. Xie Z, Liang H, Wang J, Xu X, Zhu Y, Guo A, et al. Significance of the E3 ubiquitin protein UBR5 as an oncogene and a prognostic biomarker in colorectal cancer. *Oncotarget*. 2017 Dec 8;8(64):108079–92.
196. Li CG, Mahon C, Sweeney NM, Verschueren E, Kantamani V, Li D, et al. PPAR γ Interaction with UBR5/ATMIN Promotes DNA Repair to Maintain Endothelial Homeostasis. *Cell Rep*. 2019 Jan;26(5):1333-1343.e7.
197. Schmidt L, Wiedner M, Velimezi G, Prochazkova J, Owusu M, Bauer S, et al. ATMIN is required for the ATM-mediated signaling and recruitment of 53BP1 to DNA damage sites upon replication stress. *DNA Repair (Amst)*. 2014 Dec;24:122–30.

198. Vitale I, Pietrocola F, Guilbaud E, Aaronson SA, Abrams JM, Adam D, et al. Apoptotic cell death in disease—Current understanding of the NCCD 2023. *Cell Death Differ.* 2023 May;30(5):1097–154.
199. Goldberg AL. Development of proteasome inhibitors as research tools and cancer drugs. *Journal of Cell Biology.* 2012 Nov 12;199(4):583–8.
200. Wang Z, Spoel SH. HECT ubiquitin ligases as accessory proteins of the plant proteasome. *Essays Biochem.* 2022 Aug 5;66(2):135–45.
201. Moiseeva TN, Bottrill A, Melino G, Barlev NA. DNA damage-induced ubiquitylation of proteasome controls its proteolytic activity. *Oncotarget.* 2013 Sep 30;4(9):1338–48.
202. Békés M, Langley DR, Crews CM. PROTAC targeted protein degraders: the past is prologue. *Nat Rev Drug Discov.* 2022 Mar 18;21(3):181–200.
203. Ishida T, Ciulli A. E3 Ligase Ligands for PROTACs: How They Were Found and How to Discover New Ones. *SLAS Discovery.* 2021 Apr;26(4):484–502.
204. Landré V, Rotblat B, Melino S, Bernassola F, Melino G. Screening for E3-ubiquitin ligase inhibitors: challenges and opportunities. *Oncotarget.* 2014 Sep 30;5(18):7988–8013.
205. Cassidy KB, Bang S, Kurokawa M, Gerber SA. Direct regulation of Chk1 protein stability by E3 ubiquitin ligase HUWE1. *FEBS J.* 2020 May 29;287(10):1985–99.
206. Kwon YT, Ciechanover A. The Ubiquitin Code in the Ubiquitin-Proteasome System and Autophagy. *Trends Biochem Sci.* 2017 Nov;42(11):873–86.
207. Gatti M, Pinato S, Maiolica A, Rocchio F, Prato MG, Aebersold R, et al. RNF168 Promotes Noncanonical K27 Ubiquitination to Signal DNA Damage. *Cell Rep.* 2015 Jan;10(2):226–38.
208. Kulathu Y, Komander D. Atypical ubiquitylation — the unexplored world of polyubiquitin beyond Lys48 and Lys63 linkages. *Nat Rev Mol Cell Biol.* 2012 Aug 23;13(8):508–23.
209. Deng L, Meng T, Chen L, Wei W, Wang P. The role of ubiquitination in tumorigenesis and targeted drug discovery. *Signal Transduct Target Ther.* 2020 Feb 29;5(1):11.
210. Jiricny J. Postreplicative Mismatch Repair. *Cold Spring Harb Perspect Biol.* 2013 Apr 1;5(4):a012633–a012633.
211. Kunkel TA, Erie DA. DNA MISMATCH REPAIR. *Annu Rev Biochem.* 2005 Jun 1;74(1):681–710.
212. Arlow T, Kim J, Haye-Bertolozzi JE, Martínez CB, Fay C, Zorensky E, et al. MutS α mismatch repair protein stability is governed by subunit interaction, acetylation, and ubiquitination. *G3 Genes|Genomes|Genetics.* 2021 Apr 23;11(3).
213. Cleaver JE, Lam ET, Revet I. Disorders of nucleotide excision repair: the genetic and molecular basis of heterogeneity. *Nat Rev Genet.* 2009 Nov 7;10(11):756–68.
214. Bregman DB, Halaban R, van Gool AJ, Henning KA, Friedberg EC, Warren SL. UV-induced ubiquitination of RNA polymerase II: a novel modification deficient in Cockayne syndrome cells. *Proceedings of the National Academy of Sciences.* 1996 Oct 15;93(21):11586–90.
215. Aamann MD, Muftuoglu M, Bohr VA, Stevnsner T. Multiple interaction partners for Cockayne syndrome proteins: Implications for genome and transcriptome maintenance. *Mech Ageing Dev.* 2013 May;134(5–6):212–24.
216. Fortini P, Parlanti E, Sidorkina OM, Laval J, Dogliotti E. The Type of DNA Glycosylase Determines the Base Excision Repair Pathway in Mammalian Cells. *Journal of Biological Chemistry.* 1999 May;274(21):15230–6.

217. Dogliotti E, Fortini P, Pascucci B, Parlanti E. The mechanism of switching among multiple BER pathways. In 2001. p. 3–27.
218. Bennett RAO, Demple B. DNA Base Excision Repair Pathways. In: Encyclopedia of Biological Chemistry. Elsevier; 2013. p. 1–8.
219. Ceccaldi R, Rondinelli B, D'Andrea AD. Repair Pathway Choices and Consequences at the Double-Strand Break. *Trends Cell Biol.* 2016 Jan;26(1):52–64.
220. MAO Z, BOZZELLA M, SELUANOV A, GORBUNOVA V. Comparison of nonhomologous end joining and homologous recombination in human cells. *DNA Repair (Amst).* 2008 Oct 1;7(10):1765–71.
221. Daley JM, Sung P. 53BP1, BRCA1, and the Choice between Recombination and End Joining at DNA Double-Strand Breaks. *Mol Cell Biol.* 2014 Apr 1;34(8):1380–8.
222. Krejci L, Altmannova V, Spirek M, Zhao X. Homologous recombination and its regulation. *Nucleic Acids Res.* 2012 Jul 1;40(13):5795–818.
223. Huertas P, Jackson SP. Human CtIP mediates cell cycle control of DNA end resection and double strand break repair. *J Biol Chem.* 2009 Apr 3;284(14):9558–65.
224. Khanna KK, Jackson SP. DNA double-strand breaks: signaling, repair and the cancer connection. *Nat Genet.* 2001 Mar;27(3):247–54.
225. Chang HHY, Pannunzio NR, Adachi N, Lieber MR. Non-homologous DNA end joining and alternative pathways to double-strand break repair. *Nat Rev Mol Cell Biol.* 2017 Aug 17;18(8):495–506.
226. Jackson SP. Sensing and repairing DNA double-strand breaks. *Carcinogenesis.* 2002 May 1;23(5):687–96.
227. Caracciolo D, Riillo C, Di Martino MT, Tagliaferri P, Tassone P. Alternative Non-Homologous End-Joining: Error-Prone DNA Repair as Cancer's Achilles' Heel. *Cancers (Basel).* 2021 Mar 19;13(6):1392.
228. Iliakis G, Murmann T, Soni A. Alternative end-joining repair pathways are the ultimate backup for abrogated classical non-homologous end-joining and homologous recombination repair: Implications for the formation of chromosome translocations. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis.* 2015 Nov;793:166–75.
229. Yang G, Liu C, Chen SH, Kassab MA, Hoff JD, Walter NG, et al. Super-resolution imaging identifies PARP1 and the Ku complex acting as DNA double-strand break sensors. *Nucleic Acids Res.* 2018 Apr 20;46(7):3446–57.
230. Ceccaldi R, Liu JC, Amunugama R, Hajdu I, Primack B, Petalcorin MIR, et al. Homologous-recombination-deficient tumours are dependent on Polθ-mediated repair. *Nature.* 2015 Feb 12;518(7538):258–62.
231. Lu G, Duan J, Shu S, Wang X, Gao L, Guo J, et al. Ligase I and ligase III mediate the DNA double-strand break ligation in alternative end-joining. *Proceedings of the National Academy of Sciences.* 2016 Feb 2;113(5):1256–60.
232. Paddock MN, Bauman AT, Higdon R, Kolker E, Takeda S, Scharenberg AM. Competition between PARP-1 and Ku70 control the decision between high-fidelity and mutagenic DNA repair. *DNA Repair (Amst).* 2011 Mar 7;10(3):338–43.
233. Ahrabi S, Sarkar S, Pfister SX, Pirovano G, Higgins GS, Porter ACG, et al. A role for human homologous recombination factors in suppressing microhomology-mediated end joining. *Nucleic Acids Res.* 2016 Jul 8;44(12):5743–57.

234. Lu H, Shamanna RA, de Freitas JK, Okur M, Khadka P, Kulikowicz T, et al. Cell cycle-dependent phosphorylation regulates RECQL4 pathway choice and ubiquitination in DNA double-strand break repair. *Nat Commun.* 2017 Dec 11;8(1):2039.
235. Moldovan GL, D'Andrea AD. How the Fanconi Anemia Pathway Guards the Genome. *Annu Rev Genet.* 2009 Dec 1;43(1):223–49.
236. Thompson LH, Hinz JM. Cellular and molecular consequences of defective Fanconi anemia proteins in replication-coupled DNA repair: Mechanistic insights. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis.* 2009 Jul;668(1–2):54–72.

BOX 1- The multiple roles of ubiquitination

Ubiquitination is a post-translational modification consisting in the attachment of ubiquitin molecule to specific lysine residues of protein substrates (3). This mechanism is involved in a variety of cellular processes, such as protein degradation, signaling and DNA repair(78,152,205). As shown in the figure, a ubiquitin-activating E1 enzyme starts the reaction by the ATP-dependent formation of a thioester bond between its active cysteine and the ubiquitin molecule. The ubiquitin is afterwards transferred to a conjugating E2 enzyme prior to the recruitment of the ubiquitin-loaded E2 to a specific substrate by a E3 ligase (Box 1 figure). The precise regulation of this mechanism is triggered by this one last enzyme, which recognizes specific structural features of the substrate, to finally transfer a ubiquitin molecule to a particular lysine residue.

This reaction consists of a mono-ubiquitination, whereas poly-ubiquitination occurs when multiple ubiquitin molecules are subsequently attached. As in the figure, poly-ubiquitin chains can be assembled in various topologies, which finally influence the fate and function of the substrate (Box 1 figure) (14,15). To form poly-ubiquitin chains, ubiquitin molecules are linked through various lysine (K) residues (K6, K11, K27, K29, K33, K48, K63) (206), to finally determining a specific fate for the substrate, such as proteasomal degradation, functional alterations, subcellular localization, or formation of protein complexes.

Interestingly, specific linkages are associated with particular functions. For instance, K6-linked chains are involved in mitochondrial autophagy (mitophagy) and DDR (205), K27-linked chains, participate in the recruitment of DDR factors and in the modulation of immune responses (207). On the other hand, K48-linked as well as K11-linked chains are common signals for proteasomal degradation, marking substrates for the 26S proteasome. Differently, K63-linked chains are often involved in non-proteolytic roles such as DNA repair and signal transduction. The structural differences among these poly-ubiquitin chains then influence the activity, localization, or interaction properties of substrate proteins, thus regulating signaling pathways also involved in cell cycle progression, apoptosis, DDR and repair (15,208).

Ubiquitination is evidently critical to regulate the level and the function of proteins. The diversity of the ubiquitin linkages and the substrate specificity provided by the E3s affect a variety of processes. Depending on the function of the ubiquitination target and the effects of the modification, E3s are indeed involved in DDR, cell cycle regulation and cell death (15,16,209). Their ability to modulate the stability of key DDR factors, influencing chromatin remodeling at sites of damage, or facilitating the recruitment of repair factors, underlines the importance of the E3s in influencing cancer and therapies response, thereby contributing to protect genome integrity (15,16,98,103,209).

BOX 2- DNA Damage Response (DDR) and Repair

MMR includes factors such as MutS (mutator S) α and β (MutS protein homolog MSH2-MSH6 and MSH2-MSH3), recognizing mismatches induced by DNA polymerases, that escape proofreading (210) and insertion/deletion loops (IDLs), resulting from errors during replication or recombination (31,210). After recognition of a DNA lesion, the excision of the patch containing it occurs by the incision activity of MutL (MLH, mutator L homolog) complex (MutL α (MLH1-PMS2, Postmeiotic segregation increased 2 complex), MutL β (MLH1-PMS1), or MutL γ (MLH1-MLH3)). The strand is finally corrected by excision, DNA synthesis and ligation, involving PCNA (Proliferating cell nuclear antigen), RPA (replication protein A), RFC (replication factor C), Exonuclease I (Exo I), DNA polymerases δ and ϵ , Flap endonuclease 1 (FEN1) and additional factors (210,211).

The stabilization of MMR proteins by post translational modifications and proteasomal degradation (63,212), underlines the importance of ubiquitination events in the regulation of repair proteins' function.

Crosslinks and helix-distorting DNA adducts, such as 6-4 photoproducts (6-4PPs) and pyrimidine dimers (CPDs) induced by UV rays, activate NER, which is divided in GG (global genome) -NER and TCR (transcription-coupled) -NER, depending on whether the damage occurs in a transcriptionally active or inactive domain (31,213). In GG-NER, DNA lesions are detected by the Xeroderma pigmentosum factor C (XPC) - hRad23B/A protein complex and by the DDB1 (DNA damage-binding protein 1) -DDB2 (or XPE) complex, while the arrested transcription apparatus signals the DNA damage in TC-NER, with the involvement of Cockayne syndrome (CS) proteins CSA and CSB (31,72,213). Their participation in DNA damage-induced ubiquitination events triggers the subsequent repair process, ensuring the removal of RNA polymerase II (RNAPII) for degradation (68,70,214). The transcription factor TFIIH (transcription factor II H) acts through the helicase function of its XPB and XPD subunits. The excision of a 30 nucleotides-long DNA fragment containing the lesion occurs after the incision step performed by XPF and XPG nucleases. The DNA Pol δ/ϵ , in cooperation with PCNA and RFC, synthesize a new DNA fragment. Finally, Ligasi I, or Ligase III with the help of the BER scaffold protein XRCC1 (X-ray repair cross-complementing protein 1), operates the sealing step (31,72,213). Interestingly, ubiquitination also regulates the late stages of TC-NER, when CSB is degraded by the activity of the CSA including E3 complex (73).

Notably, the NER pathway is also activated by the oxidatively induced bulky lesions cyclopurines, which represent an exceptional type of DNA damage, considering the common view that oxidative DNA lesions are mostly repaired by the BER (35). Indeed, NER factors cooperate with BER proteins in response to oxidative stress. For instance, CSB interacts with multiple BER factors, to improve their recruitment or stimulating their function, both in the nuclear and in the mitochondrial compartment (215).

In BER, a DNA glycosylase excises the damaged base leaving an AP (apurinic/aprimidinic) site and an endonuclease creates a SSB (single-strand break). In the short-patch sub-pathway, one nucleotide is replaced by Pol β and the gap is sealed by XRCC1-Ligase III α . During the long-patch BER, Pol δ/ϵ synthesize 2-10 nucleotides and FEN1 removes the 5' flap DNA, whereas the ligation is performed by the PCNA-Ligase I complex (216,217).

During the repair process, new types of DNA damage may also arise. For example, oxidation of DNA bases caused by ROS (reactive oxygen species) activates mono- or bi-functional glycosylases involved in BER (216,217). Among bifunctional glycosylases, 8-oxoguanine DNA glycosylase-1 (OGG1) specifically responds to guanine oxidation by cleaving both the glycosidic bond and the phosphodiester backbone (216,218), with its activity regulated by ubiquitination (78). In the case of mono-functional glycosylases like MutY DNA glycosylase (MUTYH), the process involves recruiting an AP endonuclease, such as APE1, which then cleaves the phosphodiester bond, creating a SSB as an intermediate in BER (218).

SSBs are recognized by Poly (ADP-ribose) polymerase (PARP) 1, which ADP-ribosylates itself and other factors at the site of breaks, and recruits the scaffold protein XRCC1, involved in both BER and SSBs direct repair (38). The BER-induced SSBs can pose risks to transcription and replication; if left unrepaired, they may collide with replication forks, leading to DSBs, which are typically repaired by the HR pathway (37).

In response to DSBs, a signaling cascade involves kinases such as ATM (ataxia telangiectasia mutated), ATR (ataxia telangiectasia and Rad3-related), mutated in A-T, and DNA-PKcs (DNA-dependent protein kinase catalytic subunit). These kinases phosphorylate multiple substrates, promoting the recruitment of DDR factors, cell cycle arrest, and repair processes. The choice between HR and NHEJ is tightly regulated throughout the cell cycle (219). HR is mainly active during late S and G2 phases, when sister chromatids or homologous sequences are available as templates for recombination. In contrast, NHEJ operates throughout the cell cycle, especially in G1, but is less accurate and more error-prone because it lacks a homologous template and may involve degradation of DNA ends (18,220), which makes the choice of the repair mechanism significantly impacting on genome stability.

The key NHEJ factor, 53BP1 (p53 binding protein 1), was first identified as a DNA damage checkpoint protein. Alongside BRCA1 (Breast cancer gene 1), a RING E3 and a well-known tumor suppressor involved in HR and often mutated in cancer (50,221), 53BP1 plays a central role in the choice of the repair pathway following DSBs (221).

In HR, BRCA1 acts upstream of BRCA2 (Breast Cancer gene 2), facilitating the resection of DSB ends. BRCA2 then loads the recombinase RAD51 onto resected single-stranded DNA (ssDNA), initiating strand invasion for homology search and recombination (222). The process of DNA end resection begins with CtBP (C terminus-binding protein)-interacting protein (CtIP) working together with the MRN complex (MRE11, Meiotic recombination 11, RAD50, and NBS1, Nijmegen breakage syndrome protein 1) to generate short ssDNA at the break ends (223). This resection produces 3' overhangs, allowing RAD51 to form nucleofilaments that displace RPA and invade the homologous template, forming D-loop structures for homology search (222). Once a polymerase has extended the 3' end of the invading strand, the ligation by DNA Ligase I yields to a heteroduplexed DNA structure. Finally, the recombination intermediates like dHJs are resolved and the correction of the DSB is complete (31,220).

While in S phase, BRCA1 promotes the removal of 53BP1 to enable resection, in the NHEJ pathway, which does not require sequence homology, 53BP1 negatively regulates end resection during G1 (91,92,220). Differently from HR, NHEJ acts in an error-prone manner. Ku heterodimer (Ku70-Ku80) initiates NHEJ by binding to the free DNA ends of the DSB and recruiting DNA-PKcs, with which it forms DNA-PK complex(224). DNA-PKcs is recruited with the endonuclease Artemis, undergoes autophosphorylation and activates the nuclease (225). Activated by DNA binding, DNA-PKcs additionally phosphorylates Ku proteins, Ligase IV and its cofactor XRCC4 (X- ray repair cross-complementing protein 4) (224,225). The final ligation of DNA ends by Ligase IV completes the C (canonical)-NHEJ mechanism (224,226).

Recent evidence indicates the Alt (alternative)-NHEJ, also named microhomology-mediated end joining (MMEJ), as an error-prone back-up pathway when C (canonical)-NHEJ or HR fail to repair (227). This pathway requires a 2 to 20 nucleotides homology sequence to start (228). PARP1 recognizes DNA breaks and activates MRN/CtIP for DNA end resection, which causes the exposition of microhomology sequence (229). DNA ends are aligned via the microhomology sequences, and non-homologous 3' tails are degraded by XPF. PolQ fills the gaps (230) prior to ligation by Ligase III/XRCC1 (X- ray repair cross-complementing protein 1) or Ligase I (231).

Alt-NHEJ relates to the other DSBs repair pathways in a mutual regulation and competition manner: PARP1 and Ku proteins compete for DNA breaks sensing (229,232); PolQ interacts with RAD51, removing RPA from resected DSBs (230); ligation by Alt-NHEJ dependent Ligase III opposes to C-NHEJ Ligase IV activity (231). On the other hand, functional HR or C-NHEJ suppresses Alt-NHEJ (233). These findings indicate that multiple mechanisms cooperate to safeguard genomic stability by counteracting the harmful effects of aberrant Alt-NHEJ activation (227,228). As well as for other DDR mechanisms, the proper coordination of DSB repair pathway choice is essential for maintaining genome stability (234).

Additionally, the removal of interstrand crosslinks (ICLs), highly cytotoxic covalent links between the double helix, is triggered by Fanconi anemia (FA) pathway in cooperation with NER and HR factors (31). The FA protein complex, formed by FANCM and FAAP24, recruits a large multi-subunit ubiquitin ligase, termed the core complex, to DNA lesions. This structure is composed of FANCA-FANCG, FANCC-FANCE-FANCF and FANCB-FANCL-FAAP100 (235). The core complex mono-ubiquitinates FANCD2 and FANCI, which then localize to DNA repair foci together with FANCD1, FANCI, and FANCN (235,236). ICLs final repair occurs through the collaborative activity of FA, NER and HR factors (31).

Figure Legends

Figure 1. Structural domain organization of the HECT E3 family members. Structural organization of the N-lobe, the C-lobe and the hinge region (indicated with arrows) of the HECT E3s (A) ITCH, (B) NEDD4, (C) WWP1, (D) WWP2, and (E) HUWE1. The N-lobe is responsible for interacting with the E2-Ub complex, while the catalytic cysteine (in red), which forms the thioester intermediate, is located within the C-lobe. The hinge region provides the flexibility for conformational changes necessary for the catalytic function.

Figure 2. Classification of the HECT E3 ubiquitin ligase family. All members of the HECT E3 family are defined by the presence of a C-terminal HECT domain, which acts as the catalytic domain. The HECT E3 family is composed of three subfamilies. (A) The members of the NEDD4 subfamily are characterized by an N-terminal C2 domain and several WW domains. (B) The HERC subfamily includes members characterized by one or more RLD domains. (C) The “Other” HECT E3 subfamily is identified by the lack of both RLD and WW domains.


Figure 3. Roles of the NEDD4-like HECT E3 subfamily in the regulation of DDR. (A) NEDD4 modulates p53 activity through the ubiquitination of MDM2 and regulates HR repair by targeting PTEN. (B) NEDD4L affects the response to genotoxic stress by targeting Smad2/3 and promotes apoptosis. It regulates the balance between two DNA repair pathways (HR and NHEJ) by targeting 53BP1. NEDD4L reduces BER efficacy by targeting OGG1. (C) ITCH controls proliferation and apoptosis by regulating p63, p73 and c-FLIP levels. (D) WWP1 modulates cell proliferation, oxidative stress response and SSBs or DSBs repair by targeting p27^{Kip1}, ΔNp63, TXNIP and JARID1B, while suppressing p53 activity. (E) WWP2 impairs the SSBs repair through the ubiquitination of PARP1 and the transcriptional arrest at the damage site by targeting RNAPII. (F) SMURF1 inhibits apoptosis by targeting RhoB. (G) SMURF2 modulates chromatin architecture and controls the recruitment of DNA repair factors by ubiquitinating RNF20 and RNF168. SMURF1/2 promotes p53 degradation by favoring MDM2 stabilization. (H) NEDL1 activates p53 signaling, promoting apoptosis. (I) NEDL2 regulates the DDR via modulation of p73 activity. Arrows indicate activation or promotion of the pathway, while blunt lines denote inhibition.


Figure 4. Involvement of the HERC family members in cellular processes linked to genomic stability. (A) HERC1 is shown to inhibit BAK, affecting UV damage-induced apoptosis. HERC1 also interacts with MSH2, a protein involved in MMR. (B) HERC2 stabilizes p53 through oligomerization. On the other hand, it marks p53 for degradation by MDM2 at late stages of DDR. HERC2 additionally targets both RPA and BRCA1 for degradation. (C) HERC3 is associated with frameshift mutations, increased mutation rate, leading to gastric and colorectal carcinomas. (D) HERC4 is linked to SAV1, a core component of the Hippo pathway, which regulates cell proliferation and growth signaling. Dysregulation of HERC4 is associated with breast and lung cancer. (E, F) HERC5 and HERC6 are involved in innate immunity, affecting signaling cascades, interferon production, and cytokine release. They might have then a still unknown role in damage response to infections affecting genome integrity. Arrows indicate activation or promotion of the pathway, while blunt lines denote inhibition.


Figure 5. Roles of the “Other” subfamily in DDR and repair. (A) AREL1 promotes the ubiquitination and subsequent proteasomal degradation of pro-apoptotic factors, including IAP, SMAC, HtrA2, and ARTS. (B) HUWE1 targets and destabilizes key oncoproteins such as Dishevelled and C-Myc. It concurrently confers apoptotic sensitivity through its ubiquitin ligase activity on Mcl-1, HDAC2, and Cdc6. Its role in DNA integrity maintenance is highlighted by its interaction with H2AX and BRCA1 in HR repair, and its regulatory effect on NEIL1 in BER. (C)


G2E3 is necessary for an efficient DDR signaling and regulation of mitosis. (D) TRIP12 significantly influences SSBs repair by modulating the stability or activity of PARP1, while its regulatory impact on ARF underscores its contribution to appropriate cell cycle transitions. Acting on RNF168, TRIP12 also regulates the recruitment of DSBs repair factors. (E) Additionally, to TRIP12, UBR5 regulates RNF168. UBR5 is involved in DDR, facilitating DNA repair upon replication stress or oxidative stress. It interacts with ATMIN and influences 53BP1 localization for initiating NHEJ. Finally, UBR5 regulates the oxidative stress response through its effect on TXNIP. Arrows indicate activation or promotion of the pathway, while blunt lines denote inhibition.


Box 1 Figure. Different types of ubiquitination and the specific roles of ubiquitin chain linkages. (A) The key enzymes involved in the ubiquitination cycle: E1, E2 and E3. The process begins with the activation of ubiquitin and its covalent attachment to the E1. In the second step, ubiquitin is transferred to the E2 during conjugation. Finally, the poly-ubiquitin chain is transferred to the target protein through the action of the E3. (B) Formation of different types of ubiquitin chains, including mono-ubiquitination, poly-ubiquitination via the same or different ubiquitin acceptor sites and the formation of branched poly-ubiquitin chains. (C) The different lysine residues involved in the formation of ubiquitin chains determine the fate of poly-ubiquitinated proteins, including their role in DDR.













A



B



Mono-ubiquitination

Multiple mono-ubiquitination

Homotypic poly-ubiquitination

Heterotypic poly-ubiquitination

Heterotypic branched poly-ubiquitination

C

K6

- DNA damage ⚡
- Mitophagy

K11

- Degradation
- Cell cycle control
- Membrane trafficking

K27

- DNA damage ⚡
- Mitochondrial damage response

K29

- Anti-bacterial autophagy

K33

- Autophagy

K48

- Degradation

K63

- DNA damage ⚡
- Anti-bacterial autophagy