

EXTERIOR SPHERE CONDITION AND TIME OPTIMAL CONTROL FOR DIFFERENTIAL INCLUSIONS*

PIERMARCO CANNARSA[†] AND KHAI T. NGUYEN[‡]

Abstract. The minimum time function $T(\cdot)$ of smooth control systems is known to be locally semiconcave provided Petrov’s controllability condition is satisfied. Moreover, such a regularity holds up to the boundary of the target under an inner ball assumption. We generalize this analysis to differential inclusions, replacing the above hypotheses with the continuity of $T(\cdot)$ near the target, and an inner ball property for the multifunction associated with the dynamics. In such a weakened setup, we prove that the hypograph of $T(\cdot)$ satisfies, locally, an exterior sphere condition. As is well known, this geometric property ensures most of the regularity results that hold for semiconcave functions, without assuming $T(\cdot)$ to be Lipschitz.

Key words. proximal normal vectors, exterior sphere condition, differential inclusions, time optimal control, semiconcave functions

AMS subject classifications. Primary, 34A60, 49J53; Secondary, 49J15

DOI. 10.1137/110825078

1. Introduction. This paper is concerned with the time optimal control problem for the differential inclusion

$$(1.1) \quad \begin{cases} \dot{x}(t) \in F(x(t)) & \text{a.e. } t \geq 0, \\ x(0) = x_0 \in \mathbb{R}^n, \end{cases}$$

with a closed target $\mathcal{S} \subset \mathbb{R}^n$. Here, $F : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ is a convex-valued Lipschitz continuous multifunction describing the dynamics, and it will be subject to further conditions in Hamiltonian form (see section 3 below). For each trajectory $y^{x_0}(\cdot)$ of (1.1), we denote by $\theta(y^{x_0}(\cdot)) := \inf \{t \geq 0 \mid y^{x_0}(t) \in \mathcal{S}\}$ the transition time from x_0 to \mathcal{S} along $y^{x_0}(\cdot)$. Clearly, $\theta(y^{x_0}(\cdot)) \in [0, \infty]$. The *minimum time* $T(x_0)$ to reach \mathcal{S} from x_0 is defined by

$$(1.2) \quad T(x_0) := \inf \{\theta(y^{x_0}(\cdot)) \mid y^{x_0}(\cdot) \text{ is a trajectory of (1.1)}\}.$$

Observe that, in general, $T(\cdot) : \mathbb{R}^n \rightarrow [0, \infty]$. The controllable set \mathcal{C} consists of all points $x \in \mathbb{R}^n$ such that $T(x)$ is finite.

The regularity of the minimum time function, which is related to the controllability of (1.1), has been the subject of an extensive literature. Most papers study the case where F is given with a parameterization, which means that F has the form

$$(1.3) \quad F(x) = \{f(x, u) \mid u \in U\} \quad \forall x \in \mathbb{R}^n$$

with $U \subseteq \mathbb{R}^m$ compact and $f : \mathbb{R}^n \times U \rightarrow \mathbb{R}^n$ satisfying the following:

- f is continuous in (x, u) ; and

*Received by the editors February 18, 2011; accepted for publication (in revised form) October 5, 2011; published electronically December 13, 2011. This research was partially supported by the GDRE ConEDP issued by CNRS, INdAM, and the Université de Provence.

<http://www.siam.org/journals/sicon/49-6/82507.html>

[†]Dipartimento di Matematica, Università di Roma “Tor Vergata,” 00133 Rome, Italy (cannarsa@mat.uniroma2.it).

[‡]Dipartimento di Matematica Pura ed Applicata, Università di Padova, 35121 Padova, Italy (khai@math.unipd.it).

- there exists $k > 0$ so that

$$|f(x_0, u) - f(x_1, u)| \leq k|x_0 - x_1|$$

for all $u \in U$ and $x_0, x_1 \in \mathbb{R}^n$.

On the contrary, fewer results are available for systems modeled by differential inclusions such as (1.1)—an exception is [15], where a representation formula for the proximal subgradient of $T(\cdot)$ is recovered. Although Lipschitz multifunctions with convex values always admit parameterizations by Lipschitz functions (see [1, 14]), it is a challenging open problem to determine which multifunctions F admit parameterizations with *smooth* functions. Observe that a certain smoothness of the parameterization—essentially that $f(\cdot, u)$ is differentiable with $D_x f(\cdot, u)$ Lipschitz—is crucial in order to derive further regularity properties of the value function, such as *semiconcavity*, by known methods (see [3, 6]). This fact explains why parameterization theorems have so far proved of little use for regularity purposes.

Instead of searching for smooth parameterizations, in [8] the first author and Wolenski proposed an alternative strategy to obtain semiconcavity results for the value function of the *Mayer problem* for system (1.1). Unlike previous approaches, the proof of [8] exploits the semiconvexity in x of the Hamiltonian

$$(1.4) \quad H(x, p) = \sup_{v \in F(x)} \langle v, p \rangle, \quad (x, p) \in \mathbb{R}^n \times \mathbb{R}^n,$$

as well as the nonsmooth maximum principle.

As for time optimal control problems, when F is parameterized as in (1.3) and $D_x f(\cdot, u)$ is Lipschitz, the minimum time function is known to be semiconcave on $\mathcal{C} \setminus \text{int}(\mathcal{S})$ provided \mathcal{S} has the inner ball property, and $T(\cdot)$ is dominated by the distance from \mathcal{S} (see [7, 6]). The latter assumption is equivalent to *Petrov’s controllability condition*: for some constant $\mu > 0$ and all points $x \in \partial\mathcal{S}$,

$$(1.5) \quad \min_{u \in U} \langle f(x, u), \nu \rangle \leq -\mu|\nu|$$

for all proximal normal vectors ν to \mathcal{S} at x . This is a strong hypothesis on the control system since, when $f(x, u) = u$, it is equivalent to the fact that U contains an open neighborhood of the origin. Nevertheless, it is also necessary for the semiconcavity of $T(\cdot)$ up to the boundary of the target, because it is equivalent to the Lipschitz continuity of $T(\cdot)$ in a neighborhood of \mathcal{S} —a direct consequence of semiconcavity. On the other hand, the assumption that \mathcal{S} satisfies a uniform interior sphere condition¹ can be removed by the method introduced in [4], where the local semiconcavity of $T(\cdot)$ in $\mathcal{C} \setminus \mathcal{S}$ is derived from an analogous geometric property of $f(x, U)$. Moreover, the approach of [8] to obtain the semiconcavity of the value function can be adapted to the minimum time function for (1.1), keeping Petrov’s condition and the inner ball property of \mathcal{S} as standing assumptions (see [5]).

The main purpose of this paper is to study the regularity of $T(\cdot)$ for the general system (1.1), assuming neither the inner ball property of \mathcal{S} nor Petrov’s condition (1.5). In view of the above discussion, the expected regularity of $T(\cdot)$ will be weaker than semiconcavity, since the Lipschitz continuity of $T(\cdot)$ near \mathcal{S} is no longer guaranteed. Similarly, the lack of the inner ball property of \mathcal{S} will result in the fact that the regularity of $T(\cdot)$ will be just local in $\mathcal{C} \setminus \mathcal{S}$.

¹In this paper, the expressions “ \mathcal{S} has the inner ball property” and “ \mathcal{S} satisfies a uniform interior sphere condition” have the same meaning.

A weaker class of functions enjoying most of the differential properties of semiconcave functions is the class of all continuous functions u whose hypograph is φ -convex; see, e.g., [2, 10]. For any function u of the above class, Colombo and Marigonda [10] proved the existence of second order Taylor's expansions a.e., showed the local BV regularity of the gradient, and studied the structure of the corresponding singular set. For parameterized control systems and targets satisfying an internal sphere condition, Colombo and the second author [11] obtained φ -convexity results for the hypograph of $T(\cdot)$, assuming that $T(\cdot)$ is continuous and the proximal normal cone to the hypograph of $T(\cdot)$ is pointed at every point. Furthermore, the second author removed the pointedness assumption in [12], proving that, for any function u in the above class, the points where the proximal normal cone to the hypograph of u is not pointed form a closed set with zero Lebesgue measure. Consequently, in the above φ -convex setup, $T(\cdot)$ retains most of the regularity properties of semiconcave functions.

Therefore, the main results of our paper can also be described as the extension of the analysis of [11, 12] to systems given in the general form (1.1), replacing the geometric assumption on \mathcal{S} with a different property of the dynamics. This goal will be achieved in two steps. First, we will show that if $T(\cdot)$ is continuous, then the hypograph of $T(\cdot)$ satisfies an exterior sphere condition provided \mathcal{S} has the inner ball property (Theorem 4.1). Then, we shall remove the assumption that \mathcal{S} has such a property adapting the approach of [4] to system (1.1), that is, showing that the sublevel sets $\{T(\cdot) \leq t\}$ acquire the inner ball property for $t > 0$ small enough provided $F(\cdot)$ satisfies the same geometric property (see Theorem 4.7 and Corollary 4.8 below).

The outline of this paper is the following. In section 2, we introduce our notation and recall some results from nonsmooth analysis. Our standing assumptions on F and H are described in section 3, together with basic properties of the Hamiltonian and estimates for solutions to certain differential inclusions. Finally, section 4 contains our main results and their proofs.

2. Notation and preliminary results. Let $Q \subset \mathbb{R}^n$ be a closed set and let $x \in Q$. We say that a vector $v \in \mathbb{R}^n$ is a *proximal normal* to Q at x (and write $v \in N_Q^P(x)$) if there exists $\sigma \geq 0$ such that

$$(2.1) \quad \langle v, y - x \rangle \leq \sigma |y - x|^2 \quad \forall y \in Q.$$

Equivalently, $v \in N_Q^P(x)$ if and only if there exists $\lambda > 0$ such that $d_Q(x + \lambda v) = \lambda |v|$, where $d_Q(\cdot)$ denotes the distance function from Q . We say that $v \in N_Q^P(x)$ is *realized* by a ball of radius $\rho > 0$ if (2.1) holds true for $\sigma = \frac{|v|}{2\rho}$.

The Clarke normal cone to Q at x , $N_Q^C(x)$, is defined as

$$N_Q^C(x) = \overline{\text{co}}\{\xi \in \mathbb{R}^n \mid \exists x_i \rightarrow x, \xi_i \rightarrow \xi, \xi_i \in N_Q^P(x_i)\},$$

where $\overline{\text{co}}$ denotes the closed convex hull.

Now, suppose $f : \mathbb{R}^n \rightarrow (-\infty, +\infty]$ is lower semicontinuous and denote by

$$\text{epi}(f) = \{(x, \beta) \mid \beta \geq f(x)\}$$

its (closed) epigraph. For any $x \in \text{dom}(f) := \{x \mid f(x) < \infty\}$, the set of proximal subgradients of f at x , $\partial_P f(x)$, equals

$$\partial_P f(x) = \{\xi \in \mathbb{R}^n \mid (\xi, -1) \in N_{\text{epi}(f)}^P(x, f(x))\}.$$

Similarly, the set of Clarke subgradients of f at x , $\partial f(x)$, is given by

$$\partial f(x) = \{ \xi \in \mathbb{R}^n \mid (\xi, -1) \in N_{\text{epi}(f)}^C(x, f(x)) \}.$$

The property of *semiconcavity* has several characterizations (see, e.g., [6]), and we now proceed to recall some of them. For a convex set $K \subseteq \mathbb{R}^n$, a function $f : K \rightarrow \mathbb{R}$, and a constant $c \geq 0$, the following properties are equivalent:

- (1) for all $x_0, x_1 \in K$ and $0 \leq \lambda \leq 1$, one has

$$(1 - \lambda)f(x_0) + \lambda f(x_1) - f(x_\lambda) \leq c\lambda(1 - \lambda)|x_1 - x_0|^2,$$

where $x_\lambda = (1 - \lambda)x_0 + \lambda x_1$;

- (2) f is continuous and, for all $x \in K$ and $z \in \mathbb{R}^n$ with $x \pm z \in K$, one has

$$f(x + z) + f(x - z) - 2f(x) \leq 2c|z|^2;$$

- (3) the map $x \mapsto f(x) - c|x|^2$ is concave.

In any of the above cases, we say that f is (linearly) semiconcave on K with constant c , and we call f semiconvex on K if $-f$ is semiconcave.

The following lemma is proved in [8].

LEMMA 2.1. *Suppose that $f : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}$ is Lipschitz in (x, y) on a boxed neighborhood $U_x \times U_y = \{(x, y) \mid \max\{|x - \bar{x}|, |y - \bar{y}|\} < \delta\}$ of the point $(\bar{x}, \bar{y}) \in \mathbb{R}^n \times \mathbb{R}^m$ and that, for each $y \in U_y$, the function $x \rightarrow f(x, y)$ is semiconvex on U_x with constant independent of y . Then, for any $\xi = (\xi_x, \xi_y) \in \partial f(\bar{x}, \bar{y})$, one has $\xi_x \in \partial_x f(\bar{x}, \bar{y})$.*

We now introduce a concept which plays a major role in what follows.

DEFINITION 2.2. *Let $Q \subset \mathbb{R}^n$ be closed and let $\theta(\cdot) : \partial Q \rightarrow (0, \infty)$ be continuous. We say that Q satisfies the $\theta(\cdot)$ -exterior sphere condition if and only if, for every $x \in \partial Q$, there exists a nonzero vector $v_x \in N_Q^P(x)$ which is realized by a ball of radius $\theta(x)$.*

Denote by Q' the closure of the complement of Q . We say that the set Q satisfies the $\theta(\cdot)$ -interior sphere condition if Q' satisfies the $\theta(\cdot)$ -exterior sphere condition. If Q satisfies the $\theta(\cdot)$ -interior sphere condition for some constant function $\theta(\cdot) = \theta_0$, then we also say that Q has the *inner ball property* (of radius θ_0).

Finally, for any function $f : \Omega \rightarrow \mathbb{R}$ denote by

$$\text{hypo}(f) = \{(x, \beta) \mid \beta \leq f(x)\}$$

the hypograph of f . Let $\theta : \Omega \rightarrow (0, \infty)$ be a continuous function. Adapting Definition 2.2, we say $\text{hypo}(f)$ satisfies the $\theta(\cdot)$ -exterior sphere condition if for every $x \in \Omega$ there exists a nonzero vector $v_x \in N_{\text{hypo}(f)}^P(x, f(x))$ which is realized by a ball in \mathbb{R}^{n+1} of radius $\theta(x)$. We now recall a result from [12].

PROPOSITION 2.3. *Let $\Omega \subset \mathbb{R}^N$ be an open set and let $f : \Omega \rightarrow \mathbb{R}$ be continuous. Assume that $\text{hypo}(f)$ satisfies the $\theta(\cdot)$ -exterior sphere condition, where $\theta : \Omega \rightarrow (0, \infty)$ is a given continuous function. Then there exists a sequence of sets $\Omega_h \subseteq \Omega$, with Ω_h compact in $\text{dom}(f)$, such that the following hold:*

- (1) the union of Ω_h covers \mathcal{L}^N -almost all of $\text{dom}(f)$;
- (2) for all $x \in \cup_h \Omega_h$ there exist $\delta = \delta(x) > 0$ and $L = L(x) > 0$ such that f is Lipschitz on $B(x, \delta)$ with constant L and hence semiconcave on $B(x, \delta)$.

Consequently, the following holds:

- (3) f is a.e. Fréchet differentiable and admits a second order Taylor expansion around almost every point of its domain.

3. Hypotheses and some consequences. We list below our hypotheses on F and H —the problem data introduced in (1.1) and (1.4), respectively—together with some of their consequences.

Hypotheses (F).

- (F1) $F(x)$ is nonempty, convex, and compact for each $x \in \mathbb{R}^n$.
- (F2) F is Lipschitz continuous with respect to the Hausdorff distance. Thus, if K is the Lipschitz constant of F , then $K|p|$ is the Lipschitz constant of $H(\cdot, p)$, i.e.,

$$(3.1) \quad |H(y, p) - H(x, p)| \leq K|p||y - x| \quad \forall x, y \in \mathbb{R}^n, \forall p \in \mathbb{R}^n.$$

Hypotheses (H).

- (H1) There exists a constant $c_0 \geq 0$ such that $x \mapsto H(x, p)$ is semiconvex with semiconvexity constant $c_0|p|$.
- (H2) For all $p \neq 0$, the gradient $\nabla_p H(\cdot, p)$ exists and is globally Lipschitz, i.e.,

$$(3.2) \quad |\nabla_p H(x, p) - \nabla_p H(y, p)| \leq K_1|y - x| \quad \forall x, y \in \mathbb{R}^n, \forall p \in \mathbb{R}^n \setminus \{0\},$$

for some constant $K_1 \geq 0$.

Remark 3.1. In particular, (F2) implies the following:

- (F3) There exists $K_2 > 0$ such that $\max\{|v| \mid v \in F(x)\} \leq K_2(1 + |x|)$.

This in turn guarantees that solutions to (1.1) are defined on $[0, \infty)$.

Global Lipschitz continuity in both (F2) and (H2) was assumed just to simplify computations. Indeed, our results still hold if F is locally Lipschitz with respect to the Hausdorff distance, and $\nabla_p H(\cdot, p)$ is locally Lipschitz in x , uniformly so over p in $\mathbb{R}^n \setminus \{0\}$. In that case, however, (F3) has to be assumed as an extra condition.

From Lemma 2.1 we immediately obtain the following.

COROLLARY 3.2. *Suppose H satisfies assumption (H1). Then*

$$\partial H(x, p) \subseteq \partial_x H(x, p) \times \partial_p H(x, p) \quad \forall p \neq 0.$$

The following proposition is a consequence of [8, Proposition 1].

PROPOSITION 3.3. *Suppose F satisfies (F) and (H1). Then the following hold:*

- (1) *for each $x, z \in \mathbb{R}^n$, we have*

$$H(x + z, p) + H(x - z, p) - 2H(x, p) \geq -c_0|p||z|^2;$$

- (2) *for each $x, y \in \mathbb{R}^n$, and $\xi \in \partial_x H(x, p)$, we have*

$$H(y, p) - H(x, p) - \langle \xi, y - x \rangle \geq -c_0|p||y - x|^2.$$

The differentiability statement in assumption (H2) is equivalent to the argmax set of $v \mapsto \langle v, p \rangle$ being a singleton, which equals $\nabla_p H(x, p)$ and will also be denoted by $F_p(x)$. In view of (H2), for all $p \neq 0$ the function $F_p(\cdot)$ is globally Lipschitz with constant K_1 . The main use of (H2) is given by the following result, whose proof is straightforward.

PROPOSITION 3.4. *Assume (F) and (H), and let $p(\cdot)$ be an absolutely continuous arc on $[0, T]$, with $p(t) \neq 0$ for all $t \in [0, T]$. Then, for each $x \in \mathbb{R}^n$, the problem*

$$(3.3) \quad \begin{cases} \dot{x}(t) = F_{p(t)}(x(t)) & \text{a.e. } t \in [0, T], \\ x(0) = x \end{cases}$$

has a unique solution $y(\cdot, x)$. Moreover, $x \mapsto y(t, x)$ is Lipschitz on \mathbb{R}^n and

$$(3.4) \quad |y(t, z) - y(t, x)| \leq e^{K_1 t} |z - x| \quad \forall x, z \in \mathbb{R}^n, \forall t \in [0, T].$$

We conclude this section with some simple consequences of Gronwall's lemma.

LEMMA 3.5. *Let $G : [0, T] \times \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ be an upper semicontinuous multifunction. Assume $G(t, \cdot)$ satisfies hypotheses (F1), (F2) uniformly in $t \in [0, T]$ and is such that, for some $K_0 > 0$,*

$$|v| \leq K_0 |p| \quad \forall v \in G(t, p), \forall (t, p) \in [0, T] \times \mathbb{R}^n.$$

Let $p(\cdot)$ be a solution of the differential inclusion

$$(3.5) \quad \begin{cases} \dot{p}(t) \in G(t, p(t)) & \text{a.e. } t \in [0, T], \\ p(0) = p_0. \end{cases}$$

Then

$$e^{-K_0 t} |p(0)| \leq |p(t)| \leq e^{K_0 t} |p(0)| \quad \forall t \in [0, T].$$

Moreover, for all $0 \leq t_1 \leq t_2 \leq T$,

$$e^{-K_0(t_2-t_1)} |p(t_2)| \leq |p(t_1)| \leq e^{K_0(t_2-t_1)} |p(t_2)|$$

and

$$|p(t_2) - p(t_1)| \leq K_0 e^{K_0(t_2-t_1)} (t_2 - t_1) |p(t_2)|.$$

Proof. Since $p(t) = p(0) + \int_0^t \dot{p}(s) ds$, we have

$$|p(t)| \leq |p(0)| + \int_0^t |\dot{p}(s)| ds \leq |p(0)| + K_0 \int_0^t |p(s)| ds.$$

So, using Gronwall's inequality, we get $|p(t)| \leq e^{K_0 t} |p(0)|$.

We are now going to prove that $e^{-K_0 t} |p(0)| \leq |p(t)|$ for all $t > 0$. Fixing $t > 0$, we define $g(s) := p(t - s)$ for all $s \in [0, t]$. Since $\dot{g}(s) = -\dot{p}(t - s)$ for almost all $s \in [0, t]$, we have $g(s) = g(0) + \int_0^s \dot{g}(\tau) d\tau$ for all $s \in [0, t]$. Thus,

$$\begin{aligned} |g(s)| &\leq |g(0)| + \int_0^s |\dot{g}(\tau)| d\tau = |g(0)| + \int_0^s |\dot{p}(t - \tau)| d\tau \\ &\leq |g(0)| + K_0 \int_0^s |p(t - \tau)| d\tau = |g(0)| + K_0 \int_0^s |g(\tau)| d\tau. \end{aligned}$$

Again by Gronwall's inequality, we obtain $|g(s)| \leq e^{K_0 s} |g(0)|$ for all $s \in [0, t]$. In particular, $|g(t)| \leq e^{K_0 t} |g(0)|$. The proof is completed noting that $g(t) = p(0)$ and $g(0) = p(t)$. \square

COROLLARY 3.6. *Let $p(\cdot)$ be a solution of (3.5). Then either $p(t) = 0$ for all $t \in [0, T]$ or $p(t) \neq 0$ for all $t \in [0, T]$.*

LEMMA 3.7. *Let $y(\cdot, x_0)$ be a solution of (1.1). Then, for all $t > 0$, the following hold:*

- (i) $|y(t, x_0)| \leq (|x_0| + 1)e^{K_2 t} - 1$;
- (ii) $|y(t, x_0) - x_0| \leq (|x_0| + 1)(e^{K_2 t} - 1) \leq K_2(|x_0| + 1)e^{K_2 t} t$.

Proof. Since

$$y(t, x_0) = x_0 + \int_0^t \dot{y}(s, x_0) ds,$$

recalling (F3) we have

$$|y(t, x_0)| \leq |x_0| + K_2 t + K_2 \int_0^t |y(s, x_0)| ds.$$

Hence, Gronwall’s inequality yields (i). Then, observing that

$$|y(t, x_0) - x_0| \leq K_2 \int_0^t (1 + |y(s, x_0)|) ds,$$

(ii) follows using (i) in the above estimate. \square

4. Main results.

4.1. Part I. In this part, we will assume that \mathcal{S} is nonempty and closed and has the inner ball property with balls of radius $\rho_0 > 0$. Moreover, assumptions (F) and (H) are also assumed throughout. Recall that c_0, K, K_1, K_2 are the constants in (H1), (F2), (H2), (F3). Let us define, for any $r > 0$,

$$\mathcal{S}'(r) = \{x \mid T(x) \geq r\}, \quad \mathcal{S}' = (\mathbb{R}^n \setminus \mathcal{S}) \cup \partial \mathcal{S}, \quad \mathcal{C} = \{x \in \mathbb{R}^n \setminus \mathcal{S} \mid T(x) < +\infty\}$$

and let $T|_{\mathcal{O}} : \mathcal{O} \rightarrow \mathbb{R}$ be the restriction of T to \mathcal{O} ; i.e., $T|_{\mathcal{O}}(x) = T(x)$ for $x \in \mathcal{O}$.

Our main results are the following theorem, together with Corollary 4.3.

THEOREM 4.1. *Assume (F) and (H). Suppose further that \mathcal{S} is nonempty and closed and has the inner ball property with balls of radius $\rho_0 > 0$ and that $T(\cdot)$ is continuous in an open subset \mathcal{O} of \mathcal{C} . Then, the hypograph of $T|_{\mathcal{O}}(\cdot)$ satisfies a $\rho_T(\cdot)$ -exterior sphere condition for some continuous function $\rho_T(\cdot) : \mathcal{O} \rightarrow (0, \infty)$.*

Remark 4.2. The function $\rho_T(\cdot)$ can be explicitly computed and depends only on $x, T(x)$, and on c_0, K, K_1, K_2, ρ_0 .

Consequently, under the assumptions of Theorem 4.1, $T|_{\mathcal{O}}(\cdot)$ enjoys the regularity properties described in Proposition 2.3. Moreover, the following corollary follows from Theorem 4.1 and [13, Theorem 21].

COROLLARY 4.3. *Under the assumptions of Theorem 4.1, if $T(\cdot)$ is locally Lipschitz in \mathcal{O} , then $T(\cdot)$ is locally semiconcave in \mathcal{O} .*

The main part of the proof of Theorem 4.1 is divided into three lemmas.

LEMMA 4.4. *Suppose $\bar{x} \in \mathcal{O}$ is not a local maximum of $T(\cdot)$. Let $r = T(\bar{x})$ and let $\bar{x}^+(\cdot)$ be an optimal trajectory steering \bar{x} to \mathcal{S} in time r , and set $\bar{x}^-(s) = \bar{x}^+(r - s)$. Then, there exists an arc $\bar{p}(\cdot)$ defined on $[0, r]$, with $\bar{p}(s) \neq 0$ for all $s \in [0, r]$, such that*

$$(4.1) \quad \begin{cases} -\dot{\bar{p}}(s) \in \partial_x H(\bar{x}^-(s), -\bar{p}(s)) & \text{a.e. } s \in [0, r], \\ \dot{\bar{x}}^-(s) = -F_{-\bar{p}(s)}(\bar{x}^-(s)) & \text{a.e. } s \in [0, r]. \end{cases}$$

Moreover, $-\bar{p}(r - t) \in N_{\mathcal{S}'(r-t)}^P(\bar{x}^+(t))$ is realized by a ball of radius $\rho(r - t)$ for all $t \in [0, r]$, i.e.,

$$(4.2) \quad \left\langle \frac{-\bar{p}(r - t)}{|\bar{p}(r - t)|}, \bar{y} - \bar{x}^+(t) \right\rangle \leq \frac{1}{2\rho(r - t)} |\bar{y} - \bar{x}^+(t)|^2 \quad \forall \bar{y} \in \mathcal{S}'(r - t),$$

where

$$(4.3) \quad \rho(s) = \frac{\rho_0}{1 + 2c_0\rho_0s} e^{-(K+2K_1)s}.$$

Proof. Set $\bar{x}_1 = \bar{x}^+(r)$. Of course, $\bar{x}_1 \in \partial\mathcal{S}$. Since \mathcal{S} satisfies the ρ_0 -internal sphere condition, there exists a proximal normal vector $v \neq 0$ to \mathcal{S}' at \bar{x}_1 such that $\overline{B}(\bar{x}_1 + \rho_0 \frac{v}{|v|}, \rho_0) \subseteq \mathcal{S}$, i.e.,

$$(4.4) \quad \left\langle \frac{v}{|v|}, z - \bar{x}_1 \right\rangle \leq \frac{1}{2\rho_0} |z - \bar{x}_1|^2 \quad \forall z \in \mathcal{S}'.$$

Now, consider the reversed differential inclusion with initial data

$$(4.5) \quad \begin{cases} \dot{y}(s) \in -F(y(s)) & \text{a.e. } s \in [0, r], \\ y(0) \in \overline{B}(\bar{x}_1 + \rho_0 \frac{v}{|v|}, \rho_0) \subseteq \mathcal{S}. \end{cases}$$

The Hamiltonian associated with $-F$ is defined by

$$(4.6) \quad H^-(x, p) := \sup_{v \in -F(x)} \langle v, p \rangle = \sup_{w \in F(x)} \langle w, -p \rangle = H(x, -p).$$

Let us recall that the attainable set from $\overline{B}(\bar{x}_1 + \rho_0 \frac{v}{|v|}, \rho_0)$, denoted by $\mathcal{A}^-(r)$, is defined to be the set of all points $y(r)$ where $y(\cdot)$ is a trajectory satisfying (4.5). Since $\bar{x}^-(\cdot)$ is a solution of (4.5) with initial point $y(0) = \bar{x}_1$, and $T(\cdot)$ does not have a local maximum at the point \bar{x} , one has that $\bar{x}^-(r) = \bar{x}$ is on the boundary of $\mathcal{A}^-(r)$. Indeed, suppose \bar{x} is not on the boundary of $\mathcal{A}^-(r)$; then there exists $\epsilon > 0$ such that $B(\bar{x}, \epsilon) \subset \mathcal{A}^-(r)$. Thus, $T(y) \leq r = T(\bar{x})$ for all $y \in B(\bar{x}, \epsilon)$, and we get a contradiction since $T(\cdot)$ does not have a local maximum at \bar{x} . Now, since \bar{x} is on the boundary of $\mathcal{A}^-(r)$, by [9, Theorem 3.5.4], there is an arc $\bar{p}(\cdot)$, such that $\|\bar{p}(\cdot)\|_\infty > 0$, satisfying

$$(4.7) \quad (-\dot{\bar{p}}(s), \dot{\bar{x}}^-(s)) \in \partial H^-(\bar{x}^-(s), \bar{p}(s)) \quad \text{a.e. } s \in [0, r],$$

and

$$(4.8) \quad \bar{p}(0) \in N_{\overline{B}(\bar{x}_1 + \rho_0 \frac{v}{|v|}, \rho_0)}(\bar{x}_1).$$

From (4.7), (4.6), and Corollary 3.2, we have

$$(4.9) \quad -\dot{\bar{p}}(s) \in \partial_x H(\bar{x}^-(s), -\bar{p}(s)) \quad \text{a.e. } s \in [0, r].$$

Moreover, owing to (3.1), for all $v \in \partial_x H(x, p)$ we have $|v| \leq K|p|$. Therefore, applying Lemma 3.5 to $G(s, -\bar{p}(s)) = \partial_x H(\bar{x}^-(s), -\bar{p}(s))$, we get

$$(4.10) \quad e^{-Ks} |\bar{p}(0)| \leq |\bar{p}(s)| \leq e^{Ks} |\bar{p}(0)| \quad \forall s \in [0, r].$$

Since $\|\bar{p}(\cdot)\|_\infty > 0$, we have $\bar{p}(s) \neq 0$ for all $s \in [0, r]$. Therefore, from (4.7) and Corollary 3.2 we get

$$(4.11) \quad \dot{\bar{x}}^-(s) = -\partial_p H(\bar{x}^-(s), -\bar{p}(s)) = -F_{-\bar{p}(s)}(\bar{x}^-(s)) \quad \text{a.e. } s \in [0, r].$$

We are now going to prove (4.2). Fix $t \in [0, r]$ and let $\bar{y} \in \mathcal{S}'(r - t)$, i.e., $T(\bar{y}) \geq r - t$. Let $\bar{y}^+(\cdot)$ be the solution of the Cauchy problem

$$(4.12) \quad \begin{cases} \dot{\bar{y}}^+(s) = F_{-\bar{p}(r-t-s)}(\bar{y}^+(s)) & \text{a.e. } s \in [0, r - t], \\ \bar{y}^+(0) = \bar{y}. \end{cases}$$

Note that $\bar{y}_1 := \bar{y}^+(r - t) \in \mathcal{S}'$. Then, $\bar{y}^-(s) := \bar{y}^+(r - t - s)$ satisfies $\bar{y}^-(r - t) = \bar{y}$ and

$$(4.13) \quad \begin{cases} \dot{\bar{y}}^-(s) = -F_{-\bar{p}(s)}(\bar{y}^-(s)) & \text{a.e. } s \in [0, r - t], \\ \bar{y}^-(0) = \bar{y}_1. \end{cases}$$

From (4.11), (4.13), and (3.4), we have

$$(4.14) \quad |\bar{y}^-(s) - \bar{x}^-(s)| \leq e^{K_1(r-t)} |\bar{y}_1 - \bar{x}_1| \quad \forall s \in [0, r - t].$$

In order to prove (4.2), observe that

$$(4.15) \quad \begin{aligned} & \langle -\bar{p}(r - t), \bar{y}^-(r - t) - \bar{x}^-(r - t) \rangle \\ &= \langle -\bar{p}(0), \bar{y}^-(0) - \bar{x}^-(0) \rangle + \int_0^{r-t} \frac{d}{ds} \langle -\bar{p}(s), \bar{y}^-(s) - \bar{x}^-(s) \rangle ds. \end{aligned}$$

Moreover,

$$\begin{aligned} & \frac{d}{ds} \langle -\bar{p}(s), \bar{y}^-(s) - \bar{x}^-(s) \rangle \\ &= \langle -\dot{\bar{p}}(s), \bar{y}^-(s) - \bar{x}^-(s) \rangle + \langle -\bar{p}(s), \dot{\bar{y}}^-(s) - \dot{\bar{x}}^-(s) \rangle \\ &= \langle -\dot{\bar{p}}(s), \bar{y}^-(s) - \bar{x}^-(s) \rangle + \langle -\bar{p}(s), -F_{-\bar{p}(s)}(\bar{y}^-(s)) + F_{-\bar{p}(s)}(\bar{x}^-(s)) \rangle \\ &= \langle -\dot{\bar{p}}(s), \bar{y}^-(s) - \bar{x}^-(s) \rangle - H(\bar{y}^-(s), -\bar{p}(s)) + H(\bar{x}^-(s), -\bar{p}(s)). \end{aligned}$$

Recalling (4.9) and Proposition 3.3 it follows that

$$\langle -\dot{\bar{p}}(s), \bar{y}^-(s) - \bar{x}^-(s) \rangle - H(\bar{y}^-(s), -\bar{p}(s)) + H(\bar{x}^-(s), -\bar{p}(s)) \leq c_0 |\bar{p}(s)| |\bar{y}^-(s) - \bar{x}^-(s)|^2.$$

Therefore,

$$(4.16) \quad \frac{d}{ds} \langle -\bar{p}(s), \bar{y}^-(s) - \bar{x}^-(s) \rangle \leq c_0 |\bar{p}(s)| |\bar{y}^-(s) - \bar{x}^-(s)|^2.$$

Owing to (4.8) and the fact that $\bar{p}(0) \neq 0$, we have $-\frac{\bar{p}(0)}{|\bar{p}(0)|} = \frac{v}{|v|}$. Thus, by (4.4) and the fact that $\bar{y}_1 \in \mathcal{S}'$ we obtain

$$(4.17) \quad \left\langle -\frac{\bar{p}(0)}{|\bar{p}(0)|}, \bar{y}_1 - \bar{x}_1 \right\rangle \leq \frac{1}{2\rho_0} |\bar{y}_1 - \bar{x}_1|^2.$$

Combining (4.15), (4.16), (4.17) and noting that $\bar{x}^-(0) = \bar{x}_1$, $\bar{y}^-(0) = \bar{y}_1$, $\bar{x}^-(r - t) = \bar{x}^+(t)$, $\bar{y}^-(r - t) = \bar{y}$, we conclude that

$$\langle -\bar{p}(r - t), \bar{y} - \bar{x}^+(t) \rangle \leq \frac{|\bar{p}(0)|}{2\rho_0} |\bar{y}_1 - \bar{x}_1|^2 + c_0 \int_0^{r-t} |\bar{p}(s)| |\bar{y}^-(s) - \bar{x}^-(s)|^2 ds.$$

Thus, by (4.10) and (4.14),

$$\left\langle \frac{-\bar{p}(r - t)}{|\bar{p}(r - t)|}, \bar{y} - \bar{x}^+(t) \right\rangle \leq \left(\frac{1}{2\rho_0} + c_0(r - t) \right) e^{(K+2K_1)(r-t)} |\bar{y} - \bar{x}^+(t)|^2.$$

So, (4.2) follows (4.3), and the proof is complete. \square

LEMMA 4.5. *Suppose $\bar{x} \in \mathcal{O}$ is not a local maximum of $T(\cdot)$. Let $r = T(\bar{x})$ and let $\bar{x}^+(\cdot)$ be an optimal trajectory steering \bar{x} to \mathcal{S} in time r . If $\bar{p}(\cdot)$ is the arc in Lemma 4.4, then $H(\bar{x}, -\bar{p}(r)) \geq 0$.*

Proof. Fixing $t \in (0, r]$, by (4.2) and the fact that $\bar{x} \in \mathcal{S}'(r - t)$, we have

$$\langle -\bar{p}(r - t), \bar{x} - \bar{x}^+(t) \rangle \leq \frac{1}{2\rho(r - t)} |\bar{p}(r - t)| |\bar{x} - \bar{x}^+(t)|^2.$$

Equivalently,

$$\left\langle -\bar{p}(r - t), \int_0^t -F_{-\bar{p}(r-s)}(\bar{x}^+(s)) ds \right\rangle \leq \frac{1}{2\rho(r - t)} |\bar{p}(r - t)| \left| \int_0^t -F_{-\bar{p}(r-s)}(\bar{x}^+(s)) ds \right|^2.$$

Dividing both sides of the above inequality by t , we get

$$\left\langle -\bar{p}(r - t), \frac{\int_0^t -F_{-\bar{p}(r-s)}(\bar{x}^+(s)) ds}{t} \right\rangle \leq \frac{1}{2\rho(r - t)} |\bar{p}(r - t)| \frac{\left| \int_0^t -F_{-\bar{p}(r-s)}(\bar{x}^+(s)) ds \right|^2}{t}.$$

As $t \rightarrow 0$, we obtain

$$\langle -\bar{p}(r), -F_{-\bar{p}(r)}(\bar{x}) \rangle \leq 0.$$

This implies that $H(\bar{x}, -\bar{p}(r)) \geq 0$. \square

LEMMA 4.6. *Suppose $\bar{x} \in \mathcal{O}$ is not a local maximum of $T(\cdot)$. Let $r = T(\bar{x})$ and let $\bar{x}^+(\cdot)$ be an optimal trajectory steering \bar{x} to \mathcal{S} in time r . Let $\bar{p}(\cdot)$ be the arc given by Lemma 4.4, and set $\lambda = H(\bar{x}, -\bar{p}(r))$. Then there exists a positive constant ρ_T such that $(-\bar{p}(r), \lambda) \in N_{\text{hypo}(T_{|\mathcal{O}})}^P(\bar{x}, T_{|\mathcal{O}}(\bar{x}))$ is realized by a ball of radius ρ_T ; i.e., for all $\bar{y} \in \mathcal{O}$ and $\beta \leq T(\bar{y})$,*

$$(4.18) \quad \left\langle \frac{(-\bar{p}(r), \lambda)}{|(-\bar{p}(r), \lambda)|}, (\bar{y} - \bar{x}, \beta - r) \right\rangle \leq \frac{1}{2\rho_T} (|\bar{y} - \bar{x}|^2 + |\beta - r|^2).$$

Moreover, $\rho_T = \rho_T(\bar{x})$, where $\rho_T(\cdot) : \mathcal{O} \rightarrow (0, \infty)$ is a continuous function that can be computed explicitly.

Proof. Let $\bar{y} \in \mathcal{O}$. Two cases may occur:

- (i) $T(\bar{y}) < T(\bar{x})$;
- (ii) $T(\bar{y}) \geq T(\bar{x})$.

First case: $T(\bar{y}) =: r_1 < r = T(\bar{x})$. Let $\bar{x}_1 = \bar{x}^+(r - r_1)$ and write

$$(4.19) \quad \langle -\bar{p}(r), \bar{y} - \bar{x} \rangle = \langle -\bar{p}(r), \bar{y} - \bar{x}_1 \rangle + \langle -\bar{p}(r), \bar{x}_1 - \bar{x} \rangle.$$

Recalling Lemma 4.4 and noting that $\bar{y} \in \mathcal{S}'(r_1)$, we can estimate the first term in the right-hand side of the above identity as follows:

$$\begin{aligned} \langle -\bar{p}(r), \bar{y} - \bar{x}_1 \rangle &= \langle -\bar{p}(r_1), \bar{y} - \bar{x}_1 \rangle + \langle -\bar{p}(r) + \bar{p}(r_1), \bar{y} - \bar{x}_1 \rangle \\ &\leq \frac{1}{2\rho(r_1)} |\bar{p}(r_1)| |\bar{y} - \bar{x}_1|^2 + |\bar{p}(r) - \bar{p}(r_1)| |\bar{y} - \bar{x}_1|. \end{aligned}$$

From Lemmas 3.5 and 3.7, we have that

$$|\bar{p}(r_1)| \leq e^{K(r-r_1)} |\bar{p}(r)|, \quad |\bar{p}(r) - \bar{p}(r_1)| \leq Ke^{K(r-r_1)}(r - r_1) |\bar{p}(r)|$$

and

$$|\bar{y} - \bar{x}_1| \leq |\bar{y} - \bar{x}| + |\bar{x}_1 - \bar{x}| \leq |\bar{y} - \bar{x}| + K_2(|\bar{x}| + 1)e^{K_2(r-r_1)}(r - r_1).$$

Thus, observing that $\rho(r_1) \geq \rho(r)$, one can get the estimate

$$(4.20) \quad \langle -\bar{p}(r), \bar{y} - \bar{x}_1 \rangle \leq L_1(|x|, r)|\bar{p}(r)|(|\bar{y} - \bar{x}|^2 + |r - r_1|^2),$$

where

$$L_1(|x|, r) = \frac{1 + K_2^2(|x| + 1)^2 e^{2K_2r}}{2\rho(r)} e^{Kr} + KK_2(|x| + 1)e^{(K+K_2)r} + 2Ke^{Kr}.$$

We rewrite the rightmost term of (4.19) as

$$\begin{aligned} \langle -\bar{p}(r), \bar{x}_1 - \bar{x} \rangle &= \left\langle -\bar{p}(r), \int_0^{r-r_1} F_{-\bar{p}(r-s)}(\bar{x}^+(s)) ds \right\rangle \\ &= \int_0^{r-r_1} \langle -\bar{p}(r), F_{-\bar{p}(r-s)}(\bar{x}^+(s)) \rangle ds \end{aligned}$$

and observe that

$$\begin{aligned} \langle -\bar{p}(r), F_{-\bar{p}(r-s)}(\bar{x}^+(s)) \rangle &= \langle -\bar{p}(r), F_{-\bar{p}(r-s)}(\bar{x}^+(s)) - F_{-\bar{p}(r)}(\bar{x}^+(s)) \rangle \\ &\quad + \langle -\bar{p}(r), F_{-\bar{p}(r)}(\bar{x}^+(s)) - F_{-\bar{p}(r)}(\bar{x}) \rangle + \langle -\bar{p}(r), F_{-\bar{p}(r)}(\bar{x}) \rangle. \end{aligned}$$

Moreover, recalling that $\lambda = H(\bar{x}, -\bar{p}(r))$, we have

$$\langle -\bar{p}(r), F_{-\bar{p}(r)}(\bar{x}) \rangle = H(\bar{x}, -\bar{p}(r)) = \lambda,$$

$$\begin{aligned} \langle -\bar{p}(r), F_{-\bar{p}(r)}(\bar{x}^+(s)) - F_{-\bar{p}(r)}(\bar{x}) \rangle &\leq K |\bar{p}(r)| |\bar{x}^+(s) - \bar{x}| \\ &\leq KK_2(|\bar{x}| + 1)e^{K_2r} |\bar{p}(r)| s, \end{aligned}$$

and

$$\begin{aligned} &\langle -\bar{p}(r), F_{-\bar{p}(r-s)}(\bar{x}^+(s)) - F_{-\bar{p}(r)}(\bar{x}^+(s)) \rangle \\ &= \langle -\bar{p}(r), F_{-\bar{p}(r-s)}(\bar{x}^+(s)) \rangle - H(\bar{x}^+(s), -\bar{p}(r)) \\ &= \langle -\bar{p}(r) + \bar{p}(r-s), F_{-\bar{p}(r-s)}(\bar{x}^+(s)) \rangle + H(\bar{x}^+(s), -\bar{p}(r-s)) - H(\bar{x}^+(s), -\bar{p}(r)) \\ &\leq 2K_2(|\bar{x}^+(s)| + 1) |\bar{p}(r) - \bar{p}(r-s)| \leq 2KK_2(|\bar{x}| + 1)e^{(K_2+K)r} |\bar{p}(r)| s. \end{aligned}$$

Therefore,

$$\langle -\bar{p}(r), F_{-\bar{p}(r-s)}(\bar{x}^+(s)) \rangle \leq \lambda + L_2(|\bar{x}|, r) |\bar{p}(r)| s,$$

where $L_2(|\bar{x}|, r) = KK_2(|\bar{x}| + 1)(2e^{Kr} + 1)e^{K_2r}$. Thus, in view of the above estimates,

$$(4.21) \quad \langle -\bar{p}(r), \bar{x}_1 - \bar{x} \rangle \leq \lambda(r - r_1) + \frac{L_2(|\bar{x}|, r)}{2} |\bar{p}(r)| |r - r_1|^2.$$

Combining (4.19), (4.20), and (4.21), we get

$$\langle -\bar{p}(r), \bar{y} - \bar{x} \rangle + \lambda(r_1 - r) \leq \frac{2L_1(|\bar{x}|, r) + L_2(|\bar{x}|, r)}{2} |\bar{p}(r)| (|\bar{y} - \bar{x}|^2 + |r_1 - r|^2).$$

From Lemma 4.5, we have that $\lambda \geq 0$. Therefore, since $r_1 < r$, we conclude that

$$(4.22) \quad \left\langle \frac{(-\bar{p}(r), \lambda)}{|(-\bar{p}(r), \lambda)|}, (\bar{y} - \bar{x}, \beta - r) \right\rangle \leq \frac{2L_1(|\bar{x}|, r) + L_2(|\bar{x}|, r)}{2} (|\bar{y} - \bar{x}|^2 + |\beta - r|^2)$$

for all $\beta \leq r_1$. So, if $T(\bar{y}) < T(\bar{x})$, then (4.18) holds true provided ρ_T is such that

$$(4.23) \quad \rho_T \leq \frac{1}{2L_1(|\bar{x}|, r) + L_2(|\bar{x}|, r)}.$$

Second case: $T(\bar{y}) = r_1 \geq r = T(\bar{x})$.

In view of Lemmas 4.4 and 4.5, we already know that (4.18) holds for all $\beta \leq r$ provided $\rho_T \leq \rho(r)$. So, we just need to prove (4.18) for $r < \beta \leq r_1$. Let $\bar{y}^+(\cdot)$ be the solution of

$$(4.24) \quad \begin{cases} \dot{\bar{y}}(s) \in F_{-\bar{p}(r)}(\bar{y}(s)) & \text{a.e. } s \in [0, r_1 - \beta], \\ \bar{y}(0) = \bar{y}. \end{cases}$$

Set $\bar{y}_1 = \bar{y}^+(\beta - r)$ and compute

$$(4.25) \quad \langle -\bar{p}(r), \bar{y} - \bar{x} \rangle = \langle -\bar{p}(r), \bar{y} - \bar{y}_1 \rangle + \langle -\bar{p}(r), \bar{y}_1 - \bar{x} \rangle.$$

Since $r < \beta \leq r_1$, one can see that $T(\bar{y}_1) \geq r$. Thus, $\bar{y}_1 \in \mathcal{S}'(r)$. Then, recalling Lemma 4.4 we get

$$\langle -\bar{p}(r), \bar{y}_1 - \bar{x} \rangle \leq \frac{1}{2\rho(r)} |\bar{p}(r)| |\bar{y}_1 - \bar{x}|^2.$$

Using Lemma 3.7, we also have

$$\begin{aligned} |\bar{y}_1 - \bar{x}| &\leq |\bar{y}_1 - \bar{y}| + |\bar{y} - \bar{x}| \\ &\leq K_2(|\bar{y}| + 1)e^{K_2(\beta-r)}|\beta - r| + |\bar{y} - \bar{x}|. \end{aligned}$$

So,

$$(4.26) \quad \langle -\bar{p}(r), \bar{y}_1 - \bar{x} \rangle \leq \frac{K_2^2(|\bar{y}| + 1)^2 e^{2K_2(\beta-r)} + 1}{2\rho(r)} |\bar{p}(r)| (|\bar{y} - \bar{x}|^2 + |\beta - r|^2).$$

On the other hand, recalling (3.2) we have

$$\begin{aligned} \langle -\bar{p}(r), \bar{y} - \bar{y}_1 \rangle &= \left\langle -\bar{p}(r), \int_0^{\beta-r} -F_{-\bar{p}(r)}(\bar{y}^+(s)) ds \right\rangle \\ &= \int_0^{\beta-r} \left\langle -\bar{p}(r), -F_{-\bar{p}(r)}(\bar{y}^+(s)) \right\rangle ds \\ &= \int_0^{\beta-r} \left\langle -\bar{p}(r), -F_{-\bar{p}(r)}(\bar{y}^+(s)) + F_{-\bar{p}(r)}(\bar{x}) \right\rangle ds + \int_0^{\beta-r} \left\langle -\bar{p}(r), -F_{-\bar{p}(r)}(\bar{x}) \right\rangle ds \\ &\leq K_1 |\bar{p}(r)| \int_0^{\beta-r} |\bar{y}^+(s) - \bar{x}| ds + \int_0^{\beta-r} -H(x, -\bar{p}(r)) ds \\ &= K_1 |\bar{p}(r)| \int_0^{\beta-r} |\bar{y}^+(s) - \bar{x}| ds + \lambda(r - \beta). \end{aligned}$$

Owing to Lemma 3.7, for all $s \in [0, \beta - r]$

$$\begin{aligned} |\bar{y}^+(s) - \bar{x}| &\leq |\bar{y}^+(s) - \bar{y}| + |\bar{y} - \bar{x}| \\ &\leq K_2(|\bar{y}| + 1)e^{K_2(\beta-r)}|\beta - r| + |\bar{y} - \bar{x}|. \end{aligned}$$

Therefore,

$$(4.27) \quad \langle -\bar{p}(r), \bar{y} - \bar{y}_1 \rangle \leq \lambda(r - \beta) + K_1[1 + K_2(|\bar{y}| + 1)e^{K_2(\beta-r)}] |\bar{p}(r)| (|\bar{y} - \bar{x}|^2 + |\beta - r|^2).$$

Combining (4.25), (4.26), and (4.27), we get

$$\left\langle \frac{(-\bar{p}(r), \lambda)}{|(-\bar{p}(r), \lambda)|}, (\bar{y} - \bar{x}, \beta - r) \right\rangle \leq L_3 (|\bar{y} - \bar{x}|^2 + |\beta - r|^2),$$

where $L_3 = \frac{K_2^2(|\bar{y}|+1)^2 e^{2K_2(\beta-r)} + 1}{2\rho(r)} + K_1[1 + K_2(|\bar{y}| + 1)e^{K_2(\beta-r)}]$. The dependence of L_3 on $|\bar{y}|$ can be easily disposed of taking

$$L_4(|\bar{x}|, r) = \frac{K_2^2(|\bar{x}| + 2)^2 e^{2K_2} + 1}{2\rho(r)} + K_1[1 + K_2(|\bar{x}| + 2)e^{K_2}] + 1.$$

Then, the above inequality yields

$$(4.28) \quad \left\langle \frac{(-\bar{p}(r), \lambda)}{|(-\bar{p}(r), \lambda)|}, (\bar{y} - \bar{x}, \beta - r) \right\rangle \leq L_4(|\bar{x}|, r) (|\bar{y} - \bar{x}|^2 + |\beta - r|^2).$$

Recalling (4.23) and (4.28), and taking

$$(4.29) \quad \rho_T := \left(\max \{2L_1(|\bar{x}|, T(\bar{x})) + L_2(|\bar{x}|, T(\bar{x})), 2L_4(|\bar{x}|, T(\bar{x}))\} \right)^{-1},$$

we obtain (4.18). Finally, since $T(\cdot)$ is continuous on \mathcal{O} , one can easily see that if we set $\rho_T(\bar{x}) = \rho_T$, then $\rho_T(\cdot)$ is also continuous on \mathcal{O} . The proof is complete. \square

Proof of Theorem 4.1. Let $\bar{x} \in \mathcal{O}$. Let $r = T(\bar{x})$ and let $\bar{x}^+(\cdot)$ be an optimal trajectory steering \bar{x} to \mathcal{S} in time r . By the dynamic programming principle, $T(\bar{x}^+(t)) = r - t$ for all $t \in (0, r)$. This implies that $\bar{x}^+(t)$ is not a local maximum of $T(\cdot)$ for all $t \in (0, r)$. Therefore, by applying Lemma 4.6, we obtain that for all $t > 0$ sufficiently small, there exists a unit vector $\bar{q}(t) \in N_{\text{hypo}(T|_{\mathcal{O}})}^P(\bar{x}^+(t), T|_{\mathcal{O}}(\bar{x}^+(t)))$ realized by a ball of radius $\rho_T(\bar{x}^+(t))$ where $\rho_T(\cdot)$ is given by (4.29); i.e., for all $\bar{y} \in \mathcal{O}$ and $\beta \leq T(\bar{y})$,

$$(4.30) \quad \begin{aligned} \left\langle \bar{q}(t), (\bar{y} - \bar{x}^+(t), \beta - T(\bar{x}^+(t))) \right\rangle \\ \leq \frac{1}{2\rho_T(\bar{x}^+(t))} (|\bar{y} - \bar{x}^+(t)|^2 + |\beta - T(\bar{x}^+(t))|^2). \end{aligned}$$

Since $\bar{q}(t)$ is a unit vector in \mathbb{R}^{n+1} for all $t > 0$ sufficiently small, there exists a sequence $\{t_k\}$ which converges to 0^+ such that the sequence $\{\bar{q}(t_k)\}$ converges to a unit vector \bar{q} in \mathbb{R}^{n+1} . Taking $t = t_k$ and then letting $k \rightarrow \infty$ in (4.30), by the continuity of $T(\cdot)$ and $\rho_T(\cdot)$ in \mathcal{O} , we obtain that for all $\bar{y} \in \mathcal{O}$ and $\beta \leq T(\bar{y})$,

$$\left\langle \bar{q}, (\bar{y} - \bar{x}, \beta - T(\bar{x})) \right\rangle \leq \frac{1}{2\rho_T(\bar{x})} (|\bar{y} - \bar{x}|^2 + |\beta - T(\bar{y})|^2),$$

where $\rho_T(\cdot)$ is given by (4.29). Therefore, $\bar{q} \in N_{\text{hypo}(T|_{\mathcal{O}})}^P(\bar{x}, T|_{\mathcal{O}}(\bar{x}))$ is realized by a ball of radius $\rho_T(T(\bar{x}))$. The proof is complete. \square

We conclude this part with an example where Petrov’s controllability condition does not hold, and the minimum time function T is just continuous. Moreover, multifunction F admits no C^1 parameterization even though F and H satisfy assumptions (F) and (H). Therefore, the results in [6, 11, 12] do not apply to this example while our results do.

Example 1. Set

$$\gamma(t) = \begin{cases} (1, t), & t \leq 0, \\ (1 - \sqrt{-t^2 + 2t}, t), & 0 \leq t \leq 1, \\ (0, t), & t \geq 1. \end{cases}$$

We set the target \mathcal{S} to be the right part of $\mathbb{R}^2 \setminus \{\gamma\}$ (see Figure 1) and the differential inclusion to be

$$(\dot{x}_1(t), \dot{x}_2(t)) \in F(x_1(t), x_2(t)) = \left\{ (u_1, h(x_2(t))u_2) \mid u_1, u_2 \in [0, 1] \right\},$$

where

$$h(x_2) = \begin{cases} 0 & \text{if } x_2 \leq 1, \\ x_2 - 1 & \text{if } x_2 \geq 1. \end{cases}$$




FIG. 1.

Observe first that \mathcal{S} has the inner ball property. Observe furthermore that for $0 < t \leq 1$, the point $z_t = (1 - \sqrt{-t^2 + 2t}, t)$ is on the boundary of \mathcal{S} , and

$$\min_{v \in F(z_t)} \langle v, \nu \rangle = -\sqrt{-t^2 + 2t} |\nu|,$$

where ν is the proximal vector to \mathcal{S} . Therefore, since $\lim_{t \rightarrow 0^+} \sqrt{-t^2 + 2t} = 0$, one can see that Petrov’s controllability condition (1.5) does not hold in a neighborhood of $(1, 0)$. Moreover, the minimum time function T equals

$$T(x_1, x_2) = \begin{cases} 1 - x_1 & \text{if } x_1 \leq 1, \ x_2 \leq 0, \\ 1 - \sqrt{-x_2^2 + 2x_2} - x_1 & \text{if } x_1 \leq 1 - \sqrt{-x_2^2 + 2x_2}, \ 0 < x_2 \leq 1, \\ -x_1 & \text{if } x_1 \leq 0, \ x_2 > 1. \end{cases}$$

So, T is continuous but is not Lipschitz at points $(x_1, 0)$ for $x_1 \leq 1$.

We next show that F admits no C^1 parameterization. We first recall a criterion from [8, p. 3]: if F admits a C^1 parameterization, then the Hamiltonian H (see (1.4)) necessarily has the property

$$(4.31) \quad H(x, p) = -H(x, -p) \implies \partial_x H(x, p) = -\partial_x H(x, -p),$$

where ∂_x denotes the Clarke partial subgradient in x . In this example, the Hamiltonian H is computed as

$$H((x_1, x_2), (p_1, p_2)) = \begin{cases} 0, & p_1 < 0, p_2 < 0, \\ p_1, & p_1 \geq 0, p_2 < 0, \\ h(x_2)p_2, & p_1 < 0, p_2 \geq 0, \\ p_1 + h(x_2)p_2, & p_1 \geq 0, p_2 \geq 0. \end{cases}$$

At the point $(x_1, x_2) = (1, 1)$, one has that $H((1, 1), (0, -1)) = H((1, 1), (0, 1)) = 0$. However,

$$\partial_x H((1, 1), (0, -1)) = (0, 0) \quad \text{and} \quad \partial_x H((1, 1), (0, 1)) = (0, [0, 1]).$$

Since (4.31) is violated at the point $(1, 1)$, there is no C^1 parameterization of F .

Finally, since h is a convex function, one can also prove that F and H satisfy assumptions (F) and (H). Therefore, by applying Theorem 4.1, the hypograph of T satisfies a $\rho_T(\cdot)$ -exterior sphere condition.

4.2. Part II. In this second part, we will study the attainable set $\mathcal{A}(T)$ from 0 for the reversed differential inclusion

$$(4.32) \quad \begin{cases} \dot{x}(t) \in -F(x(t)) & \text{a.e.}, \\ x(0) = 0. \end{cases}$$

For any $T > 0$, such a set is defined by

$$\mathcal{A}(T) := \{y(t) \mid t \in [0, T] \text{ and } y(\cdot) \text{ is a solution of (4.32)}\}.$$

Let us recall that c_0, K , and K_1 are the constants appearing in (H1), (3.1), and (3.2), respectively.

THEOREM 4.7. *Assume F satisfies (F) and (H). In addition, suppose that, for some $R > 0$ and all $x \in \mathbb{R}^n$, $F(x)$ has the inner ball property of radius R . If $T > 0$ and $e^{-3KT} > 2c_0RT^2$, then the attainable set $\mathcal{A}(T)$ has the inner ball property of radius*

$$(4.33) \quad R(T) = R \frac{(e^{-3KT} - 2c_0RT^2)}{(1 + KT + K_1T)^2}.$$

Proof. Let $\bar{x} \in \partial\mathcal{A}(T)$ and let $\bar{x}^-(\cdot)$ be a trajectory of (4.32) steering 0 to \bar{x} in time T . By the Pontryagin maximum principle, there exists an arc $\bar{p}(\cdot)$ defined in $[0, T]$, with $\bar{p}(s) \neq 0$ for all $s \in [0, T]$, such that

$$(4.34) \quad \begin{cases} -\dot{\bar{p}}(s) \in \partial_x H(\bar{x}^-(s), -\bar{p}(s)) & \text{a.e. } s \in [0, T], \\ \dot{\bar{x}}^-(s) = -F_{-\bar{p}(s)}(\bar{x}^-(s)) & \text{a.e. } s \in [0, T]. \end{cases}$$

We want to prove that, for $r_0 := R(T)$ (where $R(T)$ is defined in (4.33)),

$$(4.35) \quad B\left(\bar{x} + r_0T \frac{-\bar{p}(T)}{|\bar{p}(T)|}, r_0T\right) \subseteq \mathcal{A}(T).$$

Equivalently,

$$\bar{x} - r_0 T \left(\frac{\bar{p}(T)}{|\bar{p}(T)|} - \theta \right) \in \mathcal{A}(T) \quad \forall \theta \in B(0, 1).$$

Let $\theta \in B(0, 1)$. Considering the adjoint equation associated with $\bar{p}(\cdot)$, that is,

$$(4.36) \quad \begin{cases} \dot{\bar{z}}(s) = -\frac{\langle \dot{\bar{p}}(s), \bar{z}(s) \rangle}{|\bar{p}(s)|^2} \bar{p}(s) & \text{a.e.}, \\ \bar{z}(T) = \frac{\bar{p}(T)}{|\bar{p}(T)|} - \theta, \end{cases}$$

one can see that

$$(4.37) \quad \langle \dot{\bar{z}}(s), \bar{p}(s) \rangle = -\langle \dot{\bar{p}}(s), \bar{z}(s) \rangle \quad \text{for a.e. } s \in [0, T].$$

This implies that $\frac{d}{ds} \langle \bar{z}(s), \bar{p}(s) \rangle = 0$ for a.e. $s \in [0, T]$. Therefore, $\langle \bar{z}(s), \bar{p}(s) \rangle$ is constant for all $s \in [0, T]$. In particular,

$$(4.38) \quad \langle \bar{z}(s), \bar{p}(s) \rangle = \langle \bar{z}(T), \bar{p}(T) \rangle.$$

On the other hand, from (4.36) we have $|\dot{\bar{z}}(s)| \leq K|\bar{z}(s)|$. Thus, recalling Lemma 3.5 we obtain

$$(4.39) \quad e^{-K(t_2-t_1)} |\bar{z}(t_2)| \leq |\bar{z}(t_1)| \leq e^{K(t_2-t_1)} |\bar{z}(t_2)| \quad \forall 0 \leq t_1 \leq t_2 \leq T.$$

Setting

$$(4.40) \quad \bar{y}_\theta(s) = \bar{x}(s) - r_0 s \bar{z}(s),$$

we have $\bar{y}_\theta(T) = \bar{x} - r_0 T \left(\frac{\bar{p}(T)}{|\bar{p}(T)|} - \theta \right)$. Thus, our aim is now to prove that $\bar{y}_\theta(T) \in \mathcal{A}(T)$. Since $\bar{y}_\theta(0) = \bar{x}(0) = 0$, we need only show

$$\dot{\bar{y}}_\theta(s) \in -F(\bar{y}_\theta(s)) \quad \text{for a.e. } s \in [0, T].$$

Observe that $F_{-\bar{p}(s)}(\bar{y}_\theta(s)) \in \partial F(\bar{y}_\theta(s))$. Since $F(\bar{y}_\theta(s))$ is convex and has the inner ball property of radius R , we have that $\bar{p}(s)$ is an inner normal vector to $\partial F(\bar{y}_\theta(s))$ at the point $F_{-\bar{p}(s)}(\bar{y}_\theta(s))$. Thus, $-\dot{\bar{y}}_\theta(s) \in F(\bar{y}_\theta(s))$ (equivalently, $\dot{\bar{y}}_\theta(s) \in -F(\bar{y}_\theta(s))$) if $-\dot{\bar{y}}_\theta(s) \in B(F_{-\bar{p}(s)}(\bar{y}_\theta(s)) + R \frac{\bar{p}(s)}{|\bar{p}(s)|}, R)$. Therefore,² our conclusion will follow from

$$(4.41) \quad \left\langle \frac{\bar{p}(s)}{|\bar{p}(s)|}, -\dot{\bar{y}}_\theta(s) - F_{-\bar{p}(s)}(\bar{y}_\theta(s)) \right\rangle \geq \frac{1}{2R} |\dot{\bar{y}}_\theta(s) + F_{-\bar{p}(s)}(\bar{y}_\theta(s))|^2.$$

Equivalently,

$$(4.42) \quad \left\langle -\frac{\bar{p}(s)}{|\bar{p}(s)|}, \dot{\bar{y}}_\theta(s) + F_{-\bar{p}(s)}(\bar{y}_\theta(s)) \right\rangle \geq \frac{1}{2R} |\dot{\bar{y}}_\theta(s) + F_{-\bar{p}(s)}(\bar{y}_\theta(s))|^2.$$

We are now going to prove (4.42). On account of (4.40), we have

$$\dot{\bar{y}}_\theta(s) = -F_{-\bar{p}(s)}(\bar{x}(s)) - r_0 \bar{z}(s) - r_0 s \dot{\bar{z}}(s).$$

²Observe that for all $R > 0$ and $x \in \mathbb{R}^N$, $y \in \overline{B}(x + Rv, R) \Leftrightarrow \langle v, y - x \rangle \geq \frac{1}{2R} |y - x|^2$, where $v \in \mathbb{R}^N$ is any unit vector.

Thus,

$$\begin{aligned} & \left\langle -\frac{\bar{p}(s)}{|\bar{p}(s)|}, \dot{y}_\theta(s) + F_{-\bar{p}(s)}(\bar{y}_\theta(s)) \right\rangle \\ &= \left\langle -\frac{\bar{p}(s)}{|\bar{p}(s)|}, F_{-\bar{p}(s)}(\bar{y}_\theta(s)) - F_{-\bar{p}(s)}(\bar{x}(s)) - r_0\bar{z}(s) - r_0s\dot{\bar{z}}(s) \right\rangle \\ &= \frac{1}{|\bar{p}(s)|} [H(\bar{y}_\theta(s), -\bar{p}(s)) - H(\bar{x}(s), -\bar{p}(s))] \\ &\quad + \frac{r_0}{|\bar{p}(s)|} \langle \bar{p}(s), \bar{z}(s) \rangle + r_0s \frac{1}{|\bar{p}(s)|} \langle \bar{p}(s), \dot{\bar{z}}(s) \rangle. \end{aligned}$$

Recalling (4.37), (4.40), and (4.38), we conclude that

$$\begin{aligned} & \left\langle -\frac{\bar{p}(s)}{|\bar{p}(s)|}, \dot{y}_\theta(s) + F_{-\bar{p}(s)}(\bar{y}_\theta(s)) \right\rangle \\ &= \frac{1}{|\bar{p}(s)|} [H(\bar{y}_\theta(s), -\bar{p}(s)) - H(\bar{x}(s), -\bar{p}(s)) - \langle -\dot{\bar{p}}(s), \bar{y}_\theta(s) - \bar{x}(s) \rangle] + \frac{r_0}{|\bar{p}(s)|} \langle \bar{p}(T), \bar{z}(T) \rangle \\ &\quad \geq -c_0 |\bar{y}_\theta(s) - \bar{x}(s)|^2 + r_0 \frac{|\bar{p}(T)|}{|\bar{p}(s)|} \left\langle \frac{\bar{p}(T)}{|\bar{p}(T)|}, \frac{\bar{p}(T)}{|\bar{p}(T)|} - \theta \right\rangle \\ &\quad \geq -c_0 r_0^2 s^2 |\bar{z}(s)|^2 + \frac{r_0}{2} \frac{|\bar{p}(T)|}{|\bar{p}(s)|} \left| \frac{\bar{p}(T)}{|\bar{p}(T)|} - \theta \right|^2 \\ &\quad = -c_0 r_0^2 s^2 |\bar{z}(s)|^2 + \frac{r_0}{2} \frac{|\bar{p}(T)|}{|\bar{p}(s)|} |\bar{z}(T)|^2. \end{aligned}$$

Recalling Lemma 3.5 and (4.39), we obtain

$$(4.43) \quad \left\langle -\frac{\bar{p}(s)}{|\bar{p}(s)|}, \dot{y}_\theta(s) + F_{-\bar{p}(s)}(\bar{y}_\theta(s)) \right\rangle \geq \frac{r_0}{2} (-2c_0 r_0 T^2 + e^{-3KT}) |\bar{z}(s)|^2.$$

Observe that $0 < r_0 = R(T) = R \frac{(e^{-3KT} - 2c_0 RT^2)}{(1 + KT + K_1 T)^2} \leq R$. Then,

$$(4.44) \quad \left\langle -\frac{\bar{p}(s)}{|\bar{p}(s)|}, \dot{y}_\theta(s) + F_{-\bar{p}(s)}(\bar{y}_\theta(s)) \right\rangle \geq \frac{r_0}{2} (-2c_0 RT^2 + e^{-3KT}) |\bar{z}(s)|^2.$$

On the other hand,

$$\begin{aligned} |\dot{y}_\theta(s) + F_{-\bar{p}(s)}(\bar{y}_\theta(s))| &\leq |F_{-\bar{p}(s)}(\bar{y}_\theta(s)) - F_{-\bar{p}(s)}(\bar{x}(s))| + r_0 |\bar{z}(s)| + r_0 s |\dot{\bar{z}}(s)| \\ &\leq K_1 r_0 s |\bar{z}(s)| + r_0 |\bar{z}(s)| + K r_0 s |\bar{z}(s)| \\ &\leq r_0 (K_1 T + KT + 1) |\bar{z}(s)|. \end{aligned}$$

Thus,

$$\left\langle -\frac{\bar{p}(s)}{|\bar{p}(s)|}, \dot{y}_\theta(s) + F_{-\bar{p}(s)}(\bar{y}_\theta(s)) \right\rangle \geq \frac{e^{3KT} - 2c_0 RT^2}{2r_0 (K_1 T + KT + 1)^2} |\dot{y}_\theta(s) + F_{-\bar{p}(s)}(\bar{y}_\theta(s))|^2,$$

and (4.42) follows. The proof is complete. \square

Finally, let us denote by $\mathcal{A}(x, T)$ the attainable set from x in time T for the differential inclusion in (4.32). One can see from Theorem 4.7 that there exists a time

$T_0 > 0$ such that for all $0 < T < T_0$, the set $\mathcal{A}(x, T)$ has the inner ball property of radius $R(T)$ given by (4.33). Moreover, for any closed set $\mathcal{S} \subset \mathbb{R}^N$, let us set

$$\mathcal{A}(\mathcal{S}, T) = \bigcup_{x \in \mathcal{S}} \mathcal{A}(x, T).$$

COROLLARY 4.8. *Suppose that \mathcal{S} is nonempty and closed. Under the assumptions in Theorem 4.7, there exists $T_0 > 0$ such that, for all $0 < T < T_0$, then the set $\mathcal{A}(\mathcal{S}, T)$ has the inner ball property of radius $R(T)$ given by (4.33).*

Applying Theorem 4.1 and the above results to the minimum time function for a general target, we obtain the following theorem together with useful corollaries.

THEOREM 4.9. *Assume (F), (H), and suppose $F(x)$ has the inner ball property of radius R for some $R > 0$ and all $x \in \mathbb{R}^n$. Suppose further that \mathcal{S} is nonempty and closed and that $T(\cdot)$ is continuous in an open subset \mathcal{O} of \mathcal{C} . Then, the hypograph of $T|_{\mathcal{O}}(\cdot)$ satisfies the $\rho_T(\cdot)$ -exterior sphere condition for some continuous function $\rho_T(\cdot) : \mathcal{O} \rightarrow (0, \infty)$.*

Proof. We define, for any $0 < t < T_0$,

$$\mathcal{O}^t = \{x \in \mathcal{O} \mid T(x) > t\}, \quad \mathcal{S}^t = \mathcal{A}(\mathcal{S}, t),$$

and $T^t(\cdot)$ is the minimum time function for the differential inclusion (1.1) with the target \mathcal{S}_t . One can see that

$$T^t(x) = T(x) - t \quad \forall x \in \mathcal{O}^t.$$

Since \mathcal{S}^t has the inner ball property of radius $R(t) > 0$, by applying Theorem 4.1 to $T^t(\cdot)$, we obtain that the hypograph of $T^t|_{\mathcal{O}^t}(\cdot)$ satisfies a $\rho_T^t(\cdot)$ -exterior sphere condition for some continuous function $\rho_T^t(\cdot) : \mathcal{O}^t \rightarrow (0, \infty)$. Hence, the hypograph of $T|_{\mathcal{O}}(\cdot)$ satisfies the $\rho_T^t(\cdot)$ -exterior sphere condition. Observe that, since $\mathcal{O}^{t_1} \subseteq \mathcal{O}^{t_2} \subseteq \mathcal{O}$ for $0 < t_1 < t_2 < T_0$ and $\bigcup_{t \in (0, T_0)} \mathcal{O}^t = \mathcal{O}$, one can prove that $T|_{\mathcal{O}}(\cdot)$ satisfies a $\rho_T(\cdot)$ -exterior sphere condition for some continuous function $\rho_T(\cdot) : \mathcal{O} \rightarrow (0, \infty)$. \square

COROLLARY 4.10. *Under the assumptions of Theorem 4.9, $T|_{\mathcal{O}}(\cdot)$ satisfies properties (1)–(3) in Proposition 2.3.*

COROLLARY 4.11. *Under the assumptions of Theorem 4.9, if $T(\cdot)$ is locally Lipschitz in \mathcal{O} , then $T(\cdot)$ is locally semiconcave in \mathcal{O} .*

Acknowledgments. Part of this work was completed during the C.I.M.E. course “Control of partial differential equations” (Cetraro, July 19–23, 2010) and the trimester on “Control of partial differential equations and applications” held at Institute Henri Poincaré (Paris, Oct.–Dec. 2010). The authors wish to express their gratitude to the above institutions for their support and hospitality. The authors are grateful to the anonymous referees for their careful reading of the manuscript, and for their comments, which highly improved the quality of this paper.

REFERENCES

[1] J.-P. AUBIN AND H. FRANKOWSKA, *Set-Valued Analysis*, Birkhäuser, Boston, 1990.
 [2] A. CANINO, *On p-convex sets and geodesics*, J. Differential Equations, 75 (1988), pp. 118–157.
 [3] P. CANNARSA AND H. FRANKOWSKA, *Some characterizations of optimal trajectories in control theory*, SIAM J. Control Optim., 29 (1991), pp. 1322–1347.
 [4] P. CANNARSA AND H. FRANKOWSKA, *Interior sphere property of attainable sets and time optimal control problems*, ESAIM Control Optim. Calc. Var., 12 (2006), pp. 350–370.

- [5] P. CANNARSA, F. MARINO, AND P. R. WOLENSKI, *Semiconcavity of the minimum time function for differential inclusions*, Dyn. Contin. Discrete Impuls. Syst. Ser. B, to appear.
- [6] P. CANNARSA AND C. SINISTRARI, *Semiconcave Functions, Hamilton-Jacobi Equations, and Optimal Control*, Birkhäuser, Boston, 2004.
- [7] P. CANNARSA AND C. SINISTRARI, *Convexity properties of the minimum time function*, Calc. Var. Partial Differential Equations, 3 (1995), pp. 273–298.
- [8] P. CANNARSA AND P. R. WOLENSKI, *Semiconcavity of the value function for a class of differential inclusion*, Discrete Contin. Dyn. Syst., 29 (2011), pp. 453–466.
- [9] F. H. CLARKE, *Optimization and Nonsmooth Analysis*, Canad. Math. Soc. Ser. Monogr. Adv. Texts, John Wiley & Sons, New York, 1983.
- [10] G. COLOMBO AND A. MARIGONDA, *Differentiability properties for a class of non-convex functions*, Calc. Var. Partial Differential Equations, 25 (2006), pp. 1–31.
- [11] G. COLOMBO AND K. T. NGUYEN, *On the structure of the minimum time function*, SIAM J. Control Optim., 48 (2010), pp. 4776–4814.
- [12] K. T. NGUYEN, *Hypographs satisfying an external sphere condition and the regularity of the minimum time function*, J. Math. Anal. Appl., 372 (2010), pp. 611–628.
- [13] C. NOUR, R. J. STERN, AND J. TAKCHE, *The θ -exterior sphere condition, φ -convexity, and local semiconcavity*, Nonlinear Anal., 73 (2010), pp. 573–589.
- [14] A. ORNELAS, *Parametrization of Carathéodory multifunctions*, Rend. Sem. Mat. Univ. Padova, 83 (1990), pp. 33–44.
- [15] P. R. WOLENSKI AND Y. ZHUANG, *Proximal analysis and the minimal time function*, SIAM J. Control Optim., 36 (1998), pp. 1048–1072.