

Characterization, functional validation and gene expression patterns of two 14-3-3 isoforms from *Vitis vinifera*

G. FARACE¹, S. VISCONTI², C. MARCHIVE³, P. ADUCCI², M. MARRA² and C. TROSSAT-MAGNIN¹

¹Institut des Sciences de la Vigne et du Vin (ISVV), UMR EGFV, Villenave d'Ornon, France

²Department of Biology, University of Rome "Tor Vergata" Rome, Italy

³INRA-AgroParisTech, UMR1318, Versailles, France

Summary

In eukaryotic cells, 14-3-3 proteins constitute a family of conserved regulatory proteins which bind to a large number of phosphorylated partners. Although a great body of evidences accumulated in herbaceous species indicates that 14-3-3s play key roles in regulation of plant growth, information about function in tree plants is still very scarce. In this paper, we report the characterization of two novel 14-3-3s from grapevine (*Vitis vinifera* 'Cabernet Sauvignon') named *VV1_CS* and *VV2_CS*. The *VV1_CS* and *VV2_CS* cDNA were expressed in *Escherichia coli* and the recombinant proteins were shown to be functional proteins using plasma membrane H⁺-ATPase as target. Transcript analysis during grape development or under elicitor-induced stress revealed differential expression of the two 14-3-3 isoforms.

Key words: 14-3-3 proteins, *Vitis vinifera*.

Abbreviations: AHA1, *Arabidopsis* H⁺-ATPase isoform 1; FC, fusicoccin; *VV1_CS*, *Vitis vinifera* 14-3-3 isoform 1 'Cabernet-Sauvignon'; *VV2_CS*, *Vitis vinifera* 14-3-3 isoform 2 'Cabernet-Sauvignon'.

Introduction

14-3-3 proteins are a family of conserved dimeric proteins involved in the regulation of a wide range of many cellular processes in eukaryotes. Dimerization of 14-3-3 monomers into homo- or heterodimers allows these proteins to interact with many diverse targets through binding to conserved consensus motifs (YAFFE 2002). Plants have a variable number of 14-3-3 isoforms: *Arabidopsis* contains 13 isoforms (DELILLE *et al.* 2001), cotton six (ZHANG *et al.* 2010), rice eight (YAO *et al.* 2007) and barley five (SCHOONHEIM *et al.* 2007); soybean contains sixteen transcripts (LI and DHAUBHADEL 2011), while tobacco has 17 potential isoforms (KONAGAYA *et al.* 2004). The Grape Genome Browser (<http://www.genoscope.cns.fr/externe/GenomeBrowser/Vitis/>), created by a French-Italian Public Consortium for Grapevine Genome Characterization, identifies to date not less than seven putative 14-3-3 isoforms

(JAILLON *et al.* 2007). In agreement with their common origin, there is a high sequence conservation within and among species (WANG and SHAKES 1996). Within a species, 14-3-3s are divided into multiple groups or clades that may have different physiological roles (LI and DHAUBHADEL 2011, PIOTROWSKI and OECKING 1998).

In plants, 14-3-3 proteins are involved in several fundamental physiological processes, such as primary metabolism, ion transport, cell division and growth, hormonal signalling, as well as environmental adaptation, including response to light or to biotic and abiotic stresses (DENISON *et al.* 2011, ADUCCI *et al.* 2002). The role of 14-3-3s in the control of cell energy is also indicated by the fact that they can bind AMP, which prevents their interaction with other partners such as H⁺-ATPase (CAMONI *et al.* 2001), the master enzyme of ion transport across the plasma membrane (MICHELET and BOUTRY 1995). 14-3-3 isoforms exhibit tissue-specific expression patterns during barley embryo development (FULGOSI *et al.* 2002). At the systemic level, it is known that 14-3-3 proteins are involved in regulation of hormone pathways. 14-3-3s have well-known roles in brassinosteroid (GAMPALA *et al.* 2007, KIM *et al.* 2009) and abscisic acid signaling (CUTLER *et al.* 2010, SCHOONHEIM *et al.* 2007, DEL VISO *et al.* 2007, HONG *et al.* 2011) as well as in gibberellic acid biosynthesis (SCHWECHHEIMER and WILLIGE 2009, ISHIDA *et al.* 2004). A growing body of evidences also pinpoints a relevant role of 14-3-3 proteins in response of plants to abiotic and biotic stresses (YANG *et al.* 2009). Environmental stresses can regulate the expression and phosphorylation of 14-3-3s (SUN *et al.* 2011).

Most of the published results about the roles of plant 14-3-3 have been obtained in herbaceous/model species whereas data in tree/crop species are still very scarce, although a deeper understanding of 14-3-3 role in metabolic or developmental processes of crop species, such as for instance fruit development and ripening, could be very helpful for improvement of growth and productivity. Grapevine (*Vitis vinifera*) is one of the world's most widespread fruit crop, and during the last years, due to its economical relevance, it has become a model species for ripening studies. The present study describes the identification and characterization of two novel 14-3-3 isoforms from grape berries. Their expression profile in grape berries during development/ripening, as well as in leaves in response to elicitor challenge, has been analyzed.

Correspondence to: Dr. G. FARACE, Institut des Sciences de la Vigne et du Vin (ISVV), UMR EGFV, 210 Chemin de Leysotte, CS 50008, 33882 Villenave d'Ornon Cedex, France. E-mail: giojfarace@yahoo.com

Dedicated to the memory of Prof. SAID HAMDJ (1957-2006).

Material and Methods

Elaboration of the 14-3-3 phylogenetic tree: Database searches were conducted with the basic local alignments search tool of the National Center for Biotechnology Information (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) and via the DFCI Grape Gene Index (<http://compbio.dfci.harvard.edu/cgi-bin/tgi/gimain.pl?gudb=grape>). Alignment of amino acid sequences was performed using Clustal Omega (<http://www.ebi.ac.uk/Tools/msa/clustalo/>) with default parameters. Phylogenetic tree (using the maximum parsimony method with 1000 bootstrap replications) were made by the software MEGA5 (<http://www.megasoftware.net/>). Phylogenetic tree was performed by aligning 14-3-3 sequences from *Vitis vinifera* (VV1_CS, GenBank accession number ACO40495; VV2_CS, GenBank accession number ACO40494) and from *Arabidopsis thaliana*, accession numbers: At κ (AAD51783), At λ (AAD51781), At χ (AAA96254), At ϕ (AAB62224), At ω (AAA96253), At ν (AAD51782), At υ (AAB62225), At ψ (AAA96252), At μ (AAD51784), At ι (AAK11271), At \omicron (AAG47840), At ϵ (AAD51785), At π (NP_565174). The *Entamoeba histolytica* HMI-IMSS 14-3-3-1 isoform (Eh14-3a), accession number: U13418 was used as out-group because of its distance from the other 14-3-3s (PIOTROWSKI and OECKING 1998). For the analysis of the C-terminal end of grapevine 14-3-3s the PHD software was used (ROST and SANDER 1993).

Chemicals and reagents: Oligonucleotides were obtained from Eurogentec (Belgium). Glutathione Sepharose 4B from GE Healthcare (Uppsala, Sweden). Other chemicals were of analytical grade and were purchased from Sigma (St. Louis, Missouri, USA). Restriction enzymes were from Roche Diagnostics (Mannheim, Germany). Pfu turbo DNA polymerase was from Stratagene (La Jolla, California, USA). Protein standards (Precision plus protein standards) and other chemical for SDS-PAGE were from Bio-Rad (Hercules, California, USA).

Vitis 14-3-3 cDNA cloning and sequence analysis: A cDNA library of grape berries collected at ripening stage from 'Cabernet Sauvignon' variety has been constructed using Smart cDNA cloning kit Clontech (USA). ESTs cloned into *pTriplEx* were sequenced (Genome Express, France) and referenced by the DFCI Grape Gene Index.

Both VV1_CS and VV2_CS cDNA were amplified from *pTriplEx* by PCR using *Pfu* (Promega) and specific primers for the VV1_CS gene (forward primer 5'-AATGGATCCATGGCGGCAGCTCCATCC-3', reverse primer 5'-AATGGATCCTCACTGCTGTTCATCATCAC-3') and for the VV2_CS gene (forward primer 5'-CGGCAGCTCCATCCGCACGC-3', reverse primer 5'-GCCGAGGGAATYCAGCCGGG-3').

The VV1_CS and VV2_CS amplicons were subcloned into pGEM-T easy vector (Promega) and sequenced. These constructs were digested with appropriate enzymes and cloned into pGEX-2TK (GE Healthcare) at BamHI site for VV1_CS or between BamHI and EcoRI sites for VV2_CS. The authenticity of sequence constructs was confirmed by

restriction analysis and DNA sequencing (Genome express, France). Recombinant plasmids were electroporated in *E. coli* M15 strain (Stratagene, La Jolla, California, USA).

Recombinant protein expression and purification: VV1_CS and VV2_CS 14-3-3 proteins were expressed in M15 *E. coli* as fusion proteins with the glutathione S-transferase (GST), using pGEX-2TK vector and purified by affinity chromatography using Glutathione-Sepharose 4B (FULLONE *et al.* 1998).

SDS-PAGE and immunoblotting: SDS-PAGE (LAEMMLI 1970) was performed in a Mini Protean apparatus (Bio-Rad). Thrombin cleavage of the fusion protein GST-14-3-3 was performed when GST-fusion proteins were immobilized on the glutathione-sepharose beads in order to remove GST from the samples. For immunoblotting, samples of affinity-purified GST-14-3-3 (3 μ g) or crude protein extract from grape berries (25 μ g) were subjected to SDS-PAGE and blotted onto nitrocellulose membrane, using a semidry LKB apparatus (2 h, 0.8 mA cm⁻²). Blots were probed with anti-14-3-3 antibodies (1/500) (FULLONE *et al.* 1998). Polypeptides were visualized by alkaline phosphatase-conjugated secondary antibodies (1/10000) (Sigma), using enhanced chemiluminescence detection (ECL kit, GE Healthcare).

Overlay assay (Far Western Blotting): pGEX-2TK expression vector produces a GST-fused protein containing a cAMP-dependent protein kinase phosphorylation site and a thrombin site between the two polypeptides. The ³²P-labeled VV1_CS and VV2_CS proteins were obtained as already described (FULLONE *et al.* 1998). Briefly, 200 μ L of *E. coli* lysate containing GST-fused VV1_CS and VV2_CS proteins were incubated with 35 μ L of glutathione-sepharose beads. After washing with phosphate-buffered saline (140 mM NaCl, 2.7 mM KCl, 10 mM Na₂HPO₄ and 1.8 mM KH₂PO₄, pH 7.3), beads were incubated in 30 μ L of buffer A (20 mM Tris-HCl, 12 mM MgCl₂, 100 mM NaCl, pH 7.5) with 1.11 MBq of [γ -³²P]ATP (110 TBq·mmol⁻¹) and 10 units of the catalytic subunit of protein kinase A (Sigma) for 30 min at 30 °C. The beads were washed three times with 1 mL of phosphate-buffered saline and then incubated in 100 μ L of phosphate-buffered saline with 1 unit of thrombin (GE Healthcare) overnight at 22 °C. The mixture was centrifuged and the supernatant was collected. ³²P incorporation was measured in a liquid scintillation β -counter (LKB Wallac 1410). Specific activities of both labelled proteins were similar (about 90 GBq/mmol). The overlay assay was carried out as already described (FULLONE *et al.* 1998), with minor modifications. Briefly, two-phase partitioned maize plasma membranes (10 μ g protein) containing the H⁺-ATPase were subjected to SDS-PAGE and blotted onto nitrocellulose membrane, using a semidry LKB apparatus (2 h, 0.8 mA cm⁻²). The membrane was blocked with 5 % fat-free milk in 25 mM Hepes-OH, 75 mM KCl, 5 mM MgCl₂, 1 mM DTT, 0.1 mM EDTA, 0.05 % Tween-20, pH 7.5 (buffer H) and then incubated overnight at 4 °C in the same buffer, containing 2 % fat-free milk, 0.1 nmol of ³²P-labeled VV1_CS or VV2_CS (corresponding to 9 kBq·mL⁻¹). After incubation, the membrane was washed three times

with buffer H, dried and subjected to autoradiography at -80°C . Each overlay experiment was performed at least three times and similar results were obtained.

Purification of endoplasmic reticulum from yeast expressing AHA1: Plasma membrane H^+ -ATPase AHA1 isoform was expressed in *Saccharomyces cerevisiae* as previously described (REGENBERG *et al.* 1995). After cell homogenization, membranes were purified by differential centrifugation and endoplasmic reticulum, containing most of the AHA1, was isolated by sucrose gradient centrifugation (MARRA *et al.* 2000).

Phosphohydrolytic activity assay: AHA1 phosphohydrolytic activity of yeast endoplasmic reticulum (ER) membranes was assayed as already described (MARRA *et al.* 2000, CAMONI *et al.* 2012) with minor modifications: 10 μg of sucrose gradient purified yeast ER were preincubated with 10 μg of VV1_CS or VV2_CS protein in 500 μL of 50 mM Tris-Mes, 5 mM MgCl_2 , 50 mM KNO_3 , 5 mM NaN_3 , 0.2 mM ammonium molybdate, pH 7.2 containing 10 μM fusicoccin. After 20 min incubation, 2 mM ATP was added. ATP hydrolysis was measured as already described (SERRANO 1989).

Plant growth conditions: Rooted cuttings of *Vitis vinifera* L. 'Cabernet Sauvignon' were grown in a greenhouse (maximum temperature 24°C) in potting soil irrigated with water. Leaves, stems and roots were collected from 2-month-old rooted plants and frozen in liquid nitrogen until used.

Elicitation was made by spraying 1 mL of solution on the leaves containing known quantities of the molecule. Elicitors used for analysis were: jasmonic acid (150 μM in 1 % ethanol), ethephon (10 mM in water), salicylic acid (5 mM in water), H_2O_2 (10 mM in water), and ergosterol (200 μM in 2 % ethanol, 0.02 % triton) (MARCHIVE *et al.* 2007). For each time point of the kinetics, leaves from three plants were collected, combined and dipped in liquid nitrogen and stored at -80°C until used for RNA extraction. Berries from *Vitis vinifera* L. cultivar 'Cabernet Sauvignon' were sampled during the 2004 growing season. Each week, a sample of fifty randomly selected berries from three bunches was harvested. Stages of berry development were determined according to usual criteria (size, measures of soluble solids, softening and berry colour) as previously described (BOSS *et al.* 1996). For the identification of developmental stages, the "modified Eichorn-Lorenz system" (modified E-L system) was used (COOMBE 1995). Berries were frozen in liquid nitrogen and stored at -80°C until use.

RNA extraction and RT-PCR experiments: Total RNA was extracted from three independent samples of leaves, stems, roots and berries at different stages of development (REID *et al.* 2006). First strand cDNA synthesis was performed using 2 μg of total RNA, oligo(dT) as primer and M-MLV reverse transcriptase (Promega, Madison, WI, USA). PCR amplification using cDNAs as template was performed with gene-specific primer pairs listed below. VV1_CS gene: forward primer 5'-AATGGATCCATGGCGTCCACACTGCCG-3', reverse primer 5'-AATGAATTCCTATGGCTCATCCAAGT-3'. VV2_CS gene:

forward primer 5'-GCGTCCACACTGCCGAG-3', reverse primer 5'-AAATACCAGCACAGACCTCG-3'. An initial denaturation step (5 min at 95°C) was followed by 25 to 35 cycles of 5 s at 95°C , 45 s at 55°C , and 1 min at 72°C . PCR fragments were run on a 1.5 % (w/v) agarose gel containing ethidium bromide and visualized by UV. Fragments were purified and sequenced for identification (Genome Express, France). The expected amplicon sizes were 400 bp and 420 bp for VV1_CS and VV2_CS, respectively. The amplification of the elongation factor VV-EF1- γ cDNA (forward primer 5'-CAAGAGAAACCATC-CCTAGCTG-3', reverse primer 5'-TCAATCTGTCTAG-GAAAGGAAG-3') was used as an internal and normal control which has a constant profile over time (MARCHIVE *et al.* 2007). Each ethidium bromide-stained band on the gel was quantified using Quantity One software from BioRad (LIMA *et al.* 2012).

Statistical analysis: Two independent experiments with three technical replicates were performed for each assay, while only one representative experiment is shown. Error bars were produced by calculating the standard deviation. The gene expression analysis was performed on two independent experiments.

Results

Characterization and in silico analysis of *Vitis vinifera* cDNA clones VV1_CS and VV2_CS: Two 14-3-3 full-length cDNA were identified in the INRA EST bank. One cDNA, called VV1_CS, contained a 1157 bp insert encoding a 260 residue polypeptide with a calculated molecular weight of 29.42 kDa. The second one, called VV2_CS, contained a 1036 bp insert encoding a 252 residue polypeptide with a calculated molecular weight of 28.65 kDa. The VV1_CS and VV2_CS proteins showed 78.5 % identity at the amino-acid level and 67.9 % identity at nucleotide sequence level (data not shown). The two protein sequences were compared with a typical 14-3-3 of *Arabidopsis*, the 14-3-3 epsilon, and with the *Arabidopsis* 14-3-3 omega (Fig. 1). The alpha helices contained in all the 14-3-3s, as well as the five blocks of conserved sequences, are shown (WANG and SHAKES 1996, FERL *et al.* 1994). Two 14-3-3 signatory motifs are present in the two isoforms from grape berry (WANG and SHAKES 1996): RNLLSVAYKNVI in block 2 and SYKDSTLIMQLLRDNLWTSD in block 5, both identical in the two isoforms.

A phylogenetic tree comparing the two grapevine 14-3-3 isoforms with the *Arabidopsis* 14-3-3s (Fig. 1b) was made using the method of maximum parsimony. Group identification was performed according to the work of PIOTROWSKI and OECKING (1998) and an "out-group", represented by an *Entamoeba histolytica* 14-3-3 was added (PIOTROWSKI and OECKING 1998). The results show that VV1_CS and VV2_CS belong to the non-epsilon group of 14-3-3 proteins previously described (DELILLE *et al.* 2001) but they fall in two distinct groups, the Omega-group and the Kappa-group respectively for VV1_CS and VV2_CS, according to Pio-




Fig. 1: Comparison of amino acid sequences of 14-3-3 proteins. a): Sequence alignment of *V. vinifera* 14-3-3 proteins, the epsilon and omega isoform from *Arabidopsis thaliana*. Multiple sequence alignment was performed with the Clustal Omega software. Black bars indicate the position of the nine α -helices. Underlines indicate the five conserved sequences in all 14-3-3s. In grey are indicated the theoretical tenth alpha-helix. b): Phylogenetic tree of 14-3-3 proteins from *Arabidopsis thaliana* (DeLille *et al.* 2001) and *Vitis vinifera* constructed with the MEGA5 software using the maximum parsimony method. The 14-3-3 named “Eh14-3a” is the out-group from *Entamoeba histolytica*.

TROWSKI and OECKING (1997). A further analysis was done comparing the C-terminals from VV1_CS and VV2_CS and the *Arabidopsis* 14-3-3 omega isoform, which has been proposed to contain a tenth alpha helix (SHEN *et al.* 2003). The *in silico* analysis does not show any theoretical alpha-helix in the VV2_CS C-terminal region (Fig. 1a), whereas four residues in the VV1_CS C-terminal are compatible with an alpha-helix structure, similarly to what observed for the *Arabidopsis* 14-3-3 omega (SHEN *et al.* 2003).

Characterization of VV1_CS and VV2_CS proteins expressed in *E. coli*: *V. vinifera* 14-3-3 proteins were expressed in *E. coli* as GST (Glutathione S-Transferase)-fusion proteins and purified by affinity chromatography. The SDS-PAGE of purified proteins is shown in Fig. 2. Lanes 1 and 3 show the purified VV1_CS and VV2_CS proteins as GST-fusion




Fig. 2: SDS-PAGE and immunoblotting of VV1_CS and VV2_CS expressed in *E. coli* as GST-fusion proteins. Left panel (lanes 1 to 4): coomassie-stained gel. Right panel (lanes 5 and 6): immunoblotting of GST-fused VV1_CS and VV2_CS with anti-14-3-3 antibodies. GST-fused (lanes 1, 3, 5 and 6) or thrombin-cleaved (lanes 2 and 4) VV1_CS (lanes 1, 2 and 5) and VV2_CS (lanes 3, 4 and 6) proteins purified by affinity chromatography were loaded on the gel.

proteins, while lanes 2 and 4 show VV1_CS and VV2_CS proteins after thrombin cleavage, to remove the GST protein. The apparent molecular weight was of about 30 kDa for both proteins, in accordance with the calculated molecular weights of VV1_CS (29.42 kDa) and VV2_CS (28.65 kDa) proteins. Lanes 5 and 6 show the Western blotting of both GST-fused 14-3-3 proteins. Anti-14-3-3 antibodies, raised against a conserved plant 14-3-3 sequence, were able to immunodecorate both the GST-fused VV1_CS and VV2_CS proteins.

In order to verify the functionality of the recombinant proteins, their ability to interact with the H^+ -ATPase and to stimulate its activity was assayed. Interaction studies were performed by means of an “overlay assay” using the ^{32}P -labelled VV1_CS or VV2_CS proteins as probes and a crude plasma membrane preparation from maize roots containing the H^+ -ATPase as a bait. The autoradiography (Fig. 3a) demonstrated that both 14-3-3 proteins were able to bind to the H^+ -ATPase and that VV1_CS interacted to a higher extent, as compared to VV2_CS. The interaction was specific, since no binding with any other protein present in the crude plasma membrane fraction was observed.

To test the effect of 14-3-3 proteins on the H^+ -ATPase activity, a recombinant AHA1 H^+ -ATPase isoform (*Arabidopsis H⁺-ATPase isoform 1*) expressed in yeast was used. In this system, most of the functional enzyme is located at the endoplasmic reticulum (ER) (VILLALBA *et al.* 1992). A significant and reproducible stimulation of the H^+ -ATPase phosphohydrolytic activity can be induced by exogenous 14-3-3, providing that fusicoccin (FC) is added to the reaction mixture (CAMONI *et al.* 2001, VISCONTI *et al.* 2003). ER membranes were incubated with the GST-VV1_CS or




Fig. 3: Functional characterization of VV1_CS and VV2_CS. **a)**: Association of 14-3-3 proteins with the H^+ -ATPase. The overlay assay was performed with 10 μg of plasma membrane preparation from maize roots. **b)**: Effect of 14-3-3 proteins on the phosphohydrolytic activity of AHA1 H^+ -ATPase.

GST-VV2_CS fusion proteins in the presence of FC. The Fig. 3b shows that both VV1_CS and VV2_CS proteins were able to efficiently stimulate the enzyme activity and that VV2_CS brought about a higher stimulatory effect as compared to VV1_CS. Taken together, these results indicate that both recombinant 14-3-3 proteins from *V. vinifera* were able to associate with and to stimulate the H^+ -ATPase, thus demonstrating that they were functionally active. Activities of both isoforms were comparable to values reported for other plant isoforms (VISCONTI *et al.* 2008).

VV1_CS and VV2_CS gene expression profiles: RT-PCR analysis was performed, in order to investigate the *VV1_CS* and *VV2_CS* gene expression profiles in berries at different developmental stages as well as in other grape organs.

Results demonstrated that *VV1_CS* and *VV2_CS*, besides berries, were expressed also in stems, roots and leaves (Fig. 4a). Expression studies performed with grape berries at different stages of development showed that the two transcripts were detected in all developmental stages but with different expression patterns (Fig. 4b). In fact, *VV1_CS* transcript increased constantly during the green stages (stages from 27 to 33) and at ripening (stage 35) and dramatically decreased during later maturation stages (stages from 36 to 39). *VV2_CS* transcript steadily increased during green stages, reaching a maximum at post-ripening (stage 36) and remaining constant until stage 38, to decrease only at the last maturation stage.

VV1_CS and VV2_CS expression in response to elicitors: To investigate the potential involvement of the two *Vitis* 14-3-3s in the response to stress, their transcript accumulation was evaluated after treatment of grape leaves with elicitors mimicking different biotic stresses. Jasmonic acid, ethephon (an ethylene-releasing compound), salicylic acid, hydrogen peroxide and ergosterol were supplied to grape leaves and *VV1_CS* and




Fig. 4: Semiquantitative RT-PCR analysis of 14-3-3 gene expression in grape roots, stems, leaves and in berries. **a)**: Roots, stems and leaves were collected from 2-month-old rooted plants. **b)**: Grape berries at different stages of development (modified “E-L system” according to COOMBE 1995). The grey bar (stage 35) represents the central stage of “véraison”.

VV2_CS transcripts levels determined by semi-quantitative PCR analyses at different times, up to 24 h. As shown in Fig. 5, all elicitors were able to trigger accumulation of both 14-3-3 transcripts, although at different extent and with different time-courses. The highest level of transcript accumulation was brought about by ergosterol elicitation.

Discussion

14-3-3 proteins belong to a highly conserved protein family with regulatory roles in all eukaryotes, and peculiar functions in plants. 14-3-3 family is represented in different animal or plant species by a variable number of isoforms. *In silico* analysis separates isoforms into two main groups (the epsilon group and the non-epsilon group) and in more subgroups or clusters. It was proposed that the animal epsilon 14-3-3 is the most similar to plant 14-3-3s (WANG and SHAKES 1996). This led to hypothesize that the epsilon isoform is a “living fossil” that has faithfully maintained the character of an ancestral 14-3-3 isoform (WANG and SHAKES 1996, DOOLITTLE 1994).

Both the two 14-3-3 isoforms identified from *Vitis* belong to the non-epsilon group, but they are sufficiently different to fall into two distinct clusters, omega and kappa respectively (Fig. 1b), their divergences being mostly located at the C-terminal regions. In particular, a part of the VV1_CS C-terminus is similar to that of *Arabidopsis* omega isoform, which has been proposed to contain a tenth alpha-helix (SHEN *et al.* 2003).

In order to verify whether the two novel sequences represent functional proteins, the ability of recombinant isoforms to bind to and stimulate the activity of the plas-




Fig. 5: Semiquantitative RT-PCR analysis of 14-3-3 gene expression in grapevine leaves treated with elicitors. Relative *VV1_CS* and *VV2_CS* levels were compared at 2, 4, 8 and 24 h to the inoculated control (0 h) and normalized to the reference gene *EF1- γ* .

ma membrane H⁺-ATPase was assayed. *In vitro* produced 14-3-3s showed a size (about 30 kDa) comparable to that of known 14-3-3s, while activity assays showed that both *Vitis* isoforms were able to bind to and to stimulate the phosphohydrolytic activity of the plasma H⁺-ATPase, thus demonstrating that the recombinant proteins are fully functional (XU *et al.* 2012).

We also investigated the *VV1_CS* and *VV2_CS* expression in different grapevine tissues. Results indicated that *VV1_CS* and *VV2_CS* transcripts were present in roots, stems, leaves and berries, where they exhibited differential expression profiles during fruit development and ripening. In particular, analysis in berries clearly showed that *VV1_CS* and *VV2_CS* transcripts strongly increased at *véraison* (Fig. 4b), a crucial stage in the berry development, characterized by dramatic metabolic and anatomical changes (COOMBE and MCCARTHY 2000, ZHANG *et al.* 2008, TERRIER *et al.* 2001). These results strongly suggest that these 14-3-3 isoforms play a relevant role in grapevine berry ripening and corroborate the hypothesis that 14-3-3 proteins may have a general role in the regulation of plant development (DAUGHERTY *et al.* 1996, LI and DHAUBHADEL 2011). 14-3-3 proteins are involved in plant responses to biotic stresses, such as powdery mildew infection (YANG *et al.*, 2009) or to abiotic stresses (YAO *et al.* 2007). In rice it was observed that 14-3-3s are elicited by both biotic and abiotic stresses (CHEN *et al.* 2006). In the present work, compounds functioning as biotic stress elicitors have been used to investigate the potential involvement of the two novel isoforms in the response of *Vitis* leaves to biotic stress. Jasmonic acid and ethylene are molecules regulating the induced systemic resistance, which is associated to plant/bacteria interactions (CHOUHARY *et al.* 2007). Salicylic acid and hydrogen peroxide are molecules regulating the systemic acquired resistance (HAMMERSCHMIDT 2009), which is involved in the plant response to pathogens. Ergosterol, a molecule from fungal membranes, is perceived as a signal of pathogen invasion (GRANADO *et al.* 1995) and it has been proposed as a general elicitor in the pathogen associated molecular patterns (PAMPs)-triggered immunity (ROSSARD *et al.* 2010). All the tested elicitors were able to rapidly increase *Vitis* 14-3-3 transcript expression up to 24 h, thereby indicating that 14-3-3 proteins are involved in the early events of the plant defence response to pathogens (JONES and DANGL 2006). Accordingly, jasmonic acid affected transcription of 14-3-3 genes in rice (CHEN *et al.* 2006), in hybrid poplar (LAPOINTE *et al.* 2001a) and in white spruce (LAPOINTE *et al.* 2001b). Jasmonate and ethephon induced the transcription of the 14-3-3 *Hb14-3-3c* in *Hevea brasiliensis* (YANG *et al.* 2012); salicylic acid caused an increase of 14-3-3 expression in potato (AKSAMIT *et al.* 2005); H₂O₂ induced 14-3-3 gene expression in rice (CHEN *et al.* 2006, YAO *et al.* 2007) while in tomato the transcription of TFT6 14-3-3 isoform coincided with peaks of ethylene and salicylic acid production (ROBERTS *et al.* 2002). As far as ergosterol, to our knowledge, this is the first evidence of 14-3-3 transcript elicitation by this compound.

Conclusions

In conclusion, this study provides the identification and characterization of two 14-3-3 proteins from *Vitis vinifera*. Information about 14-3-3 from tree species is lacking and grapevine is a worldwide economically relevant tree cultivated species. Results allowed to ascertain that the two 14-3-3 isoforms are fully functional proteins, expressed in

different organs and that in berries they exhibit a differential pattern of expression during development and ripening. Furthermore, it has been demonstrated that in leaves 14-3-3 gene transcription is elicited by different compounds mimicking biotic stresses. These results suggest that also in grapevine 14-3-3s accomplish fundamental functions, both in plant development and adaptation, thereby prompting further studies aimed to a more detailed characterization of 14-3-3 expression profiles and biochemical functions in *Vitis vinifera*, particularly in relation to berry ripening and quality.

Acknowledgements

This work was supported by the Italian Ministry of Education, University and Research (MIUR), in the frame of PRIN 2007 (M.M.) and by INRA/ISVV Bordeaux, France. We thank M. G. PALMGREN for the gift of the yeast strain expressing the AHA1 H⁺-ATPase. G. FARACE would like to thank, in memory, Prof. SAÏD HAMDI for having warmly accepted him in Bordeaux and for constant encouragement and fruitful suggestions during his stay. And, last but not least, thanks to Prof. S. DELROT for the constructive criticism on the final version of this manuscript.

References

- ADUCCI, P.; CAMONI, L.; MARRA, M.; VISCONTI, S.; 2002: From cytosol to organelles: 14-3-3 proteins as multifunctional regulators of plant cell. *IUBMB Life* **53**, 49-55.
- AKSAMIT, A.; KOROBCZAK, A.; SKALA, J.; LUKASZEWICZ, M.; SZOPA, J.; 2005: The 14-3-3 gene expression specificity in response to stress is promoter-dependent. *Plant Cell Physiol.* **46**, 1635-1645.
- BOSS, P. K.; DAVIES, C.; ROBINSON, S. P.; 1996: Analysis of the expression of anthocyanin pathway genes in developing *Vitis vinifera* L. cv. Shiraz grape berries and the implications for pathway regulation. *Plant Physiol.* **111**, 1059-1066.
- CAMONI, L.; DI LUCENTE, C.; PALLUCCA, R.; VISCONTI, S.; ADUCCI, P.; 2012: Binding of phosphatidic acid to 14-3-3 proteins hampers their ability to activate the plant plasma membrane H⁺-ATPase. *IUBMB Life* **64**, 710-716.
- CAMONI, L.; VISCONTI, S.; MARRA, M.; ADUCCI, P.; 2001: Adenosine 5'-monophosphate inhibits the association of 14-3-3 proteins with the plant plasma membrane H⁺-ATPase. *J. Biol. Chem.* **276**, 31709-31712.
- CHEN, F.; LI, Q.; SUN, L.; HE, Z.; 2006: The rice 14-3-3 gene family and its involvement in responses to biotic and abiotic stress. *DNA Res.* **13**, 53-63.
- COOMBE, B.; 1995: Growth stages of the grapevine: adoption of a system for identifying grapevine growth stages. *Aust. J. Grape Wine Res.* **1**, 104-110.
- COOMBE, B. G.; MCCARTHY, M. G.; 2000: Dynamics of grape berry growth and physiology of ripening. *Aust. J. Grape Wine Res.* **6**, 131-135.
- CUTLER, S. R.; RODRIGUEZ, P. L.; FINKELSTEIN, R. R.; ABRAMS, S. R.; 2010: Abscisic acid: emergence of a core signaling network. *Annu. Rev. Plant Biol.* **61**, 651-679.
- DAUGHERTY, C. J.; ROONEY, M. F.; MILLER, P. W.; FERL, R. J.; 1996: Molecular organization and tissue-specific expression of an Arabidopsis 14-3-3 gene. *Plant Cell* **8**, 1239-1248.
- DEL VISO, F.; CASARETTO, J. A.; QUATRANO, R. S.; 2007: 14-3-3 Proteins are components of the transcription complex of the ATEM1 promoter in Arabidopsis. *Planta* **227**, 167-175.
- DELILLE, J. M.; SEHNKE, P. C.; FERL, R. J.; 2001: The Arabidopsis 14-3-3 family of signaling regulators. *Plant Physiol.* **126**, 35-38.
- DENISON, F. C.; PAUL, A. L.; ZUPANSKA, A. K.; FERL, R. J.; 2011: 14-3-3 proteins in plant physiology. *Semin Cell Dev. Biol.* **22**, 720-727.
- DOOLITTLE, R. F.; 1994: Convergent evolution: the need to be explicit. *Trends Biochem. Sci.* **19**, 15-18.
- FERL, R. J.; LU, G.; BOWEN, B. W.; 1994: Evolutionary implications of the family of 14-3-3 brain protein homologs in Arabidopsis thaliana. *Genetica* **92**, 129-138.
- FULGOSI, H.; SOLL, J.; DE FARIA MARASCHIN, S.; KORTHOUT, H. A.; WANG, M.; TESTERINK, C.; 2002: 14-3-3 proteins and plant development. *Plant Mol. Biol.* **50**, 1019-1029.
- FULLONE, M. R.; VISCONTI, S.; MARRA, M.; FOGLIANO, V.; ADUCCI, P.; 1998: Fusicoccin effect on the in vitro interaction between plant 14-3-3 proteins and plasma membrane H⁺-ATPase. *J. Biol. Chem.* **273**, 7698-7702.
- GAMPALA, S. S.; KIM, T. W.; HE, J. X.; TANG, W.; DENG, Z.; BAI, M. Y.; GUAN, S.; LALONDE, S.; SUN, Y.; GENDRON, J. M.; CHEN, H.; SHIBAGAKI, N.; FERL, R. J.; EHRHARDT, D.; CHONG, K.; BURLINGAME, A. L.; WANG, Z. Y.; 2007: An essential role for 14-3-3 proteins in brassinosteroid signal transduction in Arabidopsis. *Dev. Cell* **13**, 177-189.
- GRANDO, J.; FELIX, G.; BOLLER, T.; 1995: Perception of Fungal Sterols in Plants (Subnanomolar Concentrations of Ergosterol Elicit Extracellular Alkalinization in Tomato Cells). *Plant Physiol.* **107**, 485-490.
- HONG, J. Y.; CHAE, M. J.; LEE, I. S.; LEE, Y. N.; NAM, M. H.; KIM, D. Y.; BYUN, M. O.; YOON, I. S.; 2011: Phosphorylation-mediated regulation of a rice ABA responsive element binding factor. *Phytochemistry* **72**, 27-36.
- ISHIDA, S.; FUKAZAWA, J.; YUASA, T.; TAKAHASHI, Y.; 2004: Involvement of 14-3-3 signaling protein binding in the functional regulation of the transcriptional activator repression of shoot growth by gibberellins. *Plant Cell* **16**, 2641-2651.
- JAILLON, O.; AURY, J. M.; NOEL, B.; POLICRITI, A.; CLEPET, C.; CASAGRANDE, A.; CHOISNE, N.; AUBOURG, S.; VITULO, N.; JUBIN, C.; VEZZI, A.; LEGEAI, F.; HUGUENEY, P.; DASILVA, C.; HORNER, D.; MICA, E.; JUBLOT, D.; POULAIN, J.; BRUYERE, C.; BILLAULT, A.; SEGURENS, B.; GOUYVENOUX, M.; UGARTE, E.; CATTONARO, F.; ANTHOUARD, V.; VICO, V.; DEL FABBRO, C.; ALAUX, M.; DI GASPERO, G.; DUMAS, V.; FELICE, N.; PAILLARD, S.; JUMAN, I.; MOROLDO, M.; SCALABRIN, S.; CANAGUIER, A.; LE CLAINCHE, I.; MALACRIDA, G.; DURAND, E.; PESOLE, G.; LAUCOU, V.; CHATELET, P.; MERDINOGLU, D.; DELLEDONNE, M.; PEZZOTTI, M.; LECHARNY, A.; SARPELLI, C.; ARTIGUENAVE, F.; PE, M. E.; VALLE, G.; MORGANTE, M.; CABOCHE, M.; ADAM-BLONDON, A. F.; WEISSENBACH, J.; QUETIER, F.; WINCKER, P.; 2007: The grapevine genome sequence suggests ancestral hexaploidization in major angiosperm phyla. *Nature* **449**, 463-467.
- JONES, J. D.; DANGL, J. L.; 2006: The plant immune system. *Nature* **444**, 323-329.
- KIM, T. W.; GUAN, S.; SUN, Y.; DENG, Z.; TANG, W.; SHANG, J. X.; BURLINGAME, A. L.; WANG, Z. Y.; 2009: Brassinosteroid signal transduction from cell-surface receptor kinases to nuclear transcription factors. *Nat. Cell Biol.* **11**, 1254-1260.
- KONAGAYA, K. I.; MATSUSHITA, Y.; KASAHARA, M.; NYUNOYA, H.; 2004: Members of 14-3-3 protein isoforms interacting with the resistance gene product N and the elicitor of *Tobacco mosaic virus*. *J. General Plant Pathol.* **70**, 221-231.
- LAEMMLI, U. K.; 1970: Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680-685.
- LAPOINTE, G.; LUCKEVICH, M. D.; CLOUTIER, M.; SEGUIN, A.; 2001a: 14-3-3 gene family in hybrid poplar and its involvement in tree defence against pathogens. *J. Exp. Bot.* **52**, 1331-1338.
- LAPOINTE, G.; LUCKEVICH, M. D.; SÉGUIN, A.; 2001b: Investigation on the induction of 14-3-3 in white spruce. *Plant Cell Rep.* **20**, 79-84.
- LI, X.; DHAUBHADEL, S.; 2011: Soybean 14-3-3 gene family: identification and molecular characterization. *Planta* **233**, 569-582.
- LIMA, M. M.; FERRERES, F.; DIAS, A. P.; 2012: Response of *Vitis vinifera* cell cultures to *Phaeoemoniella chlamydospora*: changes in phenolic production, oxidative state and expression of defence-related genes. *Eur. J. Plant Pathol.* **132**, 133-146.
- MARCHIVE, C.; MZID, R.; DELUC, L.; BARRIEU, F.; PIRRELLO, J.; GAUTHIER, A.; CORIO-COSTET, M. F.; REGAD, F.; CAILLETEAU, B.; HAMDI, S.; LAUVERGEAT, V.; 2007: Isolation and characterization of a *Vitis vinifera* transcription factor, VvWRKY1, and its effect on responses to fungal pathogens in transgenic tobacco plants. *J. Exp. Bot.* **58**, 1999-2010.
- MARRA, M.; OLIVARI, C.; VISCONTI, S.; ALBUMI, C.; ADUCCI, P.; DE MICHELIS, M. I.; 2000: A phosphopeptide corresponding to the cytosolic Stretch connecting transmembrane segments 8 and 9 of the plasma

- membrane H⁺-ATPase binds 14-3-3 proteins and inhibits fusicocin-induced activation of the H⁺-ATPase. *Plant Biol.* 1-6.
- MICHELET, B.; BOUTRY, M.; 1995: The Plasma Membrane H⁺-ATPase (A Highly Regulated Enzyme with Multiple Physiological Functions). *Plant Physiol.* **108**, 1-6.
- PIOTROWSKI, M.; OECKING, C.; 1998: Five new 14-3-3 isoforms from *Nicotiana tabacum* L.: implications for the phylogeny of plant 14-3-3 proteins. *Planta* **204**, 127-130.
- REGENBERG, B.; VILLALBA, J. M.; LANFERMEIER, F. C.; PALMGREN, M. G.; 1995: C-terminal deletion analysis of plant plasma membrane H⁺-ATPase: yeast as a model system for solute transport across the plant plasma membrane. *Plant Cell* **7**, 1655-1666.
- REID, E. R.; OLSSON, N.; SCHLOSSER, J.; PENG, F.; LUND, S. T.; 2006: An optimized grapevine RNA isolation procedure and statistical determination of reference genes for real-time RT-PCR during berry development. *Plant Biol.* **6**, 27.
- ROBERTS, M. R.; SALINAS, J.; COLLINGE, D. B.; 2002: 14-3-3 proteins and the response to abiotic and biotic stress. *Plant Mol. Biol.* **50**, 1031-1039.
- ROSSARD, S.; ROBLIN, G.; ATANASSOVA, R.; 2010: Ergosterol triggers characteristic elicitation steps in *Beta vulgaris* leaf tissues. *J. Exp. Bot.* **61**, 1807-1816.
- ROST, B.; SANDER, C.; 1993: Prediction of protein secondary structure at better than 70 % accuracy. *J. Mol. Biol.* **232**, 584-599.
- SCHOONHEIM, P. J.; SINNIGE, M. P.; CASARETTO, J. A.; VEIGA, H.; BUNNEY, T. D.; QUATRANO, R. S.; DE BOER, A. H.; 2007: 14-3-3 adaptor proteins are intermediates in ABA signal transduction during barley seed germination. *Plant J.* **49**, 289-301.
- SCHWECHHEIMER, C.; WILLIGE, B. C.; 2009: Shedding light on gibberellic acid signalling. *Curr. Opin. Plant Biol.* **12**, 57-62.
- SERRANO, R.; 1989: H⁺-ATPase from plasma membranes of *Saccharomyces cerevisiae* and *Avena sativa* roots: purification and reconstitution. *Methods Enzymol.* **157**, 533-544.
- SHEN, W.; CLARK, A. C.; HUBER, S. C.; 2003: The C-terminal tail of Arabidopsis 14-3-3omega functions as an autoinhibitor and may contain a tenth alpha-helix. *Plant J.* **34**, 473-484.
- SUN, G.; XIE, F.; ZHANG, B.; 2011: Transcriptome-wide identification and stress properties of the 14-3-3 gene family in cotton (*Gossypium hirsutum* L.). *Funct. Integr. Genomics* **11**, 627-636.
- TERRIER, N.; SAUVAGE, F. X.; AGEORGES, A.; ROMIEU, C.; 2001: Changes in acidity and in proton transport at the tonoplast of grape berries during development. *Planta* **213**, 20-28.
- VISCONTI, S.; CAMONI, L.; FULLONE, M. R.; LALLE, M.; MARRA, M.; ADUCCI, P.; 2003: Mutational analysis of the interaction between 14-3-3 proteins and plant plasma membrane H⁺-ATPase. *J. Biol. Chem.* **278**, 8172-8178.
- VISCONTI, S.; CAMONI, L.; MARRA, M.; ADUCCI, P.; 2008: Role of the 14-3-3 C-terminal region in the interaction with the plasma membrane H⁺-ATPase. *Plant Cell Physiol.* **49**, 1887-1897.
- WANG, W.; SHAKES, D. C.; 1996: Molecular evolution of the 14-3-3 protein family. *J. Mol. Evol.* **43**, 384-398.
- XU, W.; SHI, W.; JIA, L.; LIANG, J.; ZHANG, J.; 2012: TFT6 and TFT7, two different members of tomato 14-3-3 gene family, play distinct roles in plant adaption to low phosphorus stress. *Plant Cell Environ.* **35**, 1393-1406.
- YAFFE, M. B.; 2002: How do 14-3-3 proteins work?-- Gatekeeper phosphorylation and the molecular anvil hypothesis. *FEBS Lett.* **513**, 53-57.
- YANG, X.; WANG, W.; COLEMAN, M.; ORGIL, U.; FENG, J.; MA, X.; FERL, R.; TURNER, J. G.; XIAO, S.; 2009: Arabidopsis 14-3-3 lambda is a positive regulator of RPW8-mediated disease resistance. *Plant J.* **60**, 539-550.
- YANG, Z. P.; LI, H. L.; GUO, D.; TIAN, W. M.; PENG, S. Q.; 2012: Molecular characterization of a novel 14-3-3 protein gene (Hb14-3-3c) from Hevea brasiliensis. *Mol. Biol. Rep.* **39**, 4491-4497.
- YAO, Y.; DU, Y.; JIANG, L.; LIU, J. Y.; 2007: Molecular analysis and expression patterns of the 14-3-3 gene family from *Oryza sativa*. *J. Biochem. Mol. Biol.* **40**, 349-357.
- ZHANG, J.; MA, H.; FENG, J.; ZENG, L.; WANG, Z.; CHENG, S.; 2008: Grape berry plasma membrane proteome analysis and its differential expression during ripening. *J. Exp. Bot.* **59**, 2979-2990.
- ZHANG, Z. T.; ZHOU, Y.; LI, Y.; SHAO, S. Q.; LI, B. Y.; SHI, H. Y.; LI, X. B.; 2010: Interactome analysis of the six cotton 14-3-3s that are preferentially expressed in fibres and involved in cell elongation. *J. Exp. Bot.* **61**, 3331-3344.

Received January 23, 2014